HARBOR PORPOISE (Phocoena phocoena): Morro Bay Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

In the Pacific, harbor porpoise are found in coastal and inland waters from Point Conception, California to Alaska and across to Kamchatka and Japan (Gaskin 1984). Harbor porpoise appear to have more restricted movements along the western coast of the continental U.S. than along the eastern coast. Regional differences in pollutant residues in harbor porpoise indicate that they do not move extensively California, Oregon, and Washington between (Calambokidis and Barlow 1991). That study also showed regional differences within California (although the sample size was small). This pattern stands as a sharp contrast to the eastern coast of the U.S. and Canada where harbor porpoise are believed to migrate seasonally from as far south as the Carolinas to the Gulf of Maine and Bay of Fundy (Polacheck et al. 1995). A phylogeographic analysis of genetic data from northeast Pacific harbor porpoise did not show complete concordance between DNA sequence types and geographic location (Rosel 1992). However, an analysis of molecular variance (AMOVA) of the same data with additional samples found significant genetic differences for four of the six pair-wise comparisons between the four areas investigated: California, Washington, British Columbia, and Alaska (Rosel et al. 1995). These results demonstrate that harbor porpoise along the west coast of North America are not panmictic or migratory, and movement is sufficiently restricted




Figure 1. Stock boundaries and distributional range of harbor porpoise along the California and southern Oregon coasts. Dashed line represents harbor porpoise habitat (0-200 m) in this region.

that genetic differences have evolved. Subsequent genetic analyses of samples ranging from Morro Bay, California to British Columbia indicate that there is small-scale subdivision within the U.S. portion of this range (Chivers *et al.*, 2002, 2007; Morin *et al.*, 2021). Six harbor porpoise stocks have been designated off California/Oregon/Washington, based on genetic analyses and density discontinuities identified from aerial surveys. The stock boundaries in waters off California and southern Oregon are shown in Figure 1. For the Marine Mammal Protection Act (MMPA) Stock Assessment Reports, Pacific coast harbor porpoise stocks include: 1) the Morro Bay stock (this report), 2) the Monterey Bay stock, 3) the San Francisco-Russian River stock, 4) the northern California/southern Oregon stock, 5) the northern Oregon/Washington coast stock, and 6) the Inland Washington stock. Three additional Alaskan harbor porpoise stocks are reported separately in the Alaska Stock Assessment Reports.

POPULATION SIZE

Previous estimates of abundance for California harbor porpoise were based on aerial surveys conducted between the coast and the 50-fm isobath during 1988-95 (Barlow and Forney 1994, Forney 1999). These estimates did not include an unknown number of animals found in deeper waters. Barlow (1988) found that the vast majority of harbor porpoise in California were within the 0-50-fm depth range; however, Green et al. (1992) found that 24% of harbor porpoise seen during aerial surveys of Oregon and Washington were between the 100m and 200m isobaths (55 to 109 fathoms). A systematic ship survey of depth strata out to 90 m in northern California showed that porpoise abundance declined significantly in waters deeper than 60 m (Carretta et al. 2001). Since 1999, aerial surveys have extended farther offshore (to the 200 m depth contour or a minimum of 10 nmi from shore in the region of the Morro Bay stock) to provide a more complete abundance estimate (Forney et al. 2014). A recent analysis of long-term trends in the Morro Bay harbor porpoise population (Forney et al. 2020) between 1986 and 2012 estimated a population size of 4,191

(CV=0.561) porpoises during 2012. This estimate includes a correction factor of 3.42 (1/g(0); g(0)=0.292, CV=0.366) (Laake *et al.* 1997) to adjust for groups missed by aerial observers, and it is the most recent estimate available for this stock.

Minimum Population Estimate

The minimum population estimate for the Morro Bay harbor porpoise stock is taken as the lower 20th percentile of the lognormal distribution of the abundance estimated from the 2012 aerial surveys, or 2,698 animals.

Current Population Trend

A hierarchical Bayesian analysis of harbor porpoise trends between 1986 and 2012 (Forney et al. 2020) showed a marked increase in population size after 1991, when gillnet bycatch was largely eliminated within the range of the Morro Bay stock (Figure 2). This study also concludes that unmonitored harbor porpoise bycatch extending back as far as the 1950s likely decimated this population to a greater extent than previously understood.




Figure 2. Population trends for the Morro Bay harbor porpoise stock, 1986-2012 (from Forney *et al.* 2020). Estimates represent median abundance (with 95% credible intervals) for years with survey effort (solid symbols) and without survey effort (open symbols). Shaded bars along the x-axis reflect the relative level of gillnet bycatch: high (black), low to moderate (dark gray), or none (light gray).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Based on what are argued to be biological limits of the species (i.e. females give birth first at age 4 and produce one calf per year until death), the theoretical, maximum-conceivable growth rate of a closed harbor porpoise population was estimated as 9.4% per year based on a human survivorship curve (Barlow and Boveng 1991). This is very similar to the growth rate of 9.6% per year (95% credible interval: 6.2% - 13.0%) estimated by Forney *et al.* (2020) for the Morro Bay harbor porpoise stock between 1991 and 2012, based on long-term aerial surveys. This estimated growth rate can be considered a maximum net productivity rate, because this stock was estimated to include only 571 porpoises when gillnet bycatch was reduced to low levels in 1991, and by 2012 the population had increased to an estimated 4,191 individuals.

POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) level for this stock is calculated as the minimum population size (2,698) <u>times</u> one half the estimated maximum net growth rate for this stock of harbor porpoise (½ of 9.6%) <u>times</u> a recovery factor of 0.5 (for a stock of unknown status; Wade and Angliss 1997), resulting in a PBR of 65.

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY Fishery Information

Gillnet fisheries for halibut and white seabass that historically operated in the vicinity of Morro Bay were eliminated in this stock's range in 2002 by a ban on gillnets inshore of 60 fathoms (~110 m) from Point Arguello to Point Reyes, California. The large-mesh drift gillnet fishery for swordfish and thresher shark operates too far offshore to interact with harbor porpoise in this region. In the most recent five-year period (2015-2019), no fishery-related strandings of harbor porpoise were documented within this stock's range (Carretta *et al.* 2021).

Table 1. Summary of available data on incidental mortality and serious injury of Morro Bay Stock harbor porpoise in commercial fisheries that might take this species. Mean annual takes are based on 2015-2019 data, Carretta *et al.* (2021). n/a indicates that data are not available.

Fishery Name	Year(s)	Data Type	Percent Observer Coverage	Observed Mortality	Kill/Day	Estimated Mortality (CV in parentheses)	Mean Annual Takes (CV in parentheses)
Unknown fishery	2015-2019	Stranding	n/a	none	n/a	n/a	0 (n/a)
Minimum total annual takes							0 (n/a)

STATUS OF STOCK

Harbor porpoise in California are not listed as threatened or endangered under the Endangered Species Act nor as depleted under the Marine Mammal Protection Act. Barlow and Hanan (1995) calculate the status of harbor porpoise relative to historic carrying capacity (K) using a technique called backprojection. They calculate that the central California population (including Morro Bay, Monterey Bay, and San Francisco-Russian River stocks) could have been reduced to between 30% and 97% of K by incidental fishing mortality, depending on the choice of input parameters. They conclude that there is no practical way to reduce the range of this estimate. Although Forney *et al.* 2020 documented a marked increase in the Morro Bay harbor porpoise stock, the carrying capacity of this stock is not known and the population status relative to Optimum Sustainable Population (OSP) levels must be treated as unknown.

Because the known human-caused mortality or serious injury (zero harbor porpoise per year) is less than the PBR (65), this stock is not considered a "strategic" stock under the MMPA, and fishery mortality can be considered insignificant and approaching zero mortality and serious injury rate. Harbor porpoises are sensitive to disturbance by a variety of anthropogenic sound sources, and the limited range of several U.S. West Coast harbor porpoise stocks makes them particularly vulnerable to potential impacts (see overview in Forney *et al.* 2017). A recent habitat concern along the U.S. West coast includes the use of acoustic deterrent devices ('seal bombs') that are used in commercial fishing activities off California (Simonis *et al.* 2020), especially in the Monterey Bay region.

REFERENCES

- Barlow, J. 1988. Harbor porpoise (*Phocoena phocoena*) abundance estimation in California, Oregon and Washington: I. Ship surveys. Fish. Bull. 86:417-432.
- Barlow, J. and P. Boveng. 1991. Modeling age-specific mortality for marine mammal populations. Mar. Mamm. Sci. 7(1):84-119.
- Barlow, J. and K. A. Forney. 1994. An assessment of the 1994 status of harbor porpoise in California. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-205. 17 pp.
- Barlow, J. and D. Hanan. 1995. An assessment of the status of harbor porpoise in central California. Rept. Int. Whal., Special Issue 16:123-140.
- Calambokidis, J. and J. Barlow. 1991. Chlorinated hydrocarbon concentrations and their use for describing population discreteness in harbor porpoises from Washington, Oregon, and California. pp. 101-110 In: J. E. Reynolds III and D. K. Odell (eds.) Marine mammal strandings in the United States. NOAA Tech. Rep. NMFS 98.
- Carretta, J.V., B.L. Taylor, and S.J. Chivers. 2001. Abundance and depth distribution of harbor porpoise (*Phocoena phocoena*) in northern California determined from a 1995 ship survey. U.S. Fishery Bulletin 99:29-39.
- Carretta, J.V., J. Greenman, K. Wilkinson, J. Freed, L. Saez, D. Lawson, J. Viezbicke, and J. Jannot. 2021. Sources of human-related injury and mortality for U.S. Pacific West Coast Marine Mammal Stock Assessments, 2015-2019. NOAA Technical Memorandum NMFS-SWFSC-643. 157 pp.
- Chivers, S.J., A.E. Dizon, P.J. Gearin, and K.M. Robertson. 2002. Small-scale population structure of eastern North Pacific harbour porpoises, (*Phocoena phocoena*), indicated by molecular genetic analyses. Journal of Cetacean Research and Management 4(2):111-122.
- Chivers, S.J., B. Hanson, J. Laake, P. Gearin, M.M. Muto, J. Calambokidis, D. Duffield, T. McGuire, J. Hodder, D. Greig, E. Wheeler, J. Harvey, K.M. Robertson, and B. Hancock. 2007. Additional genetic evidence for population structure of *Phocoena phocoena* off the coasts of California, Oregon, and Washington. Southwest Fisheries Science Center Administrative Report LJ-07-08. 16pp.
- Forney, K.A. 1999. The abundance of California harbor porpoise estimated from 1993-97 aerial line-transect surveys. Admin. Rep. LJ-99-02. Southwest Fisheries Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 16 pp.

- Forney K.A., J.V. Carretta, S.R. Benson. 2014. Preliminary estimates of harbor porpoise abundance in Pacific Coast waters of California, Oregon and Washington, 2007-2012. U.S. Dep. Commer., NOAA Tech Memo NMFS-SWFSC-537. 21 p.
- Forney, K.A., Southall, B.L., Slooten, E., Dawson, S., Read, A.J., Baird, R.W. and Brownell Jr, R.L., 2017. Nowhere to go: noise impact assessments for marine mammal populations with high site fidelity. Endangered species research, 32, pp. 391-413.
- Forney, K.A., J.E. Moore, J. Barlow, J.V. Carretta and S.R. Benson. 2020. A multi-decadal Bayesian trend analysis of harbor porpoise (*Phocoena phocoena*) populations off California relative to past fishery bycatch. Marine Mammal Science 2020; 1–15. https://doi.org/10.1111/mms.12764 Gaskin, D. E. 1984. The harbour porpoise (*Phocoena phocoena* L.): regional populations, status, and information on direct and indirect catches. Rep. int. Whal. Commn 34:569 586.
- Green, G. A., J. J. Brueggeman, R. A. Grotefendt, C. E. Bowlby, M. L. Bonnell, and K. C. Balcomb, III.
 1992. Cetacean distribution and abundance off Oregon and Washington, 1989-1990. Ch. 1 In: J. J.
 Brueggeman (ed.). Oregon and Washington Marine Mammal and Seabird Surveys. Minerals
 Management Service Contract Report 14-12-0001-30426 prepared for the Pacific OCS Region.
- Laake, J. L., J. C. Calambokidis, S. D. Osmek, and D. J. Rugh. 1997. Probability of detecting harbor porpoise from aerial surveys: estimating g(0). J. Wildl. Manag. 61:63-75.
- Morin P.A., B.R. Forester, K.A. Forney, C.A. Crossman, B. Hancock-Hanser, K.M. Robertson, L.G. Barrett-Lennard, R.W. Baird, J. Calambokidis, P. Gearin, M.B. Hanson, C. Schumacher, T. Harkins, M. Fontaine, B.L Taylor, and K. Parsons. 2021. Population structure in a continuously distributed coastal marine species, the harbor porpoise, based on microhaplotypes derived from poor quality samples. Molecular Ecology. https://doi.org/10.1111/mec.15827.
- Norris, K.S. and J.H. Prescott. 1961. Observations on Pacific cetaceans of Californian and Mexican waters. Univ. Calif. Publ. Zool. 63(4):291-402.
- Polacheck, T., F. W. Wenzel, and G. Early. 1995. What do stranding data say about harbor porpoise (*Phocoena phocoena*). Rep. Int. Whal. Comm., Special Issue 16:169-179.
- Rosel, P. E. 1992. Genetic population structure and systematic relationships of some small cetaceans inferred from mitochondrial DNA sequence variation. Ph.D. Dissertation, Univ. Calif. San Diego. 191pp.
- Rosel, P. E., A. E. Dizon, and M. G. Haygood. 1995. Variability of the mitochondrial control region in populations of the harbour porpoise, *Phocoena phocoena*, on inter-oceanic and regional scales. Can. J. Fish. and Aquat. Sci. 52:1210-1219.
- Simonis, A.E., Forney, K.A., Rankin, S., Ryan, J., Zhang, Y., DeVogelaere, A., Joseph, J., Margolina, T., Krumpel, A. and Baumann-Pickering, S., 2020. Seal Bomb Noise as a Potential Threat to Monterey Bay Harbor Porpoise. Frontiers in Marine Science, 7, p.142.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Woodley, T. H. and A. J. Read. 1991. Potential rates of increase of a harbour porpoise (*Phocoena phocoena*) population subjected to incidental mortality in commercial fisheries. Can. J. Fish. Aquat. Sci. 48:2429-2435.

HARBOR PORPOISE (Phocoena phocoena): Monterey Bay Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

In the Pacific, harbor porpoise are found in coastal and inland waters from Point Conception, California to Alaska and across to Kamchatka and Japan (Gaskin 1984). Harbor porpoise appear to have more restricted movements along the western coast of the continental U.S. than along the eastern coast. Regional differences in pollutant residues in harbor porpoise indicate that they do not move extensively California, Oregon, and Washington between (Calambokidis and Barlow 1991). That study also showed regional differences within California (although the sample size was small). This pattern stands as a sharp contrast to the eastern coast of the U.S. and Canada where harbor porpoise are believed to migrate seasonally from as far south as the Carolinas to the Gulf of Maine and Bay of Fundy (Polacheck et al. 1995). A phylogeographic analysis of genetic data from northeast Pacific harbor porpoise did not show complete concordance between DNA sequence types and geographic location (Rosel 1992). However, an analysis of molecular variance (AMOVA) of the same data with additional samples found significant genetic differences for four of the six pair-wise comparisons between the four areas investigated: California, Washington, British Columbia, and Alaska (Rosel et al. 1995). These results demonstrate that harbor porpoise along the west coast of North America are not panmictic or migratory, and movement is sufficiently restricted that genetic differences have evolved. Subsequent genetic analyses of samples ranging from Morro Bay,




Figure 1. Stock boundaries and distributional range of harbor porpoise along the California/southern Oregon coast. Dashed line represents harbor porpoise habitat (0-200 m) along the U.S. west coast.

California to British Columbia indicate that there is small-scale subdivision within the U.S. portion of this range (Chivers *et al.*, 2002, 2007; Morin *et al.*, 2021). Six harbor porpoise stocks have been designated off California/Oregon/Washington, based on genetic analyses and density discontinuities identified from aerial surveys. The stock boundaries in waters off California and southern Oregon are shown in Figure 1. For the Marine Mammal Protection Act (MMPA) Stock Assessment Reports, Pacific coast harbor porpoise stocks include: 1) the Morro Bay stock, 2) the Monterey Bay stock (this report), 3) the San Francisco-Russian River stock, 4) the northern California/southern Oregon stock, 5) the northern Oregon/Washington coast stock, and 6) the Inland Washington stock. Three additional Alaskan harbor porpoise stocks are reported separately in the Alaska Stock Assessment Reports.

POPULATION SIZE

Previous estimates of abundance for California harbor porpoise were based on aerial surveys conducted between the coast and the 50-fm isobath during 1988-95 (Barlow and Forney 1994, Forney 1999). These estimates did not include an unknown number of animals found in deeper waters. Barlow (1988) found that the vast majority of harbor porpoise in California were within the 0-50-fm depth range; however, Green et al. (1992) found that 24% of harbor porpoise seen during aerial surveys of Oregon and Washington were between the 100m and 200m isobaths (55 to 109 fathoms). A systematic ship survey of depth strata out to 90 m in northern California showed that porpoise abundance declined significantly in waters deeper than 60 m (Carretta et al. 2001). Starting in 1999, aerial surveys extended farther offshore (to the 200m depth contour or a minimum of 15 nmi from shore in the region of the Monterey Bay stock) to provide a more complete abundance estimate (Forney et al. 2014). A recent analysis of long-term trends in the Monterey Bay harbor