

KILLER WHALE (*Orcinus orca*): Eastern North Pacific Southern Resident Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Killer whales occur in all oceans and seas (Leatherwood and Dahlheim 1978). Although they occur in tropical and offshore waters, killer whales prefer the colder waters of both hemispheres, with greatest abundances found within 800 km of major continents (Mitchell 1975, Forney and Wade 2006). Along the west coast of North America, killer whales occur along the entire Alaskan coast (Braham and Dahlheim 1982, Hamilton *et al.* 2009), in British Columbia and Washington inland waterways (Bigg *et al.* 1990), and along the outer coasts of Washington, Oregon and California (Hamilton *et al.* 2009). Seasonal and year-round occurrence is documented for killer whales throughout Alaska (Braham and Dahlheim 1982) and in the intracoastal waterways of British Columbia and Washington, where three ecotypes have been recognized: 'resident', 'transient' and 'offshore' (Bigg *et al.* 1990, Ford *et al.* 1994), based on aspects of morphology, ecology, genetics and behavior (Ford and Fisher 1982; Baird and Stacey 1988; Baird *et al.* 1992, Hoelzel *et al.* 1998, Morin *et al.* 2010, Ford *et al.* 2014). Genetic studies of killer whales globally suggest that residents and transient ecotypes warrant subspecies recognition (Morin *et al.* 2010) and each are currently listed as unnamed subspecies of *Orcinus orca* (Committee on Taxonomy 2018).

The range of southern resident killer whales is described in the draft biological report for the Proposed Revision of the Critical Habitat Designation for Southern Resident Killer Whales (NMFS 2021a, 2021b): "The three pods of the Southern Resident DPS, identified as J, K, and L pods, reside for part of the year in the inland waterways of Washington State and British Columbia known as the Salish Sea (Strait of Georgia, Strait of Juan de Fuca, and Puget Sound), principally during the late spring, summer, and fall (Ford *et al.* 2000, Krahn *et al.* 2002). The whales also visit outer coastal waters off Washington and Vancouver Island, especially in the area between Grays Harbor and the Columbia River (Ford *et al.* 2000, Hanson *et al.* 2017), but travel as far south as central California and as far north as the Southeast Alaska. Although less is known about the whales' movements in outer coastal waters, satellite tagging, opportunistic sighting, and acoustic recording data suggest that Southern Residents spend nearly all of their time on the continental shelf, within 34 km (21.1 mi) of shore in water less than 200 m (656.2 ft) deep (Hanson *et al.* 2017)." Details of their winter range from satellite-tagging reveal whales use the entire Salish Sea (northern end of the Strait of Georgia and Puget Sound) in addition to coastal waters from the central west coast of Vancouver Island, British Columbia to Pt. Reyes in northern California. Animals from J pod were documented moving between the northern Strait of Georgia and the western entrance of the Strait of Juan de Fuca, with limited movement into coastal waters. In contrast, K and L pod movements were characterized by a coastal distribution from the western entrance to the Strait of Juan de Fuca to Pt. Reyes California (Hanson *et al.* 2017). Of the three pods comprising this stock, one (J) is commonly sighted in inshore waters in winter, while the other two (K and L) apparently spend more time offshore (Ford *et al.* 2000). Krahn *et al.* (2009) described sample pollutant ratios from K and L pod whales that were consistent with a hypothesis of time spent foraging in California waters, which is consistent with sightings of K and L pods as far south as Monterey Bay. In June 2007, whales from L-pod were sighted off Chatham Strait, Alaska, the farthest north they have ever been documented (J. Ford, pers. comm.). Southern resident killer whale attendance in their core summer habitat in the Salish Sea appears to be declining, with occurrence well-below average since 2017 (Center for Whale Research 2019). Passive autonomous acoustic recorders have provided more information on the seasonal occurrence of these pods along the west coast of




Figure 1. Approximate April - October distribution of the Eastern North Pacific Southern Resident killer whale stock (shaded area) and range of sightings (diagonal lines).

the U.S. (Hanson *et al.* 2013). In addition, satellite-linked tags were deployed in winter months on members of J, K, and L pods. Results were consistent with previous data, but provided much greater detail, showing wide-ranging use of inland waters by J Pod whales and extensive movements in U.S. coastal waters by K and L Pods.

Based on data regarding association patterns, acoustics, movements, genetic differences and potential fishery interactions, eight killer whale stocks are recognized within the Pacific U.S. EEZ: 1) the Eastern North Pacific Alaska Resident stock - occurring from Southeast Alaska to the Bering Sea, 2) the Eastern North Pacific Northern Resident stock - occurring from British Columbia through Alaska, 3) the Eastern North Pacific Southern Resident stock - occurring mainly within the inland waters of Washington State and southern British Columbia but extending from central California into southern Southeast Alaska (see Fig. 1), 4) the West Coast Transient stock - occurring from Alaska through California, 5) the Gulf of Alaska, Aleutian Islands, and Bering Sea Transient stock - occurring from southeast Alaska to the Bering Sea, 6) the AT1 Stock – found only in Prince William Sound, 7) the Eastern North Pacific Offshore stock - occurring from Southeast Alaska through California, 8) the Hawaiian stock. The Stock Assessment Reports for the Alaska Region contain information concerning the Eastern North Pacific Alaska Resident, Eastern North Pacific Northern Resident and the Gulf of Alaska, Aleutian Islands, and Bering Sea, AT1, and Eastern North Pacific Transient stocks.

POPULATION SIZE

The Eastern North Pacific Southern Resident stock is a trans-boundary stock including killer whales in inland Washington and southern British Columbia waters. In 1993, the three pods comprising this stock totaled 96 killer whales (Ford *et al.* 1994). The population increased to 99 whales in 1995, then declined to 79 whales in 2001, and most recently numbered 72 whales in 2020 (Fig. 2; Ford *et al.* 2000; Center for Whale Research 2019). The 2001-2005 counts included a whale born in 1999 (L-98) that was listed as missing during the annual census in May and June 2001 but was subsequently discovered alone in an inlet off the west coast of Vancouver Island. L-98 remained separate from L pod until 10 March 2006 when he died due to injuries associated with a vessel interaction in Nootka Sound. L-98 has been subtracted from the official 2006 and subsequent population censuses. The most recent census spanning 1 July 2019 through 1 July 2020 includes the death of an adult male, but does not include the two calves that were born in fall 2020.

Minimum Population Estimate

The abundance estimate for this stock of killer whales is a direct count of individually identifiable animals. It is thought that the entire population is censused every year. This estimate therefore serves as both a best estimate of abundance and a minimum estimate of abundance. Thus, the minimum population estimate (N_{min}) for the Eastern North Pacific Southern Resident stock of killer whales is 72 animals.




Figure 2. Population of Eastern North Pacific Southern Resident stock of killer whales, 1974-2020. Each year's count includes animals first seen and first missed; a whale is considered first missed the year after it was last seen alive (Ford *et al.* 2000; Center for Whale Research 2020).

Current Population Trend

During the live-capture fishery that existed from 1967 to 1973, it is estimated that 47 killer whales, mostly immature, were taken out of this stock (Ford *et al.* 1994). Since the first complete census of this stock in 1974 when 71 animals were identified, the number of southern resident killer whales has fluctuated. Between 1974 and the mid-1990s, the Southern Resident stock increased approximately 35% (Ford *et al.* 1994), representing a net annual growth rate of 1.8% during those years. Following the peak census count of 99 animals in 1995, the population size has declined approximately 1% annually and currently stands at 72 animals as of the 2020 census (Ford *et al.* 2000; Center for Whale Research 2020).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

A reliable estimate of the maximum net productivity rate is currently unavailable for this stock of killer whales. Matkin *et al.* (2014) estimated a maximum population annual growth rate of 1.035 for southern Alaska resident killer whales. The authors noted that the 3.5% annual rate estimated for southern Alaska residents is higher than previously measured rates for British Columbia northern residents (2.9%, Olesiuk *et al.* 1990) and “probably represents a population at r_{\max} (maximum rate of growth).” In the absence of published estimates of R_{\max} for southern resident killer whales, the maximum annual rate of 3.5% found for southern Alaska residents is used for this stock of southern resident killer whales. This reflects more information about the known life history of resident killer whales than the default R_{\max} of 4% and results in a more conservative estimate of potential biological removal (PBR).

POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) level for this stock is calculated as the minimum population size (72) times one-half the maximum net growth rate for *Alaska* resident killer whales ($\frac{1}{2}$ of 3.5%) times a recovery factor of 0.1 (for an endangered stock, Wade and Angliss 1997), resulting in a PBR of 0.13 whales per year, or approximately 1 animal every 7 years.

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fisheries Information

Salmon drift gillnet fisheries in Washington inland waters were last observed in 1993 and 1994 and no killer whale entanglements were documented, though observer coverage levels were less than 10% (Erstad *et al.* 1996, Pierce *et al.* 1994, Pierce *et al.* 1996, NWIFC 1995). Fishing effort in the inland waters drift gillnet fishery has declined considerably since 1994 because far fewer vessels participate today. Past marine mammal entanglements in this fishery included harbor porpoise, Dall’s porpoise, and harbor seals. Coastal marine tribal set gillnets also occur along the outer Washington coast and no killer whale interactions have been reported in this fishery since the inception of the observer program in 1988, though the fishery is not active every year (Gearin *et al.* 1994, Gearin *et al.* 2000, Makah Fisheries Management). No fishery-related mortality from gillnet fisheries in California waters was documented between 2015-2019 (Carretta 2021, Carretta *et al.* 2021).

An additional source of information on killer whale mortality and injury incidental to commercial fishery operations is the self-reported fisheries information required of vessel operators by the MMPA. No self-report records of killer whale mortality have been reported.

Due to a lack of observer programs, there are few data concerning the mortality of marine mammals incidental to Canadian commercial fisheries. Since 1990, there have been no reported fishery-related strandings of killer whales in Canadian waters. However, in 1994 one killer whale was reported to have contacted a salmon gillnet but did not entangle (Guenther *et al.* 1995). In 2014 a northern resident killer whale became entangled in a gillnet, was released from the net, but died the next winter (Fisheries and Oceans Canada 2018). Data regarding the level of killer whale mortality related to commercial fisheries in Canadian waters are not available.

The known total fishery mortality and serious injury for the southern resident stock of killer whales is zero, but undetected mortality and serious injury may occur.

Other Mortality

In 2012, a moderately decomposed juvenile female southern resident killer whale (L-112) was found dead near Long Beach, WA. A full necropsy was performed and the cause of death was determined to be blunt force trauma to the head, however the source of the trauma (vessel strike, intraspecific aggression, or other unknown source) could not be established (NOAA 2014). There was documentation of a whale-boat collision in Haro Strait in 2005 which resulted in a minor injury to a whale. In 2006, whale L98 was killed during a vessel interaction. It is important to note that L98 had become habituated to regularly interacting with vessels during its isolation in Nootka Sound. In spring 2016, a young adult male, L95, was found to have died of a fungal infection related to a satellite tag deployment approximately 5 weeks prior to its death. The expert panel reviewing the stranding noted that “the tag loss, tag petal retention with biofilm formation or direct pathogen implantation, and development of a fungal infection at the tag site contributed to the illness, stranding, and death of this whale.” (NMFS 2016). In fall 2016 another young adult male, J34, was found dead in the northern Georgia Strait. The necropsy indicated that “the animal had injuries consistent with blunt trauma to the dorsal side, and a hematoma indicating that it was alive at the time of injury and would have survived the initial trauma for a period of time prior to death” (Fisheries and Oceans Canada 2019). The injuries are consistent with those incurred during a vessel strike. A recent summary of killer whale strandings in the northeastern Pacific Ocean and Hawaii noted the occurrence of human interactions across all age classes (Raverty *et al.* 2020).

Habitat Issues

A population viability analysis identified several risk factors to this population, including limitation of preferred Chinook salmon prey, anthropogenic noise and disturbance resulting in decreased foraging efficiency, and high levels of contaminants, including PCBs and DDT (Ebre 2002, Clark *et al.* 2009, Krahn *et al.* 2007, 2009, Lacy *et al.* 2017). The summer range of this population, the inland waters of Washington and British Columbia, are home to a large commercial whale watch industry, and high levels of recreational boating and commercial shipping. Potential for acoustic masking effects on the whales' communication and foraging due to vessel traffic remains a concern (Erbe 2002, Clark *et al.* 2009, Lacy *et al.* 2017). In 2011, vessel approach regulations were implemented to restrict vessels from approaching closer than 200m. A genetic study of diet of southern resident killer whales from fecal remains collected during 2006-2011 noted that salmonids accounted for >98.6% of genetic sequences (Ford *et al.* 2016). Of six salmonid species documented, Chinook salmon accounted for 79.5% of the sequences, followed by coho salmon (15%). Chinook salmon dominate the diet in early summer, with coho salmon averaging >40% of the diet in late summer. Sockeye salmon were also found to be occasionally important (>18% in some samples). Non-salmonids were rarely observed. These results are consistent with those obtained from surface prey remains, and confirm the importance of Chinook salmon in this population's diet. These authors also noted the absence of pink salmon in the fecal samples. Prior studies note the prevalence of Chinook salmon in the killer whale diet, despite the relatively low abundance of this species in the region, supporting the thesis that southern resident killer whales are Chinook salmon specialists (Ford and Ellis 2006, Hanson *et al.* 2010). Recent studies of diet in other seasons and regions of their range indicate that although Chinook represent a major component of their diet almost year-round, other species also make potentially important contributions, likely when Chinook are less available (Hanson *et al.* 2021). There is evidence that reduced abundance of Chinook salmon has negatively affected this population via reduced fecundity (Ayres *et al.* 2012, Ford *et al.* 2009, Ward *et al.* 2009, Wasser *et al.* 2017). In addition, the high trophic level and longevity of the population has predisposed them to accumulate high levels of contaminants that potentially impact health (Krahn *et al.* 2007, 2009). In particular, there is evidence of high levels of flame retardants in young animals (Krahn *et al.* 2007, 2009). High DDT/PCB ratios have been found in Southern Resident killer whales, especially in K and L pods (Krahn *et al.* 2007, NMFS 2019b), which spend more time in California waters where DDTs still persist in the marine ecosystem (Sericano *et al.* 2014).

STATUS OF STOCK

Total documented annual fishery mortality and serious injury for this stock from 2015-2019 (zero) is not known to exceed 10% of the calculated PBR (0.13). Given the low PBR level, a single undetected / undocumented fishery mortality or serious injury would exceed 10% of the PBR, thus it is unknown if fishery mortality and serious injury is approaching zero mortality and serious injury rate. The documented annual level of human-caused mortality and serious injury for the most-recent 5-year period includes the death of L95 (fungal infection related to a satellite-tag) and J34 (vessel strike), or 0.4 whales annually, which exceeds the PBR (0.13). Southern Resident killer whales were formally listed as "endangered" under the ESA in 2005 and consequently the stock is automatically considered as a "strategic" stock under the MMPA. This stock was considered "depleted" (68 FR 31980, May 29, 2003) prior to its 2005 listing under the ESA (70 FR 69903, November 18, 2005).

REFERENCES

- Angliss, R. P., and K. L. Lodge. 2002. Alaska marine mammal stock assessments, 2002. U.S. Dep. Commer., NOAA Technical Memorandum NMFS-AFSC-133. 224 pp.
- Ayres K.L., Booth R.K., Hempelmann J.A., Koski K.L., Emmons C.K., Baird R.W., et al. 2012. Distinguishing the Impacts of Inadequate Prey and Vessel Traffic on an Endangered Killer Whale (*Orcinus orca*) Population. PLoS ONE 7(6): e36842.
- Baird, R. W., and P. J. Stacey. 1988. Variation in saddle patch pigmentation in populations of killer whales (*Orcinus orca*) from British Columbia, Alaska, and Washington State. Canadian Journal of Zoology 66:2582-2585.
- Baird, R. W., P. A. Abrams, and L. M. Dill. 1992. Possible indirect interactions between transient and resident killer whales: implications for the evolution of foraging specializations in the genus *Orcinus*. Oecologia 89:125-132.
- Barlow, J. and K.A. Forney. 2007. Abundance and population density of cetaceans in the California Current ecosystem. Fishery Bulletin 105:509-526.
- Bigg, M. A., P. F. Olesiuk, G. M. Ellis, J. K. B. Ford, and K. C. Balcomb III. 1990. Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Pp. 386-406, *In*: Hammond, P. S., S. A. Mizroch, and G. P. Donovan (eds.), Individual Recognition of Cetaceans: Use of Photo-identification and Other Techniques to Estimate Population Parameters. Rep. Int. Whal. Commn. Special Issue 12.

- Braham, H. W., and M. E. Dahlheim. 1982. Killer whales in Alaska documented in the Platforms of Opportunity Program. Rep. Int. Whal. Commn. 32:643-646.
- Brault, S., and H. Caswell. 1993. Pod-specific demography of killer whales (*Orcinus orca*). Ecology 74(5):1444-1454.
- Carretta, J.V. 2021. Estimates of marine mammal, sea turtle, and seabird bycatch in the California large-mesh drift gillnet fishery: 1990-2019. NOAA Technical Memorandum NMFS-SWFSC-654.
- Carretta, J.V., J. Greenman, K. Wilkinson, J. Freed, L. Saez, D. Lawson, J. Viezbicke, and J. Jannot. 2021. Sources of human-related injury and mortality for U.S. Pacific West Coast Marine Mammal Stock Assessments, 2015-2019. NOAA Technical Memorandum NMFS-SWFSC-643. 157 pp.
- Center for Whale Research 2020. <https://www.whaleresearch.com/orca-population>
- Clark, C.W., Ellison, W.T., Southall, B.L., Hatch, L., Van Parijs, S.M., Frankel, A. and Ponirakis, D., 2009. Acoustic masking in marine ecosystems: intuitions, analysis, and implication. Marine Ecology Progress Series, 395, pp.201-222.
- Clarke, J.T., C.L. Christman, A.A. Brower, and M.C. Ferguson. 2013. Distribution and Relative Abundance of Marine Mammals in the Northeastern Chukchi and Western Beaufort Seas, 2012. Annual Report, OCS Study BOEM 2013-00117. National Marine Mammal Laboratory, Alaska Fisheries Science Center, NMFS, NOAA, 7600 Sand Point Way NE, F/AKC3, Seattle, WA 98115-6349.
- Dahlheim, M. E., D.K. Ellifrit, and J.D. Swenson. 1997. Killer whales of Southeast Alaska: a catalogue of photoidentified individuals. National Marine Mammal Laboratory, AFSC, NMFS, NOAA, 7600 Sand Point Way NE, Seattle, WA 98115. 79 pp.
- Erbe, C., 2002. Underwater noise of whale-watching boats and potential effects on killer whales (*Orcinus orca*), based on an acoustic impact model. Marine Mammal Science, 18(2), pp.394-418.
- Erstad, P., S.J. Jeffries, and D.J. Pierce. 1996. 1994 Report for the Puget Sound fishery observer program in management areas 10/11 & 12/12B: nontreaty chum gill net fishery. Final Report, Washington Dept. Fish and Wildlife, Olympia, WA. 14 pp.
- Fisheries and Oceans Canada. 2018. Recovery Strategy for the Northern and Southern Resident Killer Whales (*Orcinus orca*) in Canada. Species at Risk Act Recovery Strategy Series, Fisheries & Oceans Canada, Ottawa, x + 84 pp.
- Fisheries and Oceans Canada. 2019. Necropsy results: Southern Resident Killer Whale J34.
- Ford, J.K.B. Pacific Biological Station, Department of Fisheries and Oceans, Nanaimo, BC V9R 5K6.
- Ford, J.K.B., and H. D. Fisher. 1982. Killer whale (*Orcinus orca*) dialects as an indicator of stocks in British Columbia. Rep. Int. Whal. Commn. 32:671-679.
- Ford, J.K.B., G. M. Ellis, and K. C. Balcomb. 1994. Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington State. University of British Columbia Press, Vancouver, BC, and University of Washington Press, Seattle. 102 pp.
- Ford, J.K., Ellis, G.M., Barrett-Lennard, L.G., Morton, A.B., Palm, R.S. and Balcomb III, K.C., 1998. Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. Canadian Journal of Zoology, 76(8), pp.1456-1471.
- Ford, J.K.B., G. M. Ellis, and K. C. Balcomb. 2000. Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington. 2nd edition. University of British Columbia Press, Vancouver, BC, and University of Washington Press, Seattle. 104 pp.
- Ford, J.K.B., and G.M. Ellis. 2006. Selective foraging by fish-eating killer whales *Orcinus orca* in British Columbia. Marine Ecology Progress Series, 316:185-199.
- Ford, J.K.B., G.M. Ellis, P.F. Olesiuk, and K.C. Balcomb. 2009. Linking killer whale survival and prey abundance: food limitation in the oceans' apex predator? Biol. Lett. published online before print September 15, 2009.
- Ford, J.K.B., Stredulinsky, E.H., Ellis, G.M., Durban, J.W., Pilkington, J.F., 2014. Offshore Killer Whales in Canadian Pacific Waters: Distribution, Seasonality, Foraging Ecology, Population Status and Potential for Recovery. DFO Canadian Science Advisory Secretariat Research Document 2014/088. p. vii + 55.
- Ford M.J., J. Hempelmann, M.B. Hanson, K.L. Ayres, R.W. Baird, C.K. Emmons, J.I. Lundin, G.S. Schorr, S.K. Wasser, L.K. Park. 2016. Estimation of a Killer Whale (*Orcinus orca*) Population's Diet Using Sequencing Analysis of DNA from Feces. PLoS ONE 11(1): e0144956.
- Forney, K.A. and P. Wade. 2006. Worldwide distribution and abundance of killer whales. Pages 145-162 In: "Whales, whaling and ocean ecosystems", J.A. Estes, R.L. Brownell, Jr., D.P. DeMaster, D.F. Doak, and T.M. Williams (eds), University of California Press. 418p.
- Gearin, P. J., S. R. Melin, R. L. DeLong, H. Kajimura, and M. A. Johnson. 1994. Harbor porpoise interactions with a chinook salmon set-net fishery in Washington State. Rep. Int. Whal. Commn. Special Issue 15:427-438.

- Gearin, P. J., M. E. Gosho, J. L. Laake, L. Cooke, R. L. DeLong, and K. M. Hughes. 2000. Experimental testing of acoustic alarms (pingers) to reduce bycatch of harbour porpoise, *Phocoena phocoena*, in the state of Washington. *J. Cetacean Res. Manage.* 2(1):1-9.
- George, J.C., L.M. Philo, K. Hazard, D. Withrow, G.M. Carroll and R. Suydam. 1994. Frequency of killer whale (*Orcinus orca*) attacks and ship collisions based on scarring on bowhead whales (*Balaena mysticetus*) of the Bering-Chukchi-Beaufort seas stock. *Arctic* 47(3): 246-255.
- Guenther, T. J., R.W. Baird, R.L. Bates, P.M. Willis, R.L. Hahn, and S.G. Wischniowski. 1995. Strandings and fishing gear entanglements of cetaceans on the west coast of Canada in 1994. Paper SC/47/O6 presented to the International Whaling Commission, May 1995 (unpublished). 7 pp.
- Hanson, M.B., R.W. Baird, J.K.B. Ford, J. Hempelmann-Halos, D. M. Van Doornik, J.R. Candy, C. K. Emmons, G. S. Schorr, B. Gisborne, K. L. Ayres, S. K. Wasser, K. C. Balcomb, K. Balcomb-Bartok, J. G. Sneva, and M. J. Ford. 2010. Species and stock identification of prey consumed by endangered "southern resident" killer whales in their summer range. *Endangered Species Research* 11:69–82.
- Hanson, M.B., C.K. Emmons, E.J. Ward, J.A. Nystuen, and M.O. Lammers. 2013. Assessing the coastal occurrence of endangered killer whales using autonomous passive acoustic recorders. *J. Acoust. Soc. Am.* 134 (5): 3486–3495.
- Hanson, M.B., E.J. Ward, C.K. Emmons, M.M. Holt and D.M. Holzer. 2017. Assessing the movements and occurrence of Southern Resident Killer Whales relative to the U.S. Navy's Northwest Training Range Complex in the Pacific Northwest. Prepared for: U.S. Navy, U.S. Pacific Fleet, Pearl Harbor, HI. Prepared by: National Oceanic and Atmospheric Administration, Northwest Fisheries Science Center under MIPR N00070-15-MP-4C363. 30 June 2017. 23 pp.
- Hanson, M.B., C.K. Emmons, M.J. Ford, M. Everett, K. Parsons, L.K. Park, J. Hempelmann, D.M. Van Doornik, G.S. Schorr, J.K. Jacobsen, M.F. Sears, M.S. Sears, J.G. Sneva, R.W. Baird, L. Barre. 2021. Endangered predators and endangered prey: Seasonal diet of Southern Resident killer whales. *PLoS ONE* 16(3): e0247031.
- Hoelzel, A.R. 1991. Analysis of regional mitochondrial DNA variation in the killer whale; implications for cetacean conservation. *Rep. Int. Whal. Commn. Special Issue* 13:225-233.
- Hoelzel, A.R., and G. A. Dover. 1991. Genetic differentiation between sympatric killer whale populations. *Heredity* 66:191-195.
- Hoelzel, A. R., M. E. Dahlheim, and S. J. Stern. 1998. Low genetic variation among killer whales (*Orcinus orca*) in the Eastern North Pacific, and genetic differentiation between foraging specialists. *J. Heredity* 89:121-128.
- Krahn, M.M., M.J. Ford, W.F. Perrin, P.R. Wade, R.P. Angliss, M.B. Hanson, B.L. Taylor, G. Ylitalo, M.E. Dahlheim, J.E. Stein, and R.S. Waples. 2004. 2004 Status review of Southern Resident killer whales (*Orcinus orca*) under the Endangered Species Act. U.S. Dep. Commer., NOAA Tech. Memo NMFS-NWFSC-62. 73 pp.
- Krahn, M.M., M.B. Hanson, R.W. Baird, R.H. Boyer, D.G. Burrows, C.K. Emmons, J. K.B. Ford, L. L. Jones, D. P. Noren, P. S. Ross, G. S. Schorr, T.K. Collier. 2007. Persistent organic pollutants and stable isotopes in biopsy samples (2004/2006) from Southern Resident killer whales. *Mar. Poll. Bull.* 54 (2007) 1903–1911.
- Krahn, M.M., M.B. Hanson, G.S. Schorr, C.K. Emmons, D.G. Burrows, J.L. Bolton, R.W. Baird, G.M. Ylitalo. 2009. Effects of age, sex and reproductive status on persistent organic pollutant concentrations in "Southern Resident" killer whales. *Marine Pollution Bulletin* 58: 1522–1529.
- Lacy, R.C., Williams, R., Ashe, E., Balcomb III, K.C., Brent, L.J., Clark, C.W., Croft, D.P., Giles, D.A., MacDuffee, M. and Paquet, P.C., 2017. Evaluating anthropogenic threats to endangered killer whales to inform effective recovery plans. *Scientific reports*, 7(1), pp.1-12.
- Leatherwood, J. S., and M. E. Dahlheim. 1978. Worldwide distribution of pilot whales and killer whales. Naval Ocean Systems Center, Tech. Rep. 443:1-39.
- Leatherwood, S., C.O. Matkin, J.D. Hall, and G.M. Ellis. 1990. Killer whales, *Orcinus orca*, photo-identified in Prince William Sound, Alaska 1976 to 1987. *Can. Field Nat.* 104:362-371.
- Lowry, L.F., R.R. Nelson, and K.J. Frost. 1987. Observations of killer whales, *Orcinus orca*, in western Alaska: Sightings, strandings, and predation on other marine mammals. *The Canadian Field Naturalist* 101:6-12.
- Matkin, C., G. Ellis, E. Saulitis, L. Barrett-Lennard, and D. Matkin. 1999. Killer Whales of Southern Alaska. *North Gulf Oceanic Society*. 96 pp.
- Matkin, C.O., Ward Testa, J., Ellis, G. M. and Saulitis, E. L. 2014. Life history and population dynamics of southern Alaska resident killer whales (*Orcinus orca*). *Marine Mammal Science*, 30: 460–479.
- Morin P. A., F. I. Archer, A. D. Foote, J. Vilstrup, E. E. Allen, P. R. Wade, J. W. Durban, K. M. Parsons, R. Pitman, L. Li, P. Bouffard, S. C. Abel Nielsen, M. Rasmussen, E. Willerslev, M. T. P. Gilbert, T. Harkins. 2010. Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Research*, 20:908-916.
- NOAA. 2014. Wild animal mortality investigation: Southern Resident Killer Whale L-112 final report.

- NMFS. 2021. Endangered and Threatened Wildlife and Plants; Revision of Critical Habitat for the Southern Resident Killer Whale Distinct Population Segment. Final Rule. 86 FR 41668.
- NMFS. 2019b. Proposed Revision of the Critical Habitat Designation for Southern Resident Killer Whales Draft Biological Report (to accompany the Proposed Rule). 30 pp + Appendices.
- NMFS. 2016. Southern Resident Killer Whale (*Orcinus orca*) Stranding Event Expert Review Summary, September 21, 2016. Available at: https://archive.fisheries.noaa.gov/wcr/publications/protected_species/marine_mammals/killer_whales/195_expert_panel.pdf
- Northwest Indian Fisheries Commission (NWIFC). 1995. Monitoring of marbled murrelet and marine mammal interactions with 1994 tribal gillnet fisheries in northern Puget Sound, Hood Canal, and the Strait of Juan de Fuca. Final Report to NMFS, Contract No. 52ABNF400087, and U.S. Fish and Wildlife Service. Unpubl. report. 41 pp. Available at NWIFC, 6730 Martin Way E, Olympia, WA 98516.
- Olesiuk, P.F., M.A. Bigg, and G.M. Ellis. 1990. Life history and population dynamics of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Rep. Int. Whal. Commn. Special Issue 12:209-242.
- Pierce, D. J., W. P. Ritchie, and R. Kreuziger. 1994. Preliminary findings of seabird interactions with the non-treaty salmon gill net fishery: Puget Sound and Hood Canal Washington. Unpubl. report. Washington Dept. Fish and Wildlife, Olympia, WA. 39 pp. Available at WDFW, 600 Capitol Way N, Olympia, WA 98501.
- Pierce, D. J., M. Alexandersdottir, S.J. Jeffries, P. Erstad, W. Beattie, and A. Chapman. 1996. Interactions of marbled murrelets and marine mammals with the 1994 Puget Sound sockeye gill net fishery. Final Report, Washington Dept. Fish and Wildlife, Olympia, WA. 21 pp.
- Raverty S, St. Leger J, Noren DP, Burek Huntington K, Rotstein DS, Gulland, FMD, Ford, JKB, Hanson, MB, Lambourn, DM, Huggins, J, Delaney, MA, Spaven, L, Rowles, T, Barre, L, Cottrell, P, Ellis, G, Goldstein, T, Terio K, Duffield, D, Rice, J, Gaydos, JK. 2020. Pathology findings and correlation with body condition index in stranded killer whales (*Orcinus orca*) in the northeastern Pacific and Hawaii from 2004 to 2013. PLOS ONE 15(12): e0242505. <https://doi.org/10.1371/journal.pone.0242505>
- Reeves, R.R., W.F. Perrin, B.L. Taylor, C.S. Baker, and S.L. Mesnick. 2004. Report of the workshop on shortcomings of cetacean taxonomy in relation to needs of conservation and management, April 30 – May 2, 2004, La Jolla, California. U.S. Department of Commerce NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-363. 94pp. Available from Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA. 92037.
- Sericano, J.L., T.L. Wade, S.T. Sweet, J. Ramirez and G.G. Lauenstein. 2014. Temporal trends and spatial distribution of DDT in bivalves from the coastal marine environments of the continental United States, 1986–2009. Marine Pollution Bulletin 81: 303-316.
- Stevens, T.A., D. Duffield, E. Asper, K. Hewlett, A. Bolz, L. Gage, and G. Bossart. 1989. Preliminary findings of restriction fragment differences in mitochondrial DNA among killer whales (*Orcinus orca*). Can. J. Zool. 67:2592-2595.
- Wade, P.R., and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Ward, E.J., E.E. Holmes, and K.C. Balcomb. 2009. Quantifying the effects of prey abundance on killer whale reproduction. Journal of Applied Ecology, 46(3):632-640.
- Wasser S.K., Lundin J.I., Ayres K., Seely E., Giles D., Balcomb K., et al. 2017. Population growth is limited by nutritional impacts on pregnancy success in endangered Southern Resident killer whales (*Orcinus orca*). PLoS ONE 12(6): e0179824.
- Yano, K., and M.E. Dahlheim. 1995. Killer whale, *Orcinus orca*, depredation on longline catches of bottomfish in the southeastern Bering Sea and adjacent waters. Fish. Bull. 93:355-372.