

COMMON BOTTLENOSE DOLPHIN (*Tursiops truncatus truncatus*): Gulf of Mexico Western Coastal Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Common bottlenose dolphins inhabit coastal waters throughout the northern Gulf of Mexico (i.e., U.S. Gulf of Mexico; Mullin *et al.* 1990). As a working hypothesis, it is assumed that the dolphins occupying habitats with dissimilar climatic, coastal and/or oceanographic characteristics might be restricted in their movements between habitats, and thus constitute separate stocks. Therefore, northern Gulf of Mexico coastal waters have been divided for management purposes into three stock areas: eastern, northern and western, with coastal waters defined as waters between the shore, barrier islands or presumed outer bay boundaries out to the 20-m isobath (Figure 1). The 20-m depth seaward boundary corresponds to survey strata (Scott 1990; Blaylock and Hoggard 1994; Fulling *et al.* 2003) and thus represents a management boundary rather than an ecological boundary. The Western Coastal common bottlenose dolphin stock area extends from the Mississippi River Delta to the Texas-Mexico border. This region is characterized by an arid to temperate climate, sand beaches in southern Texas, extensive coastal marshes in northern Texas and Louisiana, and varying amounts of freshwater input. Dolphins belonging to this stock are all expected to be of the coastal ecotype (Vollmer 2011). The Western Coastal Stock is trans-boundary with Mexico; however, there is no information available for abundance estimation, nor for estimating fishery-related mortality in Mexican waters.




Figure 1. Distribution of common bottlenose dolphin on-effort sightings in coastal and continental shelf waters during SEFSC aerial surveys in summer 2017, winter 2018, and fall 2018. Sightings within the boundaries of the Western Coastal Stock are denoted by the black circles. Isobaths are the 20-m, 200-m, 1,000-m, and 2,000-m depth contours.

Recently, genetic analyses of population structure in coastal, shelf, and oceanic waters of the Gulf of Mexico revealed seven demographically independent populations in the northern Gulf of Mexico, suggesting the current stock designations and boundaries in these waters do not accurately reflect the population structure (Vollmer and Rosel 2017). Sampling within the range of the Western Coastal Stock was limited and further work is necessary to determine the boundaries of these demographically independent populations.

This stock's boundaries abut other common bottlenose dolphin stocks, namely the Northern Coastal Stock, Continental Shelf Stock and several bay, sound and estuary stocks in Texas and Louisiana, and while individuals from

different stocks may occasionally overlap, it is not thought that significant mixing or interbreeding occurs between them. Fazioli *et al.* (2006) conducted photo-identification surveys of coastal waters off Tampa Bay, Sarasota Bay and Lemon Bay, Florida, over 14 months. They found both ‘inshore’ and ‘Gulf’ dolphins inhabited coastal waters but the two types used coastal waters differently. Dolphins from the inshore communities were observed occasionally in Gulf near-shore waters adjacent to their inshore range, whereas ‘Gulf’ dolphins were found primarily in open Gulf of Mexico waters with some displaying seasonal variations in their use of the study area. The ‘Gulf’ dolphins did not show a preference for waters near passes as was seen for ‘inshore’ dolphins, but moved throughout the study area and made greater use of waters offshore of waters used by ‘inshore’ dolphins. During winter months abundance of ‘Gulf’ groups decreased while abundance for ‘inshore’ groups increased. These findings support an earlier report by Irvine *et al.* (1981) of increased use of pass and coastal waters by Sarasota Bay dolphins in winter. Seasonal movements of identified individuals and abundance indices suggested that part of the ‘Gulf’ dolphin community moved out of the study area during winter, but their destination is unknown (Fazioli *et al.* 2006). In a follow-up study, Sellas *et al.* (2005) examined genetic population subdivision in the study area of Fazioli *et al.* (2006), and found evidence of significant population structure among all areas. Rosel *et al.* (2017) also identified significant genetic differentiation between estuarine residents of Barataria Bay and the adjacent coastal stock, further supporting separation of coastal and estuarine stocks.

Finally, off Galveston, Texas, Beier (2001) reported an open population of individual dolphins in coastal waters, but several individual dolphins had been sighted previously by other researchers over a 10-year period. Some coastal animals may move relatively long distances alongshore. Two bottlenose dolphins previously seen in the South Padre Island area in Texas were seen in Matagorda Bay, 285 km north, in May 1992 and May 1993 (Lynn and Würsig 2002).

POPULATION SIZE

The best abundance estimate available for the northern Gulf of Mexico Western Coastal Stock of common bottlenose dolphins is 20,759 (CV=0.13; Table 1; Garrison *et al.* 2021). This estimate is from an inverse-variance weighted average of seasonal abundance estimates from aerial surveys conducted during summer 2017 and fall 2018.

Earlier Abundance Estimates

Please see Appendix IV for a summary of abundance estimates, including earlier estimates and survey descriptions.

Recent Surveys and Abundance Estimates

The Southeast Fisheries Science Center conducted aerial surveys of continental shelf waters (shoreline to 200 m depth) along the U.S. Gulf of Mexico coast from the Florida Keys to the Texas/Mexico border during summer (June–August) 2017 and fall (October–November) 2018. The stock was only partially surveyed during a winter 2018 aerial survey, and therefore this survey was not included in the current abundance estimates (Garrison *et al.* 2021). The surveys were conducted along tracklines oriented perpendicular to the shoreline and spaced 20 km apart. The total survey effort varied during each survey due to weather conditions, and was 10,781 km (fall) and 14,590 km (summer). Each of these surveys was conducted using a two-team approach to develop estimates of visibility bias using the independent observer approach with Distance analysis (Laake and Borchers 2004). Abundance was calculated using mark-recapture distance sampling implemented in package mrds (version 2.21; Laake *et al.* 2020) in the R statistical programming language. This approach estimates both the probability of detection on the trackline and within the surveyed strip accounting for the effects of sighting conditions (e.g., sea state, glare, turbidity, and cloud cover). A different detection probability model was used for each seasonal survey (Garrison *et al.* 2021). The survey data were post-stratified into spatial boundaries corresponding to the defined boundaries of common bottlenose dolphin stocks within the surveyed area. The abundance estimates for the Western Coastal Stock of common bottlenose dolphins were based upon tracklines and sightings in waters from the shoreline to the 20-m isobath and between the Texas-Mexico border and the Mississippi River Delta. The seasonal abundance estimates for this stock were: summer – 18,601 (CV=0.30) and fall – 21,766 (CV=0.14). Due to the uncertainty in stock movements and apparent seasonal variability in the abundance of the stock, a weighted average of these seasonal estimates was taken where the weighting was the inverse of the CV. This approach weights estimates with higher precision more heavily in the final weighted mean. The resulting weighted mean and best estimate of abundance for the Western Coastal Stock of common bottlenose dolphins was 20,759 (CV=0.13; Table 1; Garrison *et al.* 2021).

Table 1. Most recent abundance estimate (*N_{est}*) and coefficient of variation (*CV*) of the northern Gulf of Mexico Western Coastal Stock of common bottlenose dolphins (0–20-m isobaths) based on summer 2017, winter 2018, and fall 2018 aerial surveys.

Years	Area	Nest	CV
2017, 2018	Gulf of Mexico	20,759	0.13

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for the Western Coastal Stock of common bottlenose dolphins is 20,759 (CV=0.13). Therefore, the minimum population estimate for the northern Gulf of Mexico Western Coastal Stock is 18,585 (Table 2).

Current Population Trend

The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long intervals between surveys. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV>0.30) remains below 80% ($\alpha=0.30$) unless surveys are conducted on an annual basis (Taylor *et al.* 2007). Two point estimates of common bottlenose dolphin abundance for the Western Coastal Stock have been made based on aerial data from surveys during 2011–2012 and 2017–2018 (Garrison *et al.* 2021). Each of these surveys had a similar design and was conducted using the same aircraft and observer configuration. The resulting inverse variance weighted best abundance estimates for seasonal surveys were: 2011–2012 – 19,381 (CV=0.20) and 2017–2018 – 20,759 (CV=0.13). A trends analysis is not possible because there are only two abundance estimates available. For further information on comparisons of old and current abundance estimates for this stock see Garrison *et al.* (2021).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are not known for this stock. The maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate and a recovery factor (Wade and Angliss 1997). The minimum population size is 18,585. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor is 0.45 because the CV of the shrimp trawl mortality estimate is greater than 0.6 (Wade and Angliss). PBR for the northern Gulf of Mexico Western Coastal Stock of common bottlenose dolphins is 167 (Table 2).

Table 2. Best and minimum abundance estimates of the northern Gulf of Mexico Western Coastal Stock of common bottlenose dolphins with Maximum Productivity Rate (*R_{max}*), Recovery Factor (*Fr*) and PBR.

Nest	Nest CV	Nmin	Fr	Rmax	PBR
20,759	0.13	18,585	0.45	0.04	167

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual human-caused mortality and serious injury for the Western Coastal Stock of common bottlenose dolphins during 2015–2019 is unknown because this stock is known to interact with unobserved fisheries (see below). The five-year unweighted mean annual mortality estimate for 2015–2019 for the commercial shrimp trawl fishery was 32 (CV=0.65; see Shrimp Trawl section below). The mean annual fishery-related mortality and serious injury during 2015–2019 for strandings identified as fishery-caused was 0.4. Mean annual mortality and serious injury during 2015–2019 due to other human-caused actions (the *Deepwater Horizon* (DWH) oil spill and foreign fisheries) was predicted to be 3.2. The minimum total mean annual human-caused mortality and serious injury for this stock during 2015–2019 was 36 (Table 3). This is considered a minimum because 1) not all fisheries that could interact with this stock are observed and/or observer coverage is very low, 2) stranding data are used as an indicator of fishery-related interactions

and not all dead animals are recovered by the stranding network (Peltier *et al.* 2012; Wells *et al.* 2015), 3) cause of death is not (or cannot be) routinely determined for stranded carcasses, 4) the estimate of fishery-related interactions includes an actual count of verified fishery-caused deaths and serious injuries and should be considered a minimum (NMFS 2016), 5) various assumptions were made in the population model used to estimate population decline for the northern Gulf of Mexico Bay Stock and Estuaries (BSE) stocks impacted by the DWH oil spill.

Fisheries Information

There are five commercial fisheries that interact, or that potentially could interact, with this stock. These include three Category II fisheries (Southeastern U.S. Atlantic, Gulf of Mexico shrimp trawl; Gulf of Mexico menhaden purse seine; and Gulf of Mexico gillnet); and two Category III fisheries (Gulf of Mexico blue crab trap/pot; and Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line)). Detailed fishery information is presented in Appendix III.

Note: Animals reported in the sections to follow were ascribed to a stock or stocks of origin following methods described in Maze-Foley et al. (2019). These include strandings, observed takes (through an observer program), fisherman self-reported takes (through the Marine Mammal Authorization Program), research takes, and opportunistic at-sea observations.

Shrimp Trawl

Between 1997 and 2019, 13 common bottlenose dolphins and nine unidentified dolphins, which could have been either common bottlenose dolphins or Atlantic spotted dolphins, became entangled in the lazy line, turtle excluder device or tickler chain gear in observed trips of the commercial shrimp trawl fishery in the Gulf of Mexico (Soldevilla *et al.* 2021). All dolphin bycatch interactions resulted in mortalities except for one unidentified dolphin that was released alive in 2009 (Maze-Foley and Garrison 2016). Soldevilla *et al.* (2015, 2016, 2021) provided mortality estimates calculated from analysis of shrimp fishery effort data and NMFS's Observer Program bycatch data. Annual mortality estimates were calculated for the years 2015–2019 from stratified annual fishery effort and bycatch rates, and the five-year unweighted mean mortality estimate was calculated for Gulf of Mexico dolphin stocks (Soldevilla *et al.* 2021). The four-area (TX, LA, MS/AL, FL) stratification method was chosen because it best approximates how fisheries operate (Soldevilla *et al.* 2015, 2016, 2021). The mean annual mortality estimate for the Western Coastal Stock of common bottlenose dolphins is 32 (CV=0.65). Limitations and biases of annual bycatch mortality estimates are described in detail in Soldevilla *et al.* (2015, 2016, 2021).

In addition, chaffing gear from a commercial shrimp trawl was recovered in a dolphin carcass that stranded during 2015. It is likely the animal ingested the gear while removing gilled fish that were caught in the trawl net. This animal was ascribed to both the Barataria Bay Estuarine System and Western Coastal stocks, and it was included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 25 August 2020; Table 4).

Menhaden Purse Seine

During 2015–2019, no interactions between the Western Coastal Stock and the menhaden purse seine fishery were documented. There is currently no observer program for the Gulf of Mexico menhaden purse seine fishery. Previously, interactions between the Western Coastal stock and this fishery have been documented by both a pilot observer program and the Marine Mammal Authorization Program. Without an ongoing observer program it is not possible to obtain statistically reliable information for this fishery on the number of sets annually, the incidental take and mortality rates, and the communities from which bottlenose dolphins are being taken.

Gillnet

No marine mammal mortalities associated with U.S. gillnet fisheries have been reported or observed for the Western Coastal Stock. There is limited observer coverage of this fishery in federal waters (e.g., Mathers *et al.* 2020), but none currently in state waters, although during 2012–2018 NMFS placed observers on commercial vessels (state permitted gillnet vessels) in the coastal state waters of Alabama, Mississippi, and Louisiana (Mathers *et al.* 2016). No takes were observed in state waters during this time. Because there is no observer program within this stock's boundaries, it is not possible to estimate the total number of interactions or mortalities associated with gillnet gear.

Blue Crab Trap/Pot

During 2015–2019, no interactions were documented for the Western Coastal Stock with crab trap/pot fisheries.

An earlier interaction was documented for this stock (from 2008). Since there is no observer program, it is not possible to estimate the total number of interactions or mortalities associated with crab traps/pots.

Hook and Line (Rod and Reel)

During 2015–2019, one mortality involving hook and line gear entanglement was documented for the Western Coastal Stock. The mortality occurred in 2018, and available evidence from the stranding record suggested the hook and line gear interaction contributed to the cause of death. The mortality was included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 25 August 2020) and in the stranding totals presented in Table 4, and in the annual human-caused mortality and serious injury total (Table 3).

It should be noted that, in general, it cannot be determined if hook and line gear originated from a commercial (i.e., charter boat and headboat) or recreational angler because the gear type used by both sources is typically the same. Also, it is not possible to estimate the total number of interactions with hook and line gear because there is no observer program. The documented interaction in this gear represents a minimum known count of interactions in the last five years.

Other Mortality

A population model was developed to estimate long-term injury to stocks affected by the DWH oil spill (see Habitat Issues section), taking into account long-term effects resulting from mortality, reproductive failure, and reduced survival rates (DWH MMIQT 2015; Schwacke *et al.* 2017). For the Western Coastal Stock, the model predicted the stock experienced a 5% (95%CI: 3–9) maximum reduction in population size due to the oil spill (DWH MMIQT 2015; DWH NRDAT 2016; Schwacke *et al.* 2017), and for the years 2015–2019, the model projected 16 mortalities (Table 3). This population model has a number of sources of uncertainty. The baseline population size was estimated from studies initiated after initial exposure to DWH oil occurred. Therefore, it is possible that the pre-spill population size was larger than this baseline level and some mortality occurring early in the event was not quantified. The duration of elevated mortality and reduced reproductive success after exposure is unknown, and expert opinion was used to predict the rate at which these parameters would return to baseline levels. Where possible, uncertainty in model parameters was included in the estimates of excess mortality by re-sampling from statistical distributions of the parameters (DWH MMIQT 2015; DWH NRDAT 2016; Schwacke *et al.* 2017).

In addition to the fishery interactions discussed above, two additional fishery-related mortalities were documented during 2015–2019. One mortality was documented in 2017 as a result of entanglement in monofilament line. It could not be determined if the line was part of a net or not. In 2018, an additional mortality was documented near the Texas/Mexico border in Mexican shark gillnet gear. Both of these interactions were included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 25 August 2020) and in the totals presented in Table 4, and also in the annual human-caused mortality and serious injury total for this stock (Table 3).

NOAA's Office of Law Enforcement has been investigating increased reports from along the northern Gulf of Mexico coast of violence against common bottlenose dolphins, including shootings via guns and bows and arrows, throwing pipe bombs and cherry bombs, and stabbings (Vail 2016). During 2015–2019, for one mortality, gunshot pellets were found during the necropsy. The gunshot occurred pre-mortem but was not believed to be the cause of death. This animal was included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 25 August 2020) and in the totals presented in Table 4, but was not included in the annual human-caused mortality and serious injury total for this stock (Table 3). From recent cases that have been prosecuted, it has been shown that fishermen became frustrated and retaliated against dolphins for removing bait or catch, or depredating their fishing gear. It is unknown whether the 2019 shooting involved depredation.

Depredation of fishing catch and/or bait is a growing problem in Gulf of Mexico coastal and estuary waters and globally, and can lead to serious injury or mortality via ingestion of or entanglement in gear (e.g., Zollett and Read 2006; Read 2008; Powell and Wells 2011; Vail 2016), as well as changes to the dolphin's activity patterns, such as decreases in natural foraging (Powell and Wells 2011). It has been suggested that provisioning, or the illegal feeding, of wild common bottlenose dolphins, may encourage depredation because provisioning conditions dolphins to approach humans and vessels, where they then may prey on bait and catches (Vail 2016). Such conditioning increases risks of subsequent injury and mortality (Christiansen *et al.* 2016). Provisioning has been documented in the literature in Florida and Texas (Bryant 1994; Samuels and Bejder 2004; Cunningham-Smith *et al.* 2006; Powell and Wells

2011). To date there are no records within the literature of provisioning for this stock area.

As part of its annual coastal dredging program, the Army Corps of Engineers conducts sea turtle relocation trawling during hopper dredging as a protective measure for marine turtles. No interactions have been documented during the most recent five years, 2015–2019. Historically, two mortalities were documented involving relocation trawling activities and common bottlenose dolphins likely belonging to the Western Coastal Stock (2005, 2007).

All mortalities and serious injuries from known sources for the Western Coastal Stock are summarized in Table 3.

Table 3. Summary of the incidental mortality and serious injury of common bottlenose dolphins (*Tursiops truncatus*) of the Western Coastal Stock. For fisheries that do not have an ongoing, federal observer program, counts of mortality and serious injury were based on stranding data, at-sea observations, or fisherman self-reported takes via the Marine Mammal Authorization Program (MMAP). For strandings, at-sea counts, and fisherman self-reported takes, the number reported is a minimum because not all strandings, at-sea cases, or gear interactions are detected. See the Annual Human-Caused Mortality and Serious Injury section for biases and limitations of mortality estimates, and the Strandings section for limitations of stranding data. NA = not applicable.

Fishery	Years	Data Type	Mean Annual Estimated Mortality and Serious Injury Based on Observer Data	5-year Minimum Count Based on Stranding, At-Sea, and/or MMAP Data
Shrimp Trawl	2015–2019	Observer Data	32 (CV=0.65)	NA
Menhaden Purse Seine	2015–2019	MMAP Fisherman self-reported takes	NA	0
Gillnet	2015–2019	Observer Data and Stranding Data	NA	0
Crab Trap/Pot	2015–2019	Stranding Data	NA	0
Hook and Line	2015–2019	Stranding Data and At-Sea Observations	NA	1
Unknown Gear	2015–2019	Stranding Data	NA	1
Mean Annual Mortality due to commercial fisheries (2015–2019)			32.4	
Other Takes (foreign fishing gear, 5-year Count)			1	
Mortality due to DWH (5-year Projection)			16	
Mean Annual Mortality due to other takes and DWH (2015–2019)			3.2	
Minimum Total Mean Annual Human-Caused Mortality and Serious Injury (2015–2019)			36	

Strandings

A total of 586 common bottlenose dolphins were found stranded in Western Coastal Stock waters of the northern Gulf of Mexico from 2015 through 2019 (Table 4; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 25 August 2020). There was evidence of human interaction (HI) for 26 of the strandings. No evidence of human interaction was detected for 63 strandings, and for the remaining 497 strandings, it could not be determined if there was evidence of human interaction. Human interactions were from several sources, including an entanglement with hook and line gear, an entanglement in a Mexican shark gillnet, and an animal with gunshot wounds. It should be noted that evidence of human interaction does not necessarily mean the interaction caused the animal's stranding or death.

The assignment of animals to a single stock is impossible in some regions where stocks overlap, especially in nearshore coastal waters (Maze-Foley *et al.* 2019). Of the 586 strandings ascribed to the Western Coastal Stock, 441 were ascribed solely to this stock. The counts in Table 4 may include some animals from the Barataria Bay Estuarine System; Terrebonne-Timbalier Bay Estuarine System; Galveston Bay, East Bay, Trinity Bay; West Bay; Calcasieu Lake; Nueces Bay, Corpus Christi Bay; or Laguna Madre stocks, and thereby overestimate the number of strandings for the Western Coastal Stock. Stranded carcasses are not routinely identified to either the offshore or coastal morphotype of common bottlenose dolphin, therefore it is possible that some of the reported strandings were of the offshore form, though that number is likely to be low (Byrd *et al.* 2014).

There are a number of other difficulties associated with the interpretation of stranding data. Stranding data underestimate the extent of human and fishery-related mortality and serious injury because not all of the dolphins that die or are seriously injured in human interactions wash ashore, or, if they do, they are not all recovered (Peltier *et al.* 2012; Wells *et al.* 2015; Carretta *et al.* 2016). Additionally, not all carcasses will show evidence of human interaction, entanglement or other fishery-related interaction due to decomposition, scavenger damage, etc. (Byrd *et al.* 2014). Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of human interaction.

Since 1990, there have been 15 bottlenose dolphin die-offs or Unusual Mortality Events (UMEs) in the northern Gulf of Mexico (<http://www.nmfs.noaa.gov/pr/health/mmume/events.html>, accessed 5 November 2020), and eight of these have occurred within the boundaries of the Western Coastal Stock and may have affected the stock. 1) From January through May 1990, a total of 344 bottlenose dolphins stranded in the northern Gulf of Mexico. Overall this represented a two-fold increase in the prior maximum recorded strandings for the same period, but in some locations (i.e., Alabama) strandings were 10 times the average number. The cause of the 1990 mortality event could not be determined (Hansen 1992), however, morbillivirus may have contributed to this event (Litz *et al.* 2014). 2) In March and April 1992, 119 bottlenose dolphins stranded in Texas, about nine times the average number. The cause of this event was not determined, but low salinity due to record rainfall combined with pesticide runoff and exposure to morbillivirus were suggested as potential contributing factors (Duignan *et al.* 1996; Colbert *et al.* 1999; Litz *et al.* 2014). 3) In 1993–1994 a UME of bottlenose dolphins likely caused by morbillivirus started in the Florida Panhandle and spread west with most of the mortalities occurring in Texas (Lipscomb 1993; Lipscomb *et al.* 1994; Litz *et al.* 2014). From February through April 1994, 236 bottlenose dolphins were found dead on Texas beaches, of which 67 occurred in a single 10-day period. 4) During February and March of 2007 an event was declared for northeast Texas and western Louisiana involving 64 bottlenose dolphins and two unidentified dolphins. Decomposition prevented conclusive analyses on most carcasses. 5) During February and March of 2008 an additional event was declared in Texas involving 111 bottlenose dolphin strandings (plus strandings of one unidentified dolphin and one melon-headed whale). Most of the animals recovered were in a decomposed state. The event has been closed, however, the investigation is ongoing. 6) A UME was declared for cetaceans in the northern Gulf of Mexico beginning 1 March 2010 and ending 31 July 2014 (Litz *et al.* 2014; http://www.nmfs.noaa.gov/pr/health/mmume/cetacean_gulfofmexico.htm, accessed 2 December 2020). It included cetaceans that stranded prior to the DWH oil spill (see Habitat Issues section below), during the spill, and after. Exposure to the DWH oil spill was determined to be the primary underlying cause of the elevated stranding numbers in the northern Gulf of Mexico after the spill (e.g., Schwacke *et al.* 2014; Venn-Watson *et al.* 2015; Colegrove *et al.* 2016; DWH NRDAT 2016). 7) A UME occurred from November 2011 to March 2012 across five Texas counties including 126 bottlenose dolphin strandings. Ninety-six animals from this stock were considered to be part of the UME. The strandings were coincident with a harmful algal bloom of *Karenia brevis*, but researchers have not determined that was the cause of the event. 8) During 1 February 2019 to 30 November 2019, a UME was declared for the area from the eastern border of Taylor County, Florida, west through Alabama, Mississippi, and Louisiana (<http://www.nmfs.noaa.gov/pr/health/mmume/>

cetacean_gulfofmexico.htm, accessed 5 November 2020). A total of 337 common bottlenose dolphins stranded during this event, with 44 of them being from the Western Coastal Stock. The largest number of mortalities occurred in eastern Louisiana and Mississippi. An investigation concluded the event was caused by exposure to low salinity waters as a result of extreme freshwater discharge from rivers. The unprecedented amount of freshwater discharge during 2019 (e.g., Gasparini and Yuill 2020) resulted in low salinity levels across the region.

Table 4. Common bottlenose dolphin strandings occurring in the Western Coastal Stock area from 2015 to 2019, including the number of strandings for which evidence of human interaction (HI) was detected and number of strandings for which it could not be determined (CBD) if there was evidence of HI. Data are from the NOAA National Marine Mammal Health and Stranding Response Database (unpublished data, accessed 25 August 2020). Please note HI does not necessarily mean the interaction caused the animal's death.

Stock	Category	2015	2016	2017	2018	2019	Total
Western Coastal Stock	Total Stranded	94	100	143	123	126 ^d	586
	Human Interaction						
	---Yes	3 ^a	2	6 ^b	8 ^c	7 ^e	26
	---No	3	6	22	16	16	63
	---CBD	88	92	115	99	103	497

a. Includes 1 interaction with chaffing gear from a commercial shrimp trawl (mortality).

b. Includes 2 fisheries interactions (FIs).

c. Includes FIs, including 1 interaction with hook and line gear (mortality) and 1 interaction with a Mexican shark gillnet (mortality).

d. 44 strandings were part of the UME event in the northern Gulf of Mexico.

e. Includes 1 FI and 1 animal with evidence of gunshot wounds (mortality).

HABITAT ISSUES

The *Deepwater Horizon* MC252 drilling platform, located approximately 80 km southeast of the Mississippi River Delta in waters about 1500 m deep, exploded on 20 April 2010. The rig sank, and over 87 days up to ~3.2 million barrels of oil were discharged from the wellhead until it was capped on 15 July 2010 (DWH NRDAT 2016).

Shortly after the oil spill, the Natural Resource Damage Assessment (NRDA) process was initiated under the Oil Pollution Act of 1990. A variety of NRDA research studies were conducted to determine potential impacts of the spill on marine mammals. These studies estimated that 23% (95%CI: 16–32) of the Western Coastal Stock of common bottlenose dolphins in the Gulf were exposed to oil, that 10% (95%CI: 5–15) of females suffered from reproductive failure, and 8% (95%CI: 3–13) suffered adverse health effects (DWH MMIQT 2015). A population model estimated that the stock experienced a 5% maximum reduction in population size (see Other Mortality section above).

In 2014, a vessel collision in Galveston Bay near Texas City released approximately 168,000 gallons of intermediate fuel oil. Through the National Resource Damage Assessment (NRDA) process, impacts of this spill are currently being evaluated and will include impacts to common bottlenose dolphins of the Western Coastal Stock (NOAA DAARP 2018).

The nearshore habitat occupied by the three coastal stocks is adjacent to areas of high human population and in some areas, such as Tampa Bay, Florida, Galveston, Texas, and Mobile, Alabama, is highly industrialized. Concentrations of anthropogenic chemicals such as PCBs and DDT and its metabolites vary from site to site, and can reach levels of concern for bottlenose dolphin health and reproduction in the southeastern U.S. (Schwacke *et al.* 2002). PCB concentrations in three stranded dolphins sampled from the Eastern Coastal Stock area ranged from 16–46µg/g wet weight. Two stranded dolphins from the Northern Coastal Stock area had the highest levels of DDT derivatives of any of the bottlenose dolphin liver samples analyzed in conjunction with a 1990 mortality investigation conducted by NMFS (Varanasi *et al.* 1992). The significance of these findings is unclear, but there is some evidence that increased exposure to anthropogenic compounds may reduce immune function in bottlenose dolphins (Lahvis *et al.* 1995), or impact reproduction through increased first-born calf mortality (Wells *et al.* 2005). Concentrations of chlorinated hydrocarbons and metals were relatively low in most of the bottlenose dolphins examined in conjunction with an anomalous mortality event in Texas bays in 1990; however, some had concentrations at levels of possible toxicological concern (Varanasi *et al.* 1992). Agricultural runoff following periods of high rainfall in 1992 was implicated in a high level of bottlenose dolphin mortalities in Matagorda Bay, which is adjacent to the Western Coastal Stock area.

The Mississippi River, which drains about two-thirds of the continental U.S., flows into the north-central Gulf of Mexico and deposits its nutrient load which is linked to the formation of one of the world's largest areas of seasonal

hypoxia (Rabalais *et al.* 1999). This area is located in Louisiana coastal waters west of the Mississippi River delta. How it affects bottlenose dolphins is not known.

STATUS OF STOCK

The common bottlenose dolphin is not listed as threatened or endangered under the Endangered Species Act, and the Gulf of Mexico Western Coastal Stock is not considered strategic under the Marine Mammal Protection Act. However, the occurrence of a UME of unprecedented size and duration has impacted the Western Coastal Stock area and is cause for concern. Total U.S. fishery-related mortality and serious injury for this stock is unknown, but at a minimum is greater than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The status of this stock relative to optimum sustainable population in the Gulf of Mexico EEZ is unknown. There are insufficient data to determine the population trends for this stock.

REFERENCES CITED

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background and a summary of the 1995 Assessments. NOAA Tech. Memo. NMFS-OPR-6. 73pp.
- Beier, A.G. 2001. Occurrence, distribution, and movement patterns of outer coastline bottlenose dolphins off Galveston, Texas. M.Sc. Thesis from Texas A&M University. 97pp.
- Blaylock, R.A. and W. Hoggard. 1994. Preliminary estimates of bottlenose dolphin abundance in southern U.S. Atlantic and Gulf of Mexico continental shelf waters. NOAA Tech. Memo NMFS-SEFSC-356. 10pp.
- Bryant, L. 1994. Report to Congress on results of feeding wild dolphins: 1989–1994. National Marine Fisheries Service, Office of Protected Resources. 23pp.
- Byrd, B.L., A.A. Hohn, G.N. Lovewell, K.M. Altman, S.G. Barco, A. Friedlaender, C.A. Harms, W.A. McLellan, K.T. Moore, P.E. Rosel and V.G. Thayer. 2014. Strandings illustrate marine mammal biodiversity and human impacts off the coast of North Carolina, USA. *Fish. Bull.* 112:1–23.
- Carretta, J.V., K. Danil, S.J. Chivers, D.W. Weller, D.S. Janiger, M. Berman-Kowalewski, K.M. Hernandez, J.T. Harvey, R.C. Dunkin, D.R. Casper, S. Stoudt, M. Flannery, K. Wilkinson, J. Huggins and D.M. Lambourn. 2016. Recovery rates of bottlenose dolphin (*Tursiops truncatus*) carcasses estimated from stranding and survival rate data. *Mar. Mamm. Sci.* 32(1):349–362.
- Christiansen, F., K.A. McHugh, L. Bejder, E.M. Siegal, D. Lusseau, E. Berens McCabe, G. Lovewell and R.S. Wells. 2016. Food provisioning increases the risk of injury and mortality in a long-lived marine top predator. *Royal Society Open Science* 3:160560.
- Colbert, A.A., G.I. Scott, M.H. Fulton, E.F. Wirth, J.W. Daugomah, P.B. Key, E.D. Strozier and S.B. Galloway. 1999. Investigation of unusual mortalities of bottlenose dolphins along the mid-Texas coastal bay ecosystem during 1992. NOAA Technical Report NMFS 147, Seattle, Washington. 23pp. <http://spo.nmfs.noaa.gov/tr147.pdf>
- Colegrove, K.M., S. Venn-Watson, J. Litz, M.J. Kinsel, K.A. Terio, E. Fougères, R. Ewing, D.A. Pabst, W.A. McLellan, S. Raverty, J. Saliki, S. Fire, G. Rappucci, S. Bowen-Stevens, L. Noble, A. Costidis, M. Barbieri, C. Field, S. Smith, R.H. Carmichael, C. Chevis, W. Hatchett, D. Shannon, M. Tumlin, G. Lovewell, W. McFee and T.K. Rowles. 2016. Fetal distress and *in utero* pneumonia in perinatal dolphins during the Northern Gulf of Mexico unusual mortality event. *Dis. Aquat. Org.* 119(1):1–16.
- Cunningham-Smith, P., D.E. Colbert, R.S. Wells and T. Speakman. 2006. Evaluation of human interactions with a wild bottlenose dolphin (*Tursiops truncatus*) near Sarasota Bay, Florida, and efforts to curtail the interactions. *Aquat. Mamm.* 32(3):346–356.
- DWH MMIQT [Deepwater Horizon Marine Mammal Injury Quantification Team]. 2015. Models and analyses for the quantification of injury to Gulf of Mexico cetaceans from the Deepwater Horizon Oil Spill, MM_TR.01_Schwacke_Quantification.of.Injury.to.GOM.Cetaceans. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, Florida 33140. PRBD Contribution #: PRBD-2020-02.
- DWH NRDAT [Deepwater Horizon Natural Resource Damage Assessment Trustees]. 2016. Deepwater Horizon oil spill: Final programmatic damage assessment and restoration plan and final programmatic environmental impact statement. <http://www.gulfspillrestoration.noaa.gov/restoration-planning/gulf-plan>
- Duignan, P.J., C. House, D.K. Odell, R.S. Wells, L.J. Hansen, M.T. Walsh, D.J. St. Aubin, B.K. Rima and J.R. Geraci. 1996. Morbillivirus infection in bottlenose dolphins: Evidence for recurrent epizootics in the western Atlantic and Gulf of Mexico. *Mar. Mamm. Sci.* 12:499–515.

- Fazioli, K.L., S. Hofmann and R.S. Wells. 2006. Use of Gulf of Mexico coastal waters by distinct assemblages of bottlenose dolphins (*Tursiops truncatus*). *Aquat. Mamm.* 32(2):212–222.
- Fulling, G.L., K.D. Mullin and C.W. Hubbard. 2003. Abundance and distribution of cetaceans in outer continental shelf waters of the U.S. Gulf of Mexico. *Fish. Bull.* 101:923–932.
- Garrison, L.P., J. Ortega-Ortiz and G. Rappucci. 2021. Abundance of coastal and continental shelf stocks of common bottlenose and Atlantic spotted dolphins in the northern Gulf of Mexico: 2017–2018. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, Florida 33140. PRD-2021-01. 25pp.
- Gasparini, N.M. and B. Yuill. 2020. High water: Prolonged flooding on the deltaic Mississippi River. *Eos* (101). 20 March 2020. <https://doi.org/10.1029/2020EO141465>
- Hansen, L.J. (ed). 1992. Report on investigation of 1990 Gulf of Mexico bottlenose dolphin strandings. NOAA-NMFS-SEFSC Contribution MIA-92/93-21. NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Irvine, A.B., M.D. Scott, R.S. Wells and J.H. Kaufmann. 1981. Movements and activities of the Atlantic bottlenose dolphin, *Tursiops truncatus*, near Sarasota, Florida. *Fish. Bull. U.S.* 79:671–688.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. Pages 108–189 in: S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake and L. Thomas (eds). *Advanced distance sampling*. Oxford University Press, New York. 434pp.
- Laake, J., D. Borchers, L. Thomas, D. Miller and J. Bishop. 2020. Package ‘mrds’: Mark-recapture distance sampling. Version 2.2.3. <http://github.com/DistanceDevelopment/mrds/>
- Lahvis, G.P., R.S. Wells, D.W. Kuehl, J.L. Stewart, H.L. Rhinehart and C.S. Via. 1995. Decreased lymphocyte responses in free-ranging bottlenose dolphins (*Tursiops truncatus*) are associated with increased concentrations of PCB’s and DDT in peripheral blood. *Environ. Health Perspect.* 103:67–72.
- Lipscomb, T.P. 1993. Some answers to questions about morbillivirus. Pages 4–5. in: R.A. Blaylock, B. Mase and D.K. Odell (eds). *Strandings*, Vol. 2, No. 3, SEFSC Miami Laboratory, Miami, Florida. 7pp.
- Lipscomb, T.P., S. Kennedy, D. Moffet and B.K. Ford. 1994. Morbilliviral disease in an Atlantic bottlenose dolphin (*Tursiops truncatus*) from the Gulf of Mexico. *J. Wildl. Dis.* 30(4):572–576.
- Litz, J.A., M.A. Baran, S.R. Bowen-Stevens, R.H. Carmichael, K.M. Colegrove, L.P. Garrison, S.E. Fire, E.M. Fougères, R. Hardy, S. Holmes, W. Jones, B.E. Mase-Guthrie, D.K. Odell, P.E. Rosel, J.T. Saliki, D.K. Shannon, S.F. Shippee, S.M. Smith, E.M. Stratton, M.C. Tumlin, H.R. Whitehead, G.A.J. Worthy and T.K. Rowles. 2014. Review of historical unusual mortality events (UMEs) in the Gulf of Mexico (1990–2009): Providing context for the complex and long-lasting northern Gulf of Mexico cetacean UME. *Dis. Aquat. Organ.* 112:161–175.
- Lynn, S.K. and B. Würsig. 2002. Summer movement patterns of bottlenose dolphins in a Texas bay. *G. Mex. Sci.* 20(1):25–37.
- Mathers, A.N., B.M. Deacy and J.K. Carlson. 2016. Characterization of catch and bycatch in the state gillnet fisheries of Alabama, Mississippi, and Louisiana, 2012–2015. NOAA Tech. Memo. NMFS-SEFSC-691. 35pp.
- Mathers, A.N., B.M. Deacy, H.E. Moncrief-Cox and J.K. Carlson. 2020. Catch and bycatch in U.S. Southeast gillnet fisheries, 2018. NOAA Tech. Memo. NMFS-SEFSC-743. 15pp.
- Maze-Foley, K. and L.P. Garrison. 2016. Serious injury determinations for small cetaceans off the southeast U.S. coast, 2007–2011. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2016-04. 16pp.
- Maze-Foley, K., B.L. Byrd, S.C. Horstman and J.R. Powell. 2019. Analysis of stranding data to support estimates of mortality and serious injury in common bottlenose dolphin (*Tursiops truncatus truncatus*) stock assessments for the Atlantic Ocean and Gulf of Mexico. NOAA Tech. Memo. NMFS-SEFSC-742. 42pp. <https://repository.library.noaa.gov/view/noaa/23151>
- Mullin, K.D., R.R. Lohoefer, W. Hoggard, C.L. Roden and C.M. Rogers. 1990. Abundance of bottlenose dolphins, *Tursiops truncatus*, in the coastal Gulf of Mexico. *Northeast Gulf Sci.* 11(2):113–122.
- NMFS [National Marine Fisheries Service]. 2016. Guidelines for preparing stock assessment reports pursuant to the 1994 amendments to the MMPA. NMFS Instruction 02-204-01, February 22, 2016. <http://www.nmfs.noaa.gov/op/pds/index.html>
- Peltier, H., W. Dabin, P. Daniel, O. Van Canneyt, G. Dorémus, M. Huon and V. Ridoux. 2012. The significance of stranding data as indicators of cetacean populations at sea: modelling the drift of cetacean carcasses. *Ecol. Indicators* 18:278–290.

- Powell, J.R. and R.S. Wells. 2011. Recreational fishing depredation and associated behaviors involving common bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Mar. Mamm. Sci.* 27(1):111–129.
- Rabalais, N.N., R.E. Turner and W.J. Wiseman, Jr. 1999. Hypoxia in the northern Gulf of Mexico: Linkage with the Mississippi River. Pages 297–322 *in*: H. Kumpf, K. Steidinger and K. Sherman (eds). *The Gulf of Mexico large marine ecosystem*. Blackwell Science, Oxford, UK.
- Read, A.J. 2008. The looming crisis: Interactions between marine mammals and fisheries. *J. Mammal.* 89(3):541–548.
- Rosel, P.E., L.A. Wilcox, C. Sinclair, T.R. Speakman, M.C. Tumlin, J.A. Litz and E.S. Zolman. 2017. Genetic assignment to stock of stranded common bottlenose dolphins in southeastern Louisiana after the *Deepwater Horizon* oil spill. *Endang. Species Res.* 33:221–234.
- Samuels, A. and L. Bejder. 2004. Chronic interactions between humans and free-ranging bottlenose dolphins near Panama City Beach, Florida, USA. *J. Cetacean Res. Manage.* 6:69–77.
- Schwacke, L.H., E.O. Voit, L.J. Hansen, R.S. Wells, G.B. Mitchum, A.A. Hohn and P.A. Fair. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the Southeast United States coast. *Environ. Toxicol. Chem.* 21:2752–2764.
- Schwacke, L.H., C.R. Smith, F.I. Townsend, R.S. Wells, L.B. Hart, B.C. Balmer, T.K. Collier, S. De Guise, M.M. Fry, L.J. Guillette, S.V. Lamb, S.M. Lane, W.E. McFee, N.J. Place, M.C. Tumlin, G.M. Ylitalo, E.S. Zolman and T.K. Rowles. 2014. Health of bottlenose dolphins (*Tursiops truncatus*) in Barataria Bay, Louisiana following the *Deepwater Horizon* oil spill. *Environ. Sci. Technol.* 48(1):93–103.
- Schwacke, L.H., L. Thomas, R.S. Wells, W.E. McFee, A.A. Hohn, K.D. Mullin, E.S. Zolman, B.M. Quigley, T.K. Rowles and J.H. Schwacke. 2017. Quantifying injury to common bottlenose dolphins from the *Deepwater Horizon* oil spill using an age-, sex- and class-structured population model. *Endanger. Spec. Res.* 33:265–279.
- Scott, G.P. 1990. Management-oriented research on bottlenose dolphins by the Southeast Fisheries Center. Pages. 623-639 *in*: S. Leatherwood and R.R. Reeves (eds). *The bottlenose dolphin*. Academic Press, San Diego, CA. 653pp.
- Sellas, A.B., R.S. Wells and P.E. Rosel. 2005. Mitochondrial and nuclear DNA analyses reveal fine scale geographic structure in bottlenose dolphins (*Tursiops truncatus*) in the Gulf of Mexico. *Conserv. Genet.* 6:715–728.
- Soldevilla, M.S., L.P. Garrison, E. Scott-Denton and J.M. Nance. 2015. Estimation of marine mammal bycatch mortality in the Gulf of Mexico shrimp otter trawl fishery. NOAA Tech. Memo. NMFS-SEFSC-672. 70pp.
- Soldevilla, M.S., L.P. Garrison, E. Scott-Denton and R.A. Hart. 2016. Estimated bycatch mortality of marine mammals in the Gulf of Mexico shrimp otter trawl fishery during 2012 to 2014. NOAA Tech. Memo. NMFS-SEFSC-697. 47pp.
- Soldevilla, M.S., L.P. Garrison, E. Scott-Denton and J. Primrose. 2021. Estimated bycatch mortality of marine mammals in the Gulf of Mexico shrimp otter trawl fishery during 2015 to 2019. NOAA Tech. Memo. NMFS-SEFSC-749. 78pp.
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. *Mar. Mamm. Sci.* 23(1):157–175.
- Vail, C.S. 2016. An overview of increasing incidents of bottlenose dolphin harassment in the Gulf of Mexico and possible solutions. *Front. Mar. Sci.* 3:110.
- Varanasi, U., K.L. Tilbury, D.W. Brown, M.M. Krahn, C.A. Wigren, R.C. Clark and S.L. Chan. 1992. Pages 56–86. *in*: L.J. Hansen (ed). *Report on investigation of 1990 Gulf of Mexico bottlenose dolphin strandings*, Southeast Fisheries Science Center Contribution MIA-92/93-21. 219pp.
- Venn-Watson, S., K.M. Colegrove, J. Litz, M. Kinsel, K. Terio, J. Saliki, S. Fire, R. Carmichael, C. Chevis, W. Hatchett, J. Pitchford, M. Tumlin, C. Field, S. Smith, R. Ewing, D. Fauquier, G. Lovewell, H. Whitehead, D. Rotstein, W. McFee, E. Fougères and T. Rowles. 2015. Adrenal gland and lung lesions in Gulf of Mexico common bottlenose dolphins (*Tursiops truncatus*) found dead following the *Deepwater Horizon* Oil Spill. *PLoS ONE* 10(5):e0126538.
- Vollmer, N.L. 2011. Population structure of common bottlenose dolphins in coastal and offshore waters of the Gulf of Mexico revealed by genetic and environmental analyses. Ph.D. Dissertation from University of Louisiana at Lafayette. 420pp.
- Vollmer, N.L., and P.E. Rosel. 2017. Fine-scale population structure of common bottlenose dolphins (*Tursiops truncatus*) in offshore and coastal waters of the US Gulf of Mexico. *Mar. Biol.* 164:160.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3–5, Seattle, WA. NOAA Tech. Memo. NMFS-OPR-12. 93pp.

- Wells, R.S., V. Tornero, A. Borrell, A. Aguilar, T.K. Rowles, H.L. Rhinehart, S. Hofmann, W.M. Jarman, A.A. Hohn and J.C. Sweeney. 2005. Integrating life history and reproductive success data to examine potential relationships with organochlorine compounds for bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Sci. Total Environ.* 349:106–119.
- Wells, R.S., J.B. Allen, G. Lovewell, J. Gorzelany, R.E. Delynn, D.A. Fauquier and N.B. Barros. 2015. Carcass-recovery rates for resident bottlenose dolphins in Sarasota Bay, Florida. *Mar. Mamm. Sci.* 31(1):355–368.
- Zollett, E.A. and A.J. Read. 2006. Depredation of catch by bottlenose dolphins (*Tursiops truncatus*) in the Florida king mackerel (*Scomberomorus cavalla*) troll fishery. *U.S. Fish. Bull.* 104:343–349.