# HARBOR PORPOISE (*Phocoena phocoena*): Gulf of Maine/Bay of Fundy Stock

## STOCK DEFINITION AND GEOGRAPHIC RANGE

This stock is found in U.S. and Canadian Atlantic waters. The distribution of harbor porpoises has been documented by sighting surveys, satellite telemetry data, passive acoustic monitoring, strandings and takes reported by NMFS observers in the Sea Sampling Programs. During summer (July to September), harbor porpoises are concentrated in the northern Gulf of Maine, southern Bay of Fundy and around the southern tip of Nova Scotia, generally in waters less than 150 m deep (Gaskin 1977; Kraus et al. 1983; Palka 1995), with lower densities in the upper Bay of Fundy and on Georges Bank (Palka 2000). During fall (October-December) and spring (April–June), harbor porpoises are widely dispersed from New Jersey to Maine, with lower densities farther north and south. During winter (January to March), intermediate densities of harbor porpoises can be found in waters off New Jersey to North Carolina, and lower densities are found in waters off New York to New Brunswick. Canada. In non-summer months they have been seen from the coastline to deep waters (>1800 m; Westgate et al. 1998), although the majority are found over the continental shelf. Passive acoustic monitoring detected harbor porpoises regularly during the period January-May offshore of Maryland (Wingfield et al. 2017). There does not appear to be a temporally coordinated migration or a specific migratory route to and from the Bay of Fundy region. However, during the fall, several satellite-tagged harbor porpoises did favor the waters around the 92-m isobath, which is consistent with observations of high rates of incidental catches in this depth range (Read and Westgate 1997). There were two stranding records from Florida during the 1980s (Smithsonian strandings database) and one in 2003 (NE Regional Office/NMFS strandings and entanglement database).




Figure 1. Distribution of harbor porpoises from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1995, 1998, 1999, 2002, 2004, 2006, 2007, 2008, 2010, 2011 and 2016 and portions of DFO's 2007 TNASS and 2016 NAISS surveys. Isobaths are the 100m, 200m, 1000m, and 4000m depth contours. Circle symbols represent shipboard sightings and squares are aerial sightings.

Gaskin (1984, 1992) proposed that there were four separate populations in the western North Atlantic: the Gulf of Maine/Bay of Fundy, Gulf of St. Lawrence, Newfoundland, and Greenland populations. Analyses involving mtDNA (Wang *et al.* 1996; Rosel *et al.* 1999a, 1999b), organochlorine contaminants (Westgate *et al.* 1997; Westgate and Tolley 1999), heavy metals (Johnston 1995), and life history parameters (Read and Hohn 1995) support Gaskin's proposal. Genetic studies using mitochondrial DNA (Rosel *et al.* 1999a) and contaminant studies using total PCBs

(Westgate and Tolley 1999) indicate that the Gulf of Maine/Bay of Fundy females were distinct from females from the other populations in the Northwest Atlantic. Gulf of Maine/Bay of Fundy males were distinct from Newfoundland and Greenland males, but not from Gulf of St. Lawrence males according to studies comparing mtDNA (Palka *et al.* 1996; Rosel *et al.* 1999a) and CHLORs, DDTs, PCBs and CHBs (Westgate and Tolley 1999). Nuclear microsatellite markers have also been applied to samples from these four populations, but this analysis failed to detect significant population sub-division in either sex (Rosel *et al.* 1999a). These patterns may be indicative of female philopatry coupled with dispersal of males. Both mitochondrial DNA and microsatellite analyses indicate that the Gulf of Maine/Bay of Fundy stock is not the sole contributor to the aggregation of porpoises found off the mid-Atlantic states during winter (Rosel *et al.* 1999a; Hiltunen 2006). Mixed-stock analyses using twelve microsatellite loci in both Bayesian and likelihood frameworks indicate that the Gulf of Maine/Bay of Fundy is the largest contributor (~60%), followed by Newfoundland (~25%) and then the Gulf of St. Lawrence (~12%), with Greenland making a small contribution (<3%). For Greenland, the lower confidence interval of the likelihood analysis includes zero. For the Bayesian analysis, the lower 2.5% posterior quantiles include zero for both Greenland and the Gulf of St. Lawrence. Intervals that reach zero provide the possibility that these populations contribute no animals to the mid-Atlantic aggregation.

This report follows Gaskin's hypothesis on harbor porpoise stock structure in the western North Atlantic, where the Gulf of Maine and Bay of Fundy harbor porpoises are recognized as a single management stock separate from harbor porpoise populations in the Gulf of St. Lawrence, Newfoundland, and Greenland. It is unlikely that the Gulf of Maine/Bay of Fundy harbor porpoise stock contains multiple demographically independent populations (Rosel *et al.* 1999a; Hiltunen 2006), but a comparison of samples from the Scotian shelf to the Gulf of Maine has not yet been made.

# POPULATION SIZE

The best current abundance estimate of the Gulf of Maine/Bay of Fundy harbor porpoise stock is the sum of the 2016 NEFSC and Department of Fisheries and Oceans Canada (DFO) surveys: 95,543 (CV=0.31; Table 1). Because the survey areas did not overlap, the estimates from the two surveys were added together and the CVs pooled using a delta method to produce a species abundance estimate for the stock area. A key uncertainty in the population size estimate is the precision and accuracy of the availability bias correction factor that was applied. More information on the spatio-temporal variability of the animals' dive profile is needed.

## **Earlier Abundance Estimates**

Please see Appendix IV for a summary of abundance estimates, including earlier estimates and survey descriptions. As recommended in the GAMMS II Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable for the determination of the current PBR.

# **Recent Surveys and Abundance Estimates**

An abundance estimate of 75,079 (CV=0.38) harbor porpoises was generated from a U.S. shipboard and aerial survey conducted during 27 June–28 September 2016 (Table 1; Palka 2020) in a region covering 425,192 km². The aerial portion included 11,782 km of tracklines that were over waters north of New Jersey from the coastline to the 100-m depth contour, throughout the U.S. waters. The shipboard portion included 4,351 km of tracklines that were in waters offshore of central Virginia to Massachusetts (waters that were deeper than the 100-m depth contour out to beyond the outer limit of the U.S. EEZ). Both sighting platforms used a two-team data collection procedure, which allows estimation of abundance to correct for perception bias of the detected species (Laake and Borchers 2004). The estimates were also corrected for availability bias.

An abundance estimate of 20,464 (CV=0.39) harbor porpoises from the Canadian Bay of Fundy/Scotian shelf region was generated from an aerial survey conducted by the Department of Fisheries and Oceans, Canada (DFO). The entire survey covered Atlantic Canadian shelf and shelf break waters extending from the northern tip of Labrador to the U.S border off southern Nova Scotia in August and September of 2016 (Lawson and Gosselin 2018). A total of 29,123 km were flown over the Gulf of St. Lawrence/Bay of Fundy/Scotian Shelf strata using two Cessna Skymaster 337s and 21,037 km were flown over the Newfound/Labrador strata using a DeHavilland Twin Otter. The harbor porpoise estimate was derived from the Skymaster data using single team multi-covariate distance sampling with left truncation (to accommodate the obscured area under the plane) where size-bias was also investigated. The Otter-based perception bias correction, which used double platform mark-recapture methods, was applied. An availability bias correction factor, which was based on published records of the cetaceans' surface intervals, was also applied.

Table 1. Summary of recent abundance estimates for the Gulf of Maine/Bay of Fundy harbor porpoise (Phocoena phocoena phocoena) by month, year, and area covered during each abundance survey and the resulting abundance estimate (Nest) and coefficient of variation (CV). The estimate considered best is in bold font.

Month/Year	Area	Nest	CV
Jun-Sep 2016	Central Virginia to Maine	75,079	0.38
Aug-Sep 2016	Gulf of St. Lawrence/Bay of Fundy/Scotian Shelf	20,464	0.39
Jun-Sep 2016	Central Virginia to Gulf of St. Lawrence/Bay of Fundy/Scotian Shelf - COMBINED	95,543	0.31

## **Minimum Population Estimate**

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for harbor porpoises is 95,543 (CV=0.31). The minimum population estimate for the Gulf of Maine/Bay of Fundy harbor porpoise is 74,034 (Table 2).

# **Current Population Trend**

A trend analysis has not been conducted for this stock. The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long survey interval. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV>0.30) remains below 80% (alpha=0.30) unless surveys are conducted on an annual basis (Taylor *et al.* 2007).

# CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Several attempts have been made to estimate potential population growth rates. Barlow and Boveng (1991), who used a re-scaled human life table, estimated the upper bound of the annual potential growth rate to be 9.4%. Woodley and Read (1991) used a re-scaled Himalayan tahr life table to estimate a likely annual growth rate of 4%. In an attempt to estimate a potential population growth rate that incorporates many of the uncertainties in survivorship and reproduction, Caswell *et al.* (1998) used a Monte Carlo method to calculate a probability distribution of growth rates. The median potential annual rate of increase was approximately 10%, with a 90% confidence interval of 3–15%. This analysis underscored the considerable uncertainty that exists regarding the potential rate of increase in this population. Moore and Read (2008) conducted a Bayesian population modeling analysis to estimate the potential population growth of harbor porpoise in the absence of bycatch mortality. Their method used fertility data, in combination with age-at-death data from stranded animals and animals taken in gillnets, and was applied under two scenarios to correct for possible data bias associated with observed bycatch of calves. Demographic parameter estimates were 'model averaged' across these scenarios. The Bayesian posterior median estimate for potential natural growth rate was 0.046. This last, most recent, value will be the one used for the purpose of this assessment.

Key uncertainties in the estimate of the maximum net productivity rate for this stock were discussed in Moore and Read (2008), which included the assumption that the age structure is stable, and the lack of data to estimate the probability of survivorship to maximum age. The authors considered the effects of these uncertainties on the estimated potential natural growth rate to be minimal.

# POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 74,034. The maximum productivity rate for this stock is 0.046. The recovery factor is 0.5 because stock's status relative to Optimum Sustainable Population (OSP) is unknown and the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the Gulf of Maine/Bay of Fundy harbor porpoise is 851 (Table 2).

Table 2. Best and minimum abundance estimates for the Gulf of Maine/Bay of Fundy harbor porpoise (Phocoena phocoena) with Maximum Productivity Rate (Rmax), Recovery Factor (Fr) and PBR.

Nest	CV	Nmin	Fr	Rmax	PBR
95,543	0.31	74,034	0.5	0.046	851

## ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual estimated average human-caused mortality and serious injury is 163 harbor porpoises per year (CV=0.13) from U.S. fisheries using observer data and an annual average of 1.6 animals from non-fishery stranding records (Table 3). Canadian bycatch information is not available.

Table 3. Total annual estimated average human-caused mortality and serious injury for the Gulf of Maine/Bay of

Fundy harbor porpoise (Phocoena phocoena phocoena).

Years	Source	Annual Avg.	CV
2015–2019	U.S. fisheries using observer data	163	0.13
2015–2019	Non-fishery human caused stranding mortalities	0.6	-
	TOTAL	164	-

A key uncertainty is the potential that the observer coverage in the Mid-Atlantic gillnet fishery may not be representative of the fishery during all times and places, since the observer coverage was relatively low for some times and areas (0.02–0.10). The effect of this is unknown. Another key uncertainty is that mortalities and serious injuries in Canadian waters are largely unquantified. There are no major known sources of unquantifiable human-caused mortality or serious injury for the U.S. waters within the Gulf of Maine/Bay of Fundy harbor porpoise stock's habitat.

## **United States**

### Northeast Sink Gillnet

Harbor porpoise bycatch in the northern Gulf of Maine occurs primarily from June to September, while in the southern Gulf of Maine and south of New England, bycatch occurs from January to May and September to December. Annual bycatch is estimated using ratio estimator techniques that account for the use of pingers (Orphanides and Hatch 2017; Orphanides 2019, 2020, 2021; Precoda and Orphanides 2022). See Table 4 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

# **Mid-Atlantic Gillnet**

Harbor porpoise bycatch in Mid-Atlantic waters occurs primarily from December to May in waters off New Jersey and less frequently in other waters ranging farther south, from New Jersey to North Carolina. Annual bycatch is estimated using ratio estimator techniques (Orphanides and Hatch 2017; Orphanides 2019, 2020, 2021; Precoda and Orphanides 2022). See Table 4 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

# **Northeast Bottom Trawl**

Since 1989, harbor porpoise mortalities have been observed in the northeast bottom trawl fishery, but many of these were not attributable to this fishery because decomposed animals are presumed to have been dead prior to being taken by the trawl. Those infrequently caught freshly dead harbor porpoises have been caught during January to April on Georges Bank or in the southern Gulf of Maine. Fishery-related bycatch rates were estimated using an annual stratified ratio-estimator (Lyssikatos and Chavez-Rosales 2022). See Table 4 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

# Canada

No current estimates exist, but harbor porpoise interactions have been documented in the Bay of Fundy sink gillnet fishery and in herring weirs between the years 1998–2001 in the lower Bay of Fundy demersal gillnet fishery (Trippel and Shepherd 2004). That fishery has declined since 2001 and it is assumed bycatch is very small, if any (H. Stone, Department of Fisheries and Oceans Canada, pers. comm.).

Table 4. From observer program data, summary of the incidental mortality of Gulf of Maine/Bay of Fundy harbor porpoise (Phocoena phocoena) by commercial fishery including the years sampled, the type of data used, the annual observer coverage, the mortalities and serious injuries recorded by on-board observers, the estimated

annual serious injury and mortality, the estimated CV of the annual mortality, and the mean annual combined

mortality with its CV.

Fishery	Years	Data Type <sup>a</sup>	Observer Coverage <sup>b</sup>	Obs. Serious Injury <sup>c</sup>	Obs. Mortality	Est. Serious Injury <sup>c</sup>	Est. Mortality	Est. Combined Mortality	Est. CVs	Mean Combined Annual Mortality
Northeast Sink Gillnet	2015 2016 2017 2018 2019	Obs. Data, Trip Logbook, Allocated Dealer Data	0.14 0.10 0.12 0.11 0.12	0 0 1 0	23 11 18 9 33	0 0 7 0	177 125 129 92 195	177 125 136 92 195	0.28 0.34 0.28 0.52 0.23	145 (0.14)
Mid- Atlantic Gillnet	2015 2016 2017 2018 2019	Obs. Data, Weighout	0.06 0.08 0.09 0.09 0.13	0 0 0 0	2 2 1 0 2	0 0 0 0	33 23 9.1 0 13	33 23 9.1 0 13	1.16 0.64 0.95 0 0.5	16 (0.68)
Northeast Bottom Trawl	2015 2016 2017 2018 2019	Obs. Data, Weighout	0.19 0.12 0.12 0.12 0.16	0 0 0 0	0 0 0 0 2	0 0 0 0	0 0 0 0 11	0 0 0 0 11	0 0 0 0 0.63	2.2 (.63)
TOTAL									163 (0.13)	

a. Observer data (Obs. Data) are used to measure bycatch rates and the data are collected within the Northeast Fisheries Observer Program. NEFSC collects Weighout (Weighout) landings data that are used as a measure of total effort for the U.S. gillnet fisheries. Mandatory vessel trip report (VTR; Trip Logbook) data are used to determine the spatial distribution of fishing effort in the Northeast sink gillnet fishery.

## Other Mortality

## **United States**

Recent harbor porpoise strandings on the U.S. Atlantic coast are documented in Table 5 (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 17 November 2020). Of the 417 stranding mortalities reported during this time period, 17 were coded as having signs of human interaction. Of these, 3 were deemed fishery interactions (assumend to be subsumed in the extrapolated fishery bycatch estimates) and 1 was a vessel strike. Most of the remaining Human Interaction (HI) cases were harassment, unlikely to have contributed to the stranding or post-mortem interactions. In only 3 cases were the non-fishery human interactions likely to have been contributing factors in the animal's mortality.

Stranding data underestimate the extent of mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

Table 5. Harbor porpoise (Phocoena phocoena phocoena) reported strandings along the U.S. and Canadian Atlantic coast, 2015–2019.

Area	2015	2016	2017	2018	2019	Total
Maine <sup>a, b, e</sup>	2	5	8	8	7	30
New Hampshire	0	1	2	2	6	11
Massachusetts <sup>a, b, d, e, h</sup>	18	8	29	13	68	136
Rhode Island b, d	2	2	0	0	2	6
Connecticut	0	0	0	0	1	1
New York a, d	3	1	12	2	13	31
New Jersey a, c, d	2	5	14	5	6	32
Delaware	0	0	6	0	3	9

b. Observer coverage for the U.S. Northeast and mid-Atlantic coastal gillnet fisheries is based on tons of fish landed. Northeast bottom trawl fishery coverages are ratios based on trips.

c. Serious injuries were evaluated for the 2015-2019 period and include both at-sea monitor and traditional observer data (Josephson et al. 2022).

Maryland	0	0	2	0	6	8
Virginia <sup>c</sup>	3	2	5	1	6	17
North Carolina <sup>b</sup>	14	1	1	3	13	32
TOTAL U.S.	44	25	79	34	131	313
Nova Scotia/Prince Edward Islandf	13	16	22	20	30	101
Newfoundland and New Brunswick <sup>g</sup>	2	0	0	0	1	3
GRAND TOTAL	59	41	101	54	162	417

a. In 2016, one animal in Maine and one animal in New Jersey were responded to and released alive. Ten animals were released alive in 2017, 6 of them in Massachusetts, 2 in Maine and 2 in New York.

#### Canada

Whales and dolphins stranded on the coast of Nova Scotia, New Brunswick and Prince Edward Island are recorded by the Marine Animal Response Society and the Nova Scotia Stranding Network. See Table 3 for details.

Harbor porpoises stranded on the coasts of Newfoundland and Labrador are reported by the Newfoundland and Labrador Whale Release and Strandings Program (Ledwell and Huntington 2015, 2017, 2018, 2019, 2020; Table 5).

## **HABITAT ISSUES**

In U.S. waters, harbor porpoise are mostly found in nearshore areas and inland waters, including bays, tidal areas, and river mouths. As a result, in addition to fishery bycatch, harbor porpoise are vulnerable to contaminants, such as PCBs (Hall *et al.* 2006), ship traffic (Oakley *et al.* 2017; Terhune 2015) and physical modifications resulting from urban and industrial development activities such as construction of docks and other over-water structures, dredging (Todd *et al.* 2015), installation of offshore windfarms (Carstensen *et al.* 2006; Brandt *et al.* 2011; Teilmann and Carstensen 2012; Dähne *et al.* 2013; Benjamins *et al.* 2017), seismic surveys and other sources of anthropogenic noise (Lucke *et al.* 2009).

Climate-related changes in spatial distribution and abundance, including poleward and depth shifts, have been documented in and predicted for a range of plankton species and commercially important fish stocks (Nye *et al.* 2009; Head *et al.* 2010; Pinsky *et al.* 2013; Poloczanska *et al.* 2013; Hare *et al.* 2016; Grieve *et al.* 2017; Morley *et al.* 2018) and cetacean species (e.g., MacLeod 2009; Sousa *et al.* 2019). There is uncertainty in how, if at all, the distribution and population size of this species will respond to these changes and how the ecological shifts will affect human impacts to the species.

# STATUS OF STOCK

Harbor porpoise in the Gulf of Maine/Bay of Fundy stock are not listed as threatened or endangered under the Endangered Species Act, and this stock is not considered strategic under the MMPA. The total U.S. fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The status of harbor porpoises, relative to OSP, in the U.S. Atlantic EEZ is unknown. Population trends for this species have not been investigated.

## REFERENCES CITED

Barlow, J. and P. Boveng. 1991. Modeling age-specific mortality for marine mammal populations. Mar. Mamm. Sci. 7: 50–65.

Benjamins, S., N. van Geel, G. Hastie, J. Elliott and B. Wilson. 2017. Harbour porpoise distribution can vary at small spatiotemporal scales in energetic habitats. Deep-Sea Res. II. http://dx.doi.org/10.1016/j.dsr2.2016.07.002i

b. Two HI cases in 2015: 1 in Rhode Island and 1 in North Carolina

c. Two HI cases in 2016: 1 in New Jersey and 1 in Virginia. The Virginia case was coded as a fishery interaction, and the New Jersey case was alive animal relocation.

d. Seven HI cases in 2017: 2 in Maine were released alive and another was a neonate with an infected laceration that required euthanization. One dead HI animal in Massachusetts was coded as a fishery interaction and another HI animal was released alive. One HI animal in New York was released alive and one dead animal in New Jersey had evidence of vessel interaction.

e. Two HI cases in 2018; both in Massachusetts. One was coded as a fishery interaction.

f. Data supplied by Nova Scotia Marine Animal Response Society (pers. comm.). One of the 2015 animals a suspected fishery interaction.

g. See Ledwell and Huntington (2015, 2017, 2018, 2019, 2020).

h. Three Massachusetts stranding mortalities in 2019 were classified as non-fishery human interaction.

- Brandt M.J., A. Diederichs, K. Betke and G. Nehls. 2011. Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. Mar Ecol Prog Ser 421:205–216. https://doi.org/10.3354/meps08888
- Carstensen J., O.D. Henriksen and J. Teilmann. 2006. Impacts of offshore wind farm construction on harbour porpoises: Acoustic monitoring of echolocation activity using porpoise detectors (T-PODs). Mar Ecol Prog Ser. 321:295–308.
- Caswell, H., S. Brault, A.J. Read and T.D. Smith. 1998. Harbor porpoise and fisheries: An uncertainty analysis of incidental mortality. Ecol. Appl. 8(4): 1226–1238.
- Dähne, M., A. Gilles, K. Lucke, V. Peschko, S. Adler, K. Krugel, J. Sundermeyer and U. Siebert. 2013. Effects of pile-driving on harbour porpoises (*Phocoena phocoena*) at the first offshore wind farm in Germany. Environ. Res. Lett. 8: 1–15.
- Gaskin, D.E. 1977. Harbour porpoise, *Phocoena phocoena* (L.), in the western approaches to the Bay of Fundy 1969–75. Rep. Int. Whal. Comm. 27:487–492.
- Gaskin, D.E. 1984. The harbor porpoise *Phocoena phocoena* (L.): Regional populations, status, and information on direct and indirect catches. Rep. Int. Whal. Comm. 34: 569–586.
- Gaskin, D.E. 1992. The status of the harbour porpoise. Can. Field-Nat. 106:36-54.
- Grieve, B.D., J.A. Hare and V.S. Saba. 2017. Projecting the effects of climate change on *Calanus finmarchicus* distribution within the US Northeast continental shelf. Sci. Rep. 7:6264.
- Hall, A.J., K. Hugunin, R. Deaville, R.J. Law, C.R. Allchin and P.D. Jepson. 2006. The risk of infection from polychlorinates biphenyl exposure in the harbor porpoise (*Phocoena phocoena*): A case-control approach. Environmental Health Perspectives 114(5):704–711.
- Hare, J.A., W.E. Morrison, M.W. Nelson, M.M. Stachura, E.J. Teeters, R.B. Griffis, M.A. Alexander, J.D. Scott, L. Alade, R.J. Bell, A.S. Chute, K.L. Curti, T.H. Curtis, D. Kurcheis, J.F. Kocik, S.M. Lucey, C.T. McCandless, L.M. Milke, D.E. Richardson, E. Robillard, H.J. Walsh, M.C. McManus, K.E. Maranick and C.A. Griswold. 2016. A vulnerability assessment of fish and invertebrates to climate change on the Northeast U.S. continental shelf. PLoS ONE. 11 (2016) e0146756. https://doi.org/10.1371/journal.pone.0146756.s014
- Head, E.J.H. and P. Pepin. 2010. Spatial and inter-decadal variability in plankton abundance and composition in the Northwest Atlantic (1958–2006). J. Plankton Res. 32:1633–1648.
- Hiltunen, K.H. 2006. Mixed-stock analysis of harbor porpoises (*Phocoena phocoena*) along the U.S. mid-Atlantic coast using microsatellite DNA markers. M.Sc. Thesis. The College of Charleston, Charleston, SC. 92pp.
- Johnston, D.W. 1995. Spatial and temporal differences in heavy metal concentrations in the tissues of harbour porpoises (*Phocoena phocoena* L.) from the western North Atlantic. M.Sc. Thesis. University of Guelph, Guelph, Ontario, Canada. 152pp.
- Josephson, E., F. Wenzel and M.C. Lyssikatos. 2022. Serious injury determinations for small cetaceans and pinnipeds caught in commercial fisheries off the northeast U.S. coast, 2015–2019. Northeast Fish. Sci. Cent. Ref. Doc. 22-03. 26pp.
- Kraus, S.D., J.H. Prescott and G.S. Stone. 1983. Harbor porpoise, *Phocoena phocoena*, in the U.S. coastal waters off the Gulf of Maine: A survey to determine seasonal distribution and abundance. NMFS. NA82FAC00027 22pp.
- Kraus, S.D., A.J. Read, A. Solow, K. Baldwin, T. Spradlin, E. Anderson and J. Williamson. 1997. Acoustic alarms reduce porpoise mortality. Nature 388(6642): 525.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. Pages 108-189 *in*: Advanced distance sampling. S. T. Buckland, D. R. Andersen, K. P. Burnham, J. L. Laake and L. Thomas (eds). Oxford University Press, New York.
- Lawson J. and J-F. Gosselin. 2018. Estimates of cetacean abundance from the 2016 NAISS aerial surveys of eastern Canadian waters, with a comparison to estimates from the 2007 TNASS. NAMMCO SC/25/AE/09.
- Ledwell, W. and J. Huntington. 2015. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles, strandings, ice entrapments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2015. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 22pp.
- Ledwell, W. and J. Huntington. 2017. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles and strandings and harassments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2016. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 23pp.

- Ledwell, W. and J. Huntington. 2018. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles, strandings, ice entrapments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2017. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 24pp.
- Ledwell, W. and J. Huntington. 2019. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles, strandings, ice entrapments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2018. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 24pp.
- Ledwell, W., J. Huntington, E. Sacrey and C. Landry. 2020. Entanglements in Fishing Gear and Strandings reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2019. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 24pp.
- Lucke, K., U. Siebert, P.A. Lepper and M-A. Blanchet. 2009. Temporary shift in masking hearing thresholds in a harbor porpoise (*Phocoena phocoena*) after exposure to seismic airgun stimuli. J. Acoust. Soc. Am. 125(6):4060–4070.
- Lyssikatos, M. S. Chavez-Rosales. 2022. Estimation of cetacean and pinniped bycatch in northeast and mid-Atlantic bottom trawl fisheries, 2015–2019. NOAA Tech. Memo. NMFS-NE-281.
- MacLeod, C.D. 2009. Global climate change, range changes and potential implications for the conservation of marine cetaceans: A review and synthesis. Endang. Species Res. 7:125–136.
- Moore, J.E. and A.J. Read. 2008. A Bayesian uncertainty analysis of cetacean demography and bycatch mortality using age-at-death data. Ecol. Appl. 18(8):1914–1931.
- Morley, J.W., R.L. Selden, R.J. Latour, T.L. Frolicher, R.J. Seagraves and M.L. Pinsky. 2018. Projecting shifts in thermal habitat for 686 species on the North American continental shelf. PLoS ONE 13(5):e0196127.
- NMFS [National Marine Fisheries Service]. 1992. Harbor porpoise in Eastern North America: Status and Research Needs. Results of a scientific workshop held May 5–8, 1992 at NEFSC, Woods Hole, MA, USA. Northeast Fish. Sci. Cent. Ref. Doc., 92-06. National Marine Fisheries Service. 28pp.
- Nye, J., J. Link, J. Hare and W. Overholtz. 2009. Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf. Mar. Ecol. Prog. Ser. 393:111–129.
- Oakley, J.A., A.T. Williams and T. Thomas. 2017. Reactions of harbor porpoise (*Phocoena phocoena*) to vessel traffic in the coastal waters of South West Wales, UK. Ocean & Coastal Management 138:158–169.
- Orphanides, C.D. 2019. Estimates of cetacean and pinniped bycatch during 2016 in the New England and Mid-Atlantic Sink Gillnet fisheries. Northeast Fish Sci Cent Ref Doc 19-04; 12pp. https://www.fisheries.noaa.gov/resource/publication-database/marine-mammal-mortality-and-serious-injury-reports
- Orphanides, C.D. 2020. Estimates of cetacean and pinniped bycatch during 2017 in the New England and Mid-Atlantic Sink Gillnet fisheries. Northeast Fish Sci Cent Ref Doc. 20-03 16pp. https://www.fisheries.noaa.gov/resource/publication-database/marine-mammal-mortality-and-serious-injury-reports
- Orphanides, C.D. 2021. Estimates of cetacean and pinniped bycatch during 2018 in the New England and Mid-Atlantic Sink Gillnet fisheries. Northeast Fish Sci Cent Ref Doc. 21-01. 16pp.
- Orphanides, C.D. and Hatch, J. 2017. Estimates of cetacean and pinniped bycatch during 2015 in the New England and Mid-Atlantic Sink Gillnet fisheries. Northeast Fish Sci Cent Ref Doc. 17-18. 21pp. https://www.fisheries.noaa.gov/resource/publication-database/marine-mammal-mortality-and-serious-injury-reports
- Precoda, K. and C.D. Orphanides. 2022. Estimates of cetacean and pinniped bycatch in the 2019 New England sink and mid-Atlantic Gillnet fisheries. Northeast Fish. Sci. Cent. Ref. Doc. 22-05. 21pp.
- Palka, D. 2020. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2016 line transect surveys conducted by the Northeast Fisheries Science Center. Northeast Fish. Sci. Cent. Ref. Doc. 20-05.
- Palka, D. 1995. Influences on spatial patterns of Gulf of Maine harbor porpoises. Pages 69–75 *in*: A.S. Blix, L. Walloe and O. Ulltang (eds). Whales, Seals, Fish and Man. Elsevier Science. Amsterdam.
- Palka, D.L., A.J. Read, A.J. Westgate and D.W. Johnston. 1996. Summary of current knowledge of harbour porpoises in US and Canadian Atlantic waters. Rep. Int. Whal. Comm. 46:559–565.
- Palka, D. 2000. Abundance of the Gulf of Maine/Bay of Fundy harbor porpoise based on shipboard and aerial surveys during 1999. Northeast Fish. Sci. Cent. Ref. Doc. 00-07. 29pp.

- https://repository.library.noaa.gov/view/noaa/3290.
- Palka, D.L. 2012. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey. Northeast Fish. Sci. Cent. Ref. Doc. 12-29. 37pp. https://repository.library.noaa.gov/view/noaa/4312
- Pinsky, M.L., B. Worm, M.J. Fogarty, J.L. Sarmiento and S.A. Levin. 2013. Marine taxa track local climate velocities. Science 341:1239–1242.
- Polacheck, T. 1989. Harbor porpoises and the gillnet fishery. Oceanus 32(1):63–70.
- Poloczanska, E.S., C.J. Brown, W.J. Sydeman, W. Kiessling, D.S. Schoeman, P.J. Moore, K. Brander, J.F. Bruno, L.B. Buckley, M.T. Burrows, C.M. Duarte, B.S. Halpern, J. Holding, C.V. Kappel, M.I. O'Connor, J.M. Pandolfi, C. Parmesan, F. Schwing, S.A. Thompson and A.J. Richardson. 2013. Global imprint of climate change on marine life. Nat. Clim. Change 3:919–925.
- Read, A.J. and A.A. Hohn. 1995. Life in the fast lane: The life history of harbour porpoises from the Gulf of Maine. Mar. Mamm. Sci. 11(4):423–440.
- Read, A.J. and A.J. Westgate. 1997. Monitoring the movements of harbour porpoises (*Phocoena phocoena*) with satellite telemetry. Ma. Biol. 130:315–22.
- Rosel, P.E., S.C. France, J.Y. Wang and T.D. Kocher. 1999a. Genetic structure of harbour porpoise *Phocoena phocoena* populations in the northwest Atlantic based on mitochondrial and nuclear markers. Mol. Ecol. 8:S41-S54.
- Rosel, P.E., R. Tiedemann and M. Walton. 1999b. Genetic evidence for limited trans-Atlantic movements of the harbor porpoise *Phocoena phocoena*. Mar. Biol. 133:583–591.
- Sousa, A., F. Alves, A. Dinis, J. Bentz, M.J. Cruz and J.P. Nunes. 2019. How vulnerable are cetaceans to climate change? Developing and testing a new index. Ecol. Indic. 98:9–18.
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. Mar. Mamm. Sci. 23(1):157–175.
- Teilmann, J. and J. Carstensen. 2012. Negative long term impacts on harbor porpoises from a large scale offshore wind farm in the Baltic—evidence of a slow recovery. Environ. Res, Lett. 7:045101.
- Terhune, J.M. 2015. Harbour porpoise presence near oil tankers. J. Canadian Ac. 43(3). 2pp.
- Thomas L., J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0. Release 2. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. http://distancesampling.org/Distance/
- Todd, V.L.G., I.B. Todd, J.C. Gardiner, E.C.N. Morrin, N.A. MacPherson, N.A. DiMarzio and F. Thomsen. 2015. A review of impacts of marine dredging activities on marine mammals. ICES J. of Mar. Sci. 72(2):328–340.
- Trippel, E.A. and T.D. Shepherd. 2004. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery from 1998–2001. Department of Fisheries and Oceans. Ottawa, Ontario. DFO Research Document 2004/2521 iv + 33 pp. http://www.fmap.ca/ramweb/papers-total/Trippel\_Shepherd\_2004.pdf
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3–5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93pp. https://repository.library.noaa.gov/view/noaa/15963
- Wang, J.Y., D.E. Gaskin and B.N. White. 1996. Mitochondrial DNA analysis of harbour porpoise, *Phocoena phocoena*, subpopulations in North American waters. Can. J. Fish. Aquat. Sci 53:1632–45.
- Westgate, A.J., D.C.G. Muir, D.E. Gaskin and M.C.S. Kingsley. 1997. Concentrations and accumulation patterns of organochlorine contaminants in the blubber of harbour porpoises, *Phocoena phocoena*, from the coast of Newfoundland, the Gulf of St. Lawrence and the Bay of Fundy/Gulf of Maine. Envir. Pollut. 95:105–119.
- Westgate, A.J., A.J. Read, T.M. Cox, T.D. Schofield, B.R. Whitaker and K.E. Anderson. 1998. Monitoring a rehabilitated harbor porpoise using satellite telemetry. Mar. Mamm. Sci. 14(3):599–604.
- Westgate, A.J. and K.A. Tolley. 1999. Geographical differences in organochlorine contaminants in harbour porpoises *Phocoena phocoena* from the western North Atlantic. Mar. Ecol. Prog. Ser. 177:255–268.
- Wingfield, J.E., M. O'Brien, V. Lyubchich, J.J. Roberts, P.N. Halpin, A.N. Rice and H. Bailey. 2017. Year-round spatiotemporal distribution of harbor porpoises within and around the Maryland wind energy area. PLoS ONE. 12(5):e0176653. https://doi.org/10.1371/journal.pone.0176653
- Woodley, T.H. and A.J. Read 1991. Potential rates of increase of a harbor porpoise (*Phocoena phocoena*) population subjected to incidental mortality in commercial fisheries. Can. J. Fish. Aquat. Sci 48:2429–35.