

LONG-FINNED PILOT WHALE (*Globicephala melas melas*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two species of pilot whales in the western Atlantic—the long-finned pilot whale, *Globicephala melas melas*, and the short-finned pilot whale, *G. macrorhynchus*. These species are difficult to differentiate at sea and cannot be reliably visually identified during either abundance surveys or observations of fishery mortality without high-quality photographs (Rone and Pace 2012); therefore, the ability to separately assess the two species in U.S. Atlantic waters is complex and requires additional information on seasonal spatial distribution. The long-finned pilot whale is distributed from North Carolina to North Africa (and the Mediterranean) and north to Iceland, Greenland and the Barents Sea (Sergeant 1962; Leatherwood *et al.* 1976; Abend 1993; Bloch *et al.* 1993; Abend and Smith 1999). The stock structure of the North Atlantic population is uncertain (ICES 1993; Fullard *et al.* 2000). Morphometric (Bloch and Lastein 1993) and genetic (Siemann 1994; Fullard *et al.* 2000) studies have provided little support for stock separation across the Atlantic (Fullard *et al.* 2000). However, Fullard *et al.* (2000) have proposed a stock structure that is related to sea-surface temperature: 1) a cold-water population west of the Labrador/North Atlantic current, and 2) a warm-water population that extends across the Atlantic in the Gulf Stream.

In U.S. Atlantic waters, pilot whales (*Globicephala* sp.) are distributed principally along the continental shelf edge off the northeastern U.S. coast in winter and early spring (CETAP 1982; Payne and Heinemann 1993; Abend and Smith 1999; Hamazaki 2002). In late spring, pilot whales move onto Georges Bank and into the Gulf of Maine and more northern waters, and remain in these areas through late autumn (CETAP 1982; Payne and Heinemann 1993). Pilot whales tend to occupy areas of high relief or submerged banks. They are also associated with the Gulf Stream wall and thermal fronts along the continental shelf




Figure 1. Distribution of long-finned (open symbols), short-finned (black symbols), and possibly mixed (gray symbols; could be either species) pilot whale sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004, 2006, 2007, 2011, and 2016 and the Department of Fisheries and Oceans Canada 2007 TNASS and 2016 NAISS surveys. The inferred distribution of the two species is preliminary and is valid for June–August only. Isobaths are the 1000-m and 3000-m depth contours. The U.S. EEZ is also displayed in green.

edge (Waring *et al.* 1992). Long-finned and short-finned pilot whales overlap spatially along the mid-Atlantic shelf break between Delaware and the southern flank of Georges Bank (Payne and Heinemann 1993; Rone and Pace 2012). Long-finned pilot whales have occasionally been observed stranded as far south as Florida, and short-finned pilot whales have occasionally been observed stranded as far north as Massachusetts. The exact latitudinal ranges of the two species therefore remain uncertain, although south of Cape Hatteras, most pilot whale sightings are expected to be short-finned pilot whales, while north of ~42°N most pilot whale sightings are expected to be long-finned pilot whales (Figure 1; Garrison and Rosel 2017).

POPULATION SIZE

The best available estimate for long-finned pilot whales in the western North Atlantic is 39,215 (CV=0.30; Table 1; Garrison 2020; Palka 2020; Lawson and Gosselin 2018). This estimate is the sum of the estimates generated from the northeast U.S. summer 2016 surveys covering U.S. waters from central Virginia to Maine and the Department of Fisheries and Oceans Canada summer 2016 survey covering Canadian waters from the U.S. to Labrador. Because the survey areas did not overlap, the estimates from the two surveys were added together and the CVs pooled using a delta method to produce a species abundance estimate for the stock area. The 2016 estimate is larger than those from 2011 because the 2016 estimate is derived from a survey area extending from Newfoundland to Florida, which is about 1,300,000 km² larger than the 2011 survey area. In addition, the 2016 survey estimates in U.S. waters were corrected for availability bias (due to diving behavior), whereas the 2011 estimates were not corrected. These survey data have been combined with an analysis of the spatial distribution of the 2 species based on genetic analyses of biopsy samples to derive separate abundance estimates (Garrison and Rosel 2017).

Key uncertainties in the population size estimate include the uncertain separation between the short-finned and long-finned pilot whales; the small negative bias due to the lack of an abundance estimate in the region between the US and the Newfoundland/Labrador survey area; and the uncertainty due to the unknown precision and accuracy of the availability bias correction factor that was applied.

Earlier Abundance Estimates

Please see appendix IV for a summary of abundance estimates including earlier estimates and survey descriptions. As recommended in the GAMMS II Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable for the determination of the current PBR. Due to changes in survey methodology, these historical data should not be used to make comparisons with more current estimates.

Recent Surveys and Abundance Estimates for *Globicephala* sp.

Abundance estimates of 8,166 (CV=0.31) and 25,114 (CV=0.27) *Globicephala* sp. were generated from vessel surveys conducted in the northeast and southeast U.S., respectively, during the summer of 2016. The Northeast survey was conducted during 27 June–25 August and consisted of 5,354 km of on-effort trackline. The majority of the survey was conducted in waters north of 38°N latitude and included trackline along the shelf break and offshore to the U.S. EEZ. Pilot whale sightings were concentrated along the shelf-break between the 1,000-m and 2,000-m isobaths and along Georges Bank (NMFS 2017). The Southeast vessel survey covered waters from Central Florida to approximately 38°N latitude between the 100-m isobaths and the U.S. EEZ during 30 June–19 August. A total of 4,399 km of trackline was covered on effort. Pilot whales were observed in high densities along the shelf-break between Cape Hatteras and New Jersey and also in waters further offshore in the mid-Atlantic and off the coast of Florida (NMFS 2017; Garrison and Palka 2018). Both the Northeast and Southeast surveys utilized two visual teams and an independent observer approach to estimate detection probability on the trackline (Laake and Borchers 2004). Mark-recapture distance sampling was used to estimate abundance. A logistic regression model was used to estimate the abundance of long-finned pilot whales from these surveys. For the northeast survey, this resulted in an abundance estimate of 10,997 (CV=0.51) long-finned pilot whales. In the southeast, the model indicated that this survey included habitats expected to exclusively contain short-finned pilot whales so no estimate for long-finned pilot whales was generated.

An abundance estimate of 28,218 (CV=0.36) long-finned pilot whales from the Newfoundland/Labrador region was generated from an aerial survey conducted by the Department of Fisheries and Oceans, Canada (DFO). This survey covered Atlantic Canadian shelf and shelf break waters extending from the northern tip of Labrador to the U.S. border off southern Nova Scotia in August and September of 2016 (Lawson and Gosselin 2018). A total of 29,123 km were flown over the Gulf of St. Lawrence/Bay of Fundy/Scotian Shelf stratum using two Cessna Skymaster 337s and 21,037 km were flown over the Newfoundland/Labrador stratum using a DeHavilland Twin Otter. The Newfoundland estimate was derived from the Twin Otter data using two-team mark-recapture multi-covariate distance sampling

methods. An availability bias correction factor, which was based on the cetaceans' surface intervals, was also applied. The Gulf of St. Lawrence/Bay of Fundy/Scotian Shelf survey detected 10 pilot whale groups, however, no abundance estimate was produced.

Spatial Distribution and Abundance Estimates for *Globicephala melas*

Biopsy samples from pilot whales were collected during summer months (June–August) from South Carolina to the southern flank of Georges Bank between 1998 and 2007. These samples were identified to species using phylogenetic analysis of mitochondrial DNA sequences. Stranded specimens that were morphologically identified to species were used to assign clades in the phylogeny to species and thereby identify all samples. The probability of a sample being from a long-finned (or short-finned) pilot whale was evaluated as a function of sea-surface temperature, latitude, and month using a logistic regression. This analysis indicated that the probability of a sample coming from a long-finned pilot whale was near 1 at water temperatures <22°C, and near 0 at temperatures >25°C. The probability of a long-finned pilot whale also increased with increasing latitude. Spatially, during summer months, this regression model predicted that all pilot whales observed in offshore waters near the Gulf Stream are most likely short-finned pilot whales. The area of overlap between the two species occurs primarily along the shelf break off the coast of New Jersey between 38°N and 40°N latitude (Garrison and Rosel 2017).

This model was used to partition the abundance estimates from surveys conducted during the summer of 2016. The sightings from the southeast shipboard surveys covering waters from Florida to New Jersey were predicted to consist entirely of short-finned pilot whales. The aerial portion of the northeast surveys covered the Gulf of Maine and the Bay of Fundy and surveys where the model predicted that only long-finned pilot whales would occur. The vessel portion of the northeast surveys recorded a mix of both species along the shelf break, and the sightings in offshore waters near the Gulf Stream were predicted to consist predominantly of short-finned pilot whales (Garrison and Rosel 2017).

Table 1. Summary of recent abundance estimates for the western North Atlantic long-finned pilot whale (*Globicephala melas melas*) by month, year, and area covered during each abundance survey, and resulting abundance estimate (*N_{est}*) and coefficient of variation (*CV*).

Month/Year	Area	N_{est}	CV
Jun–Aug 2016	Central Virginia to Lower Bay of Fundy	10,997	0.51
Aug–Sep 2016	Newfoundland/Labrador	28,218	0.36
Jun–Sep 2016	Central Virginia to Labrador - COMBINED	39,215	0.30

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for western North Atlantic long-finned pilot whales is 39,215 animals (CV=0.30). The minimum population estimate for long-finned pilot whales is 30,627.

Current Population Trend

A trend analysis has not been conducted for this stock. The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long survey interval. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV>0.30) remains below 80% (alpha=0.30) unless surveys are conducted on an annual basis (Taylor *et al.* 2007). There is current work to standardize the strata-specific previous abundance estimates to consistently represent the same regions and include appropriate corrections for perception and availability bias. These standardized abundance estimates will be used in state-space trend models that incorporate environmental factors that could potentially influence the process and observational errors for each stratum.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for long-finned pilot whales is 30,627. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor is 0.5 because this stock is of unknown status relative to optimum sustainable population (OSP) and the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the western North Atlantic long-finned pilot whale is 306 (Table 2).

Table 2. Best and minimum abundance estimates for western North Atlantic long-finned pilot whale (*Globicephala melas melas*) with Maximum Productivity Rate (R_{max}), Recovery Factor (F_r) and PBR.

Nest	CV	Nmin	F_r	R_{max}	PBR
39,215	0.30	30,627	0.5	0.04	306

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Total annual observed average fishery-related mortality or serious injury during 2015–2019 was 9.0 long-finned pilot whales (CV=0.4; Table 3). In bottom trawls and mid-water trawls and in the gillnet fisheries, mortalities were more generally observed north of 40°N latitude and in areas expected to have only long-finned pilot whales. Takes in these fisheries were therefore attributed to the long-finned pilot whales. Takes in the pelagic longline fishery were partitioned according to a logistic regression model (Garrison and Rosel 2017).

Fishery Information

Detailed fishery information is reported in Appendix III.

Earlier Interactions

See Appendix V for more information on historical takes.

United States

Longline

During 2015–2019, pilot whale interactions (all serious injuries) were apportioned between the short-finned and long-finned pilot whale stocks according to a logistic regression model (Garrison and Rosel 2017). See Table 3 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

Northeast Bottom Trawl

Fishery-related bycatch rates for years 2015–2019 were estimated using an annual stratified ratio-estimator (Lyssikatos and Chavez-Rosales 2022). See Table 3 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

Northeast Mid-Water Trawl (Including Pair Trawl)

Three pilot whales were taken in the northeast mid-water trawl fishery in 2016. Using model-based predictions and at-sea identification, these takes have all been assigned as long-finned pilot whales. Expanded estimates of fishery mortality for 2015–2019 are not available, and so for those years the raw number is provided. See Table 3 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

Canada

Unknown numbers of long-finned pilot whales have been taken in Newfoundland, Labrador, Scotian shelf and Bay of Fundy groundfish gillnets; Atlantic Canada and Greenland salmon gillnets; and Atlantic Canada cod traps (Read 1994).

Table 3. Summary of the incidental mortality and serious injury of long-finned pilot whales (*Globicephala melas melas*) by U.S. commercial fisheries including the years sampled (Years), the type of data used (Data Type), the annual observer coverage coverage (Observer Coverage), the observed mortalities and serious injuries recorded by on-board observers, the estimated annual mortality and serious injury, the combined annual estimates of mortality and serious injury (Estimated Combined Mortality), the estimated CV of the combined estimates (Est. CVs) and the

mean of the combined estimates (CV in parentheses). These are minimum observed counts as expanded estimates are not available.

Fishery	Years	Data Type ^a	Observer Coverage ^b	Observed Serious Injury ^c	Observed Mortality	Estimated Serious Injury ^e	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Combined Annual Mortality
Northeast Bottom Trawl	2015	Obs. Data, Logbook	0.19	0	0	0	0	0	na	6.9 (0.51)
	2016		0.12	0	4	0	29	29	0.58	
	2017		0.12	0	0	0	0	0	na	
	2018		0.12	0	0	0	0	0	na	
	2019		0.16	0	1	0	5.39	5.39	0.88	
Northeast Mid-Water Trawl - Including Pair Trawl ^c	2015	Obs. Data, Dealer Data, VTR Data	0.08	0	0	0	0	0	na	0.6 (na)
	2016		0.27	0	3	0	3	3	na	
	2017		0.16	0	0	0	0	0	na	
	2018		0.14	0	0	0	0	0	na	
	2019		0.28	0	0	0	0	0	na	
Pelagic Longline Fishery	2015	Obs. Data, Logbook Data	0.12	1	0	2.2	0	2.2	0.49	1.5 (0.49)
	2016		0.15	1	0	1.1	0	1.1	1.0	
	2017		0.12	1	0	3.3	0	3.3	0.98	
	2018		0.10	1	0	0.4	0	0.4	0.93	
	2019		0.10	1	0	0.4	0	0.4	1.0	
TOTAL										9.0 (0.4)

a. Observer data (Obs. Data) are used to measure bycatch rates and the data are collected within the Northeast Fisheries Observer Program (NEFOP). NEFSC collects landings data (unallocated Dealer Data and Allocated Dealer Data) which are used as a measure of total landings. Mandatory Vessel Trip Reports (VTR; Trip Logbook) are used to determine the spatial distribution of landings and fishing effort. Total landings are used as a measure of total effort for the coastal gillnet fishery.

b. The observer coverages for the northeast sink gillnet fishery are ratios based on tons of fish landed. Northeast bottom trawl and northeast mid-water trawl fishery coverages are ratios based on trips.

c. Expanded estimates are not available for this fishery.

d. Serious injuries were evaluated for the period and include both at-sea monitor and traditional observer data (Josephson *et al.* 2022).

Other Mortality

Pilot whales have a propensity to mass strand throughout their range, but the role of human activity in these events is unknown. From 2015 to 2019, 7 long-finned pilot whales (*Globicephala melas melas*) were reported stranded between Maine and Florida, including the EEZ (Table 4; NOAA National Marine Mammal Health and Stranding Response Database, accessed 17 November 2020). None of the animals had indications of human interaction.

Table 4. Pilot whale (*Globicephala melas melas*) strandings along the Atlantic coast, 2015–2019. The level of technical expertise among stranding network personnel varies, and given the potential difficulty in correctly identifying stranded pilot whales to species, reports to specific species should be viewed with caution.

State	2015	2016	2017	2018	2019	Total
Nova Scotia ^a	21	12	12	3	2	50
Newfoundland and Labrador ^b	0	0	1	0	2	3
Maine ^c	0	1	1	3	0	5
Massachusetts	0	1	1	0	0	2
Total U.S.	0	2	2	3	0	7

a. Data supplied by Nova Scotia Marine Animal Response Society (pers. comm.).

b. See Ledwell and Huntington 2015, 2017, 2018, 2019, 2020.

c. 2016 animal released alive.

Stranding data probably underestimate the extent of human and fishery-related mortality and serious injury, particularly for offshore species such as pilot whales, because not all of the whales that die or are seriously injured in human interactions wash ashore, or, if they do, they are not all recovered (Peltier *et al.* 2012; Wells *et al.* 2015).

Additionally, not all carcasses will show evidence of human interaction, entanglement or other fishery-related interaction due to decomposition, scavenger damage, etc. (Byrd *et al.* 2014). Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of human interaction.

HABITAT ISSUES

The chronic impacts of contaminants (polychlorinated biphenyls [PCBs] and chlorinated pesticides [DDT, DDE, dieldrin, etc.]) on marine mammal reproduction and health are of concern (e.g., Schwacke *et al.* 2002; Jepson *et al.* 2016; Hall *et al.* 2018). Moderate levels of these contaminants have been found in pilot whale blubber (Taruski *et al.* 1975; Muir *et al.* 1988; Weisbrod *et al.* 2000). Weisbrod *et al.* (2000) examined polychlorinated biphenyl and chlorinated pesticide concentrations in bycaught and stranded pilot whales in the western North Atlantic. Contaminant levels were similar to or lower than levels found in other toothed whales in the western North Atlantic, perhaps because they are feeding further offshore than other species (Weisbrod *et al.* 2000). Dam and Bloch (2000) found very high PCB levels in long-finned pilot whales in the Faroes. Also, high levels of toxic metals (mercury, lead, cadmium) and selenium were measured in pilot whales harvested in the Faroe Island drive fishery (Nielsen *et al.* 2000). However, the population effect of the observed levels of such contaminants on this stock is unknown.

Anthropogenic sound in the world's oceans has been shown to affect marine mammals, with vessel traffic, seismic surveys, and active naval sonars being the main anthropogenic contributors to low- and mid-frequency noise in oceanic waters (e.g., Nowacek *et al.* 2015; Gomez *et al.* 2016; NMFS 2018). The long-term and population consequences of these impacts are less well-documented and likely vary by species and other factors. Impacts on marine mammal prey from sound are also possible (Carroll *et al.* 2017), but the duration and severity of any such prey effects on marine mammals are unknown.

Climate-related changes in spatial distribution and abundance, including poleward and depth shifts, have been documented in or predicted for plankton species and commercially important fish stocks (Nye *et al.* 2009; Pinsky *et al.* 2013; Poloczanska *et al.* 2013; Grieve *et al.* 2017; Morley *et al.* 2018) and cetacean species (e.g., MacLeod 2009; Sousa *et al.* 2019). There is uncertainty in how, if at all, the distribution and population size of this species will respond to these changes and how the ecological shifts will affect human impacts to the species.

STATUS OF STOCK

The long-finned pilot whale is not listed as threatened or endangered under the Endangered Species Act, and the western North Atlantic stock is not considered strategic under the MMPA because the mean annual human-caused mortality and serious injury does not exceed PBR. Total U.S. fishery-related mortality and serious injury for long-finned pilot whales is less than 10% of the calculated PBR and, therefore, can be considered to be insignificant and approaching zero mortality and serious injury rate. The status of this stock relative to OSP in the U.S. Atlantic EEZ is unknown. A population trend analysis for this stock has not been conducted.

Based on the low levels of uncertainty described in the above sections, it is expected these uncertainties will have little effect on the designation of the status of this stock.

REFERENCES CITED

- Abend, A. 1993. Long-finned pilot whale distribution and diet as determined from stable carbon and nitrogen ratio isotope tracers. M.Sc. Thesis, University of Massachusetts, Amherst, MA. 147pp.
- Abend, A.G. and T.D. Smith. 1999. Review of distribution of the long-finned pilot whale (*Globicephala melas*) in the North Atlantic and Mediterranean. NOAA Tech. Memo. NMFS-NE-117. 22pp.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73pp.
- Bloch, D. and L. Lastein. 1993. Morphometric segregation of long-finned pilot whales in eastern and western North Atlantic. *Ophelia* 38:55–68.
- Bloch, D., M. Zachariassen and P. Zachariassen. 1993. Some external characters of the long-finned pilot whale off Faroe Island and a comparison with the short-finned pilot whale. *Rep. Int. Whal. Comm. (Special Issue)* 14:117–135.
- Byrd, B.L., A.A. Hohn, G.N. Lovewell, K.M. Altman, S.G. Barco, A. Friedlaender, C.A. Harms, W.A. McLellan, K.T. Moore, P.E. Rosel and V.G. Thayer. 2014. Strandings illustrate marine mammal biodiversity and human impacts off the coast of North Carolina, USA. *Fish. Bull.* 112:1–23.

- Carroll, A.G., R. Przeslawski, A. Duncan, M. Gunning, B. Bruce. 2017. A critical review of the potential impacts of marine seismic surveys on fish & invertebrates. *Mar. Pollut. Bull.* 114:9–24.
- CETAP [Cetacean and Turtle Assessment Program]. 1982. A characterization of marine mammals and turtles in the mid- and North Atlantic areas of the U.S. outer continental shelf, final report. University of Rhode Island. #AA551-CT8-48. Bureau of Land Management, Washington, DC. 576pp.
- Dam, M. and D. Bloch. 2000. Screening of mercury and persistent organochlorine pollutants in long-finned pilot whale (*Globicephala melas*) in the Faroe Islands. *Mar. Poll. Bull.* 40:1090–1099.
- Fullard, K.J., G. Early, M.P. Heide-Jorgensen, D. Bloch, A. Rosing-Asvid and W. Amos 2000. Population structure of long-finned pilot whales in the North Atlantic: A correlation with sea surface temperature? *Mol. Ecol.* 9: 949–958.
- Garrison, L.P. and P.E. Rosel. 2017. Partitioning short-finned and long-finned pilot whale bycatch estimates using habitat and genetic information. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2016-17. 24pp.
- Garrison, L.P. 2020. Abundance of cetaceans along the southeast U.S. east coast from a summer 2016 vessel survey. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRD Contribution # PRD-2020-04. 17pp.
- Gomez, C., J.W. Lawson, A.J. Wright, A.D. Buren, D. Tollit and V. Lesage. 2016. A systematic review on the behavioural responses of wild marine mammals to noise: The disparity between science and policy. *Can. J. Zool.* 94:801–819.
- Grieve, B.D., J.A. Hare and V.S. Saba. 2017. Projecting the effects of climate change on *Calanus finmarchicus* distribution within the US Northeast continental shelf. *Sci. Rep.* 7:6264.
- Hamazaki, T. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, No. Carolina, USA to Nova Scotia, Canada). *Mar. Mamm. Sci.* 18(4):920–939.
- Hare, J.A., W.E. Morrison, M.W. Nelson, M.M. Stachura, E.J. Teeters, R.B. Griffis, M.A. Alexander, J.D. Scott, L. Alade, R.J. Bell, A.S. Chute, K.L. Curti, T.H. Curtis, D. Kurcheis, J.F. Kocik, S.M. Lucey, C.T. McCandless, L.M. Milke, D.E. Richardson, E. Robillard, H.J. Walsh, M.C. McManus, K.E. Maranick, C.A. Griswold. 2016. A vulnerability assessment of fish and invertebrates to climate change on the Northeast U.S. continental shelf. *PLoS ONE*. 11:e0146756. <https://doi.org/10.1371/journal.pone.0146756.s014>
- Head, E.J.H. and P. Pepin. 2010. Spatial and inter-decadal variability in plankton abundance and composition in the Northwest Atlantic (1958–2006). *J. Plankton Res.* 32:1633–1648.
- ICES [International Council for the Exploration of the Sea]. 1993. Report of the study group on long-finned pilot whales, Copenhagen, Denmark, ICES. 30 August–3 September 1993. *CM* 1993/N:5.
- Jepson, P.D., R. Deaville, J.L. Barber, A. Aguilar, A. Borrell, S. Murphy, J. Barry, A. Brownlow, J. Barnett, S. Berrow and A.A. Cunningham. 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.-U.K.* 6:18573.
- Josephson, E., F. Wenzel and M.C. Lyssikatos. 2022. Serious injury determinations for small cetaceans and pinnipeds caught in commercial fisheries off the northeast U.S. coast, 2015–2019. *Northeast Fish. Sci. Cent. Ref. Doc.* 22-03. 26pp.
- Hall, A.J., B.J. McConnell, L.J. Schwacke, G.M. Ylitalo, R. Williams and T.K. Rowles. 2018. Predicting the effects of polychlorinated biphenyls on cetacean populations through impacts on immunity and calf survival. *Environ. Pollut.* 233:407–418.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. Pages 108–189 *in*: Advanced distance sampling, S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake, and L. Thomas (eds). Oxford University Press, New York, New York.
- Lawson J. and J-F. Gosselin. 2018. Estimates of cetacean abundance from the 2016 NAISS aerial surveys of eastern Canadian waters, with a comparison to estimates from the 2007 TNASS. *NAMMCO SC/25/AE/09*
- Leatherwood, S., D.K. Caldwell and H.E. Winn. 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. *NOAA Tech. Rep. NMFS CIRC.396.* 176pp.
- Ledwell, W. and J. Huntington. 2015. Incidental entrapments in fishing gear and stranding reported to and responded to by the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2015. Report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 22pp.
- Ledwell, W. and J. Huntington. 2017. Incidental entrapments in fishing gear and stranding reported to and responded to by the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale

- Release and Strandings program during 2016. Report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 23pp.
- Ledwell, W. and J. Huntington. 2018. Incidental entrapments in fishing gear and stranding reported to and responded to by the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2017. Report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 23pp.
- Ledwell, W. and J. Huntington. 2019. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles, strandings, ice entrapments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2018. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 24pp.
- Ledwell, W., J. Huntington, E. Sacrey and C. Landry. 2020. Entanglements in Fishing Gear and Strandings reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2019. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 24pp.
- Lyssikatos, M.C. and S. Chavez-Rosales. 2022. Estimates of cetacean and pinniped bycatch in Northeast and mid-Atlantic bottom trawl fisheries, 2015–2019. *Northeast Fish. Sci. Cent. Ref. Doc. NMFS-NE 281*. DOI: 10.25923/6sj9-yw17
- MacLeod, C.D. 2009. Global climate change, range changes and potential implications for the conservation of marine cetaceans: A review and synthesis. *Endang. Species Res.* 7:125–136.
- Morley, J.W., R.L. Selden, R.J. Latour, T.L. Frolicher, R.J. Seagraves and M.L. Pinsky. 2018. Projecting shifts in thermal habitat for 686 species on the North American continental shelf. *PLoS ONE* 13(5):e0196127.
- Muir, D.C.G., R. Wagemann, N.P. Grift, R.J. Norstrom, M. Simon and J. Lien. 1988. Organochlorine chemical and heavy metal contaminants in white-beaked dolphins (*Lagenorhynchus albirostris*) and pilot whales (*Globicephala melaena*) from the coast of Newfoundland, Canada. *Arch. Environ. Contam. Toxicol.* 17(5): 613–629.
- Nielsen, J.B., F. Nielsen, P.-J. Jorgensen and P. Grandjean. 2000. Toxic metals and selenium in blood from pilot whales (*Globicephala melas*) and sperm whales (*Physeter catodon*). *Mar. Poll. Bull.* 40:348–35. <http://www.nefsc.noaa.gov/nefsc/publications/crd/crd0603/crd0603.pdf>
- Nowacek, D.P., C.W. Clark, D. Mann, P.J.O. Miller, H.C. Rosenbaum, J.S. Golden, M. Jasny, J. Kraska and B.L. Southall. 2015. Marine seismic surveys and ocean noise: time for coordinated and prudent planning. *Front. Ecol. Environ.* 13:378–386.
- Nye, J., J. Link, J. Hare and W. Overholtz. 2009. Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf. *Mar. Ecol. Prog. Ser.* 393:111–129.
- Palka, D. 2020. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2016 line transect surveys conducted by the Northeast Fisheries Science Center. *Northeast Fish. Sci. Cent. Ref. Doc.* 20-05.
- Payne, P.M. and D.W. Heinemann 1993. The distribution of pilot whales (*Globicephala* sp.) in shelf/shelf edge and slope waters of the northeastern United States, 1978–1988. *Rep. Int. Whal. Comm. (Special Issue)* 14:51–68.
- Peltier, H., W. Dabin, P. Daniel, O. Van Canneyt, G. Dorémus, M. Huon and V. Ridoux. 2012. The significance of stranding data as indicators of cetacean populations at sea: modelling the drift of cetacean carcasses. *Ecol. Indic.* 18:278–290.
- Pinsky, M.L., B. Worm, M.J. Fogarty, J.L. Sarmiento and S.A. Levin. 2013. Marine taxa track local climate velocities. *Science* 341:1239–1242.
- Poloczanska, E.S., C.J. Brown, W.J. Sydeman, W. Kiessling, D.S. Schoeman, P.J. Moore, K. Brander, J.F. Bruno, L.B. Buckley, M.T. Burrows, C.M. Duarte, B.S. Halpern, J. Holding, C.V. Kappel, M.I. O'Connor, J.M. Pandolfi, C. Parmesan, F. Schwing, S.A. Thompson and A.J. Richardson. 2013. Global imprint of climate change on marine life. *Nat. Clim. Change* 3:919–925.
- Read, A.J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest. *Atlantic. Rep. Int. Whal. Comm. (Special Issue)* 15:133–147.
- Rone, B.K. and R.M. Pace, III. 2012. A simple photograph-based approach for discriminating between free-ranging long-finned (*Globicephala melas*) and short-finned (*G. macrorhynchus*) pilot whales off the east coast of the United States. *Mar. Mamm. Sci.* 28(2):254–275.

- Schwacke, L.H., E.O. Voit, L.J. Hansen, R.S. Wells, G.B. Mitchum, A.A. Hohn and P.A. Fair. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the southeast United States coast. *Env. Toxic. Chem.* 21(12):2752–2764.
- Sergeant, D.E. 1962. The biology of the pilot or pothead whale (*Globicephala melaena* (Traill)) in Newfoundland waters. *Bull. Fish. Res. Bd. Can* 132:1–84.
- Siemann, L. 1994. Mitochondrial DNA sequence variation in North Atlantic long-finned pilot whales, *Globicephala melas*. Ph.D. Thesis, Massachusetts Institute of Technology/Woods Hole Oceanographic Institution.
- Sousa, A., F. Alves, A. Dinis, J. Bentz, M.J. Cruz and J.P. Nunes. 2019. How vulnerable are cetaceans to climate change? Developing and testing a new index. *Ecol. Indic.* 98:9–18.
- Taruski, A.G., C.E. Olney and H.E. Winn 1975. Chlorinated hydrocarbons in cetaceans. *J. Fish. Res. Brd. Can.* 32(11): 2205–2209.
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. *Mar. Mamm. Sci.* 23(1):157–175.
- Thomas, L., J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0. Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. <http://distancesampling.org/Distance/>
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3–5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93pp. <https://repository.library.noaa.gov/view/noaa/15963>
- Waring, G.T., C.P. Fairfield, C.M. Ruhsam and M. Sano. 1992. Cetaceans associated with Gulf Stream features off the northeastern USA Shelf. *ICES [Int. Counc. Explor. Sea] C.M.* 1992/N:12.
- Weisbrod, A.V., D. Shea, M.J. Moore and J.J. Stegeman. 2000. Bioaccumulation patterns of polychlorinated biphenyls and chlorinated pesticides in northwest Atlantic pilot whales. *Environ. Toxicol. Chem.* 19:667–677.
- Wells, R.S., J.B. Allen, G. Lovewell, J. Gorzelany, R.E. Delynn, D.A. Fauquier and N.B. Barros. 2015. Carcass-recovery rates for resident bottlenose dolphins in Sarasota Bay, Florida. *Mar. Mamm. Sci.* 31(1):355–368.