

## COMMON BOTTLENOSE DOLPHIN (*Tursiops truncatus truncatus*): Gulf of Mexico Northern Coastal Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

Common bottlenose dolphins inhabit coastal waters throughout the northern Gulf of Mexico (i.e., U.S. Gulf of Mexico) (Mullin *et al.* 1990). As a working hypothesis, it is assumed that the dolphins occupying habitats with dissimilar climatic, coastal and oceanographic characteristics might be restricted in their movements between habitats, and thus constitute separate stocks. Therefore, northern Gulf of Mexico coastal waters have been divided for management purposes into 3 stock areas: eastern, northern and western, with coastal waters defined as waters between the shore, barrier islands or presumed outer bay boundaries out to the 20-m isobath (Figure 1). The 20-m depth seaward boundary corresponds to survey strata (Scott 1990; Blaylock and Hoggard 1994; Fulling *et al.* 2003), and thus represents a management boundary rather than an ecological boundary. The Northern Coastal bottlenose dolphin stock area extends from 84°W longitude to the Mississippi River Delta. This region is characterized by a temperate climate, barrier islands, sand beaches, coastal marshes and marsh islands, and has a relatively high level of freshwater input. It is bordered on the east by an extensive area of coastal marsh and marsh islands typical of Florida's Apalachee Bay. Dolphins belonging to this stock are all expected to be of the coastal ecotype (Vollmer 2011).

This stock's boundaries abut other bottlenose dolphin stocks, namely the Continental Shelf Stock and several bay, sound and estuary stocks, and while individuals from different stocks may occasionally overlap, it is not thought that significant mixing or interbreeding occurs between them. Fazioli *et al.* (2006) conducted photo-identification surveys of coastal waters off Tampa Bay, Sarasota Bay and Lemon Bay,




Figure 1. Locations (circles) of common bottlenose dolphin groups sighted in coastal and continental shelf waters during aerial surveys conducted in spring, summer and fall of 2011 and in winter of 2012. Dark circles indicate groups within the boundaries of the Northern Coastal Stock. The 20-m and 200-m isobaths are shown.

Florida, over 14 months. They found coastal waters were inhabited by both 'inshore' and 'Gulf' dolphins but that the 2 types used coastal waters differently. Dolphins from the inshore communities were observed occasionally in Gulf near-shore waters adjacent to their inshore range, whereas 'Gulf' dolphins were found primarily in open Gulf of Mexico waters with some displaying seasonal variations in their use of the study area. The 'Gulf' dolphins did not show a preference for waters near passes as was seen for 'inshore' dolphins, but moved throughout the study area and made greater use of waters offshore of waters used by 'inshore' dolphins. During winter months abundance of 'Gulf' groups decreased while abundance for 'inshore' groups increased. These findings support an earlier report by Irvine *et al.* (1981) of increased use of pass and coastal waters by Sarasota Bay dolphins in winter. Seasonal movements of identified individuals and abundance indices suggest that part of the 'Gulf' dolphin community moves out of the study area during winter, but their destination is unknown. In a follow-up study, Sellas *et al.* (2005) examined genetic population subdivision in the study area of Fazioli *et al.* (2006), and found evidence of significant population structure among all areas on the basis of both mitochondrial DNA control region sequence data and 9 nuclear microsatellite loci. The Sellas *et al.* (2005) findings support the separate identification of bay, sound and

estuary stocks from those occurring in adjacent Gulf coastal waters, as suggested by Wells (1986).

Off Galveston, Texas, Beier (2001) reported an open population of individual dolphins in coastal waters, but several individual dolphins had been sighted previously by other researchers over a 10-year period. Some coastal animals may move relatively long distances alongshore. Two bottlenose dolphins previously seen in the South Padre Island area in Texas were seen in Matagorda Bay, 285 km north, in May 1992 and May 1993 (Lynn and Würsig 2002).

## POPULATION SIZE

The best abundance estimate available for the northern Gulf of Mexico Northern Coastal Stock of bottlenose dolphins is 7,185 (CV=0.21; Table 1). This estimate is from an inverse-variance weighted average of seasonal abundance estimates from aerial surveys conducted during spring 2011, summer 2011, fall 2011 and winter 2012.

### Earlier abundance estimates

Please see Appendix IV for a summary of abundance estimates, including earlier estimates and survey descriptions.

### Recent surveys and abundance estimates

The Southeast Fisheries Science Center conducted aerial surveys of continental shelf waters (shoreline to 200 m depth) along the U.S. Gulf of Mexico coast from the Florida Keys to the Texas/Mexico border during spring (March-April) 2011, summer (July-August) 2011, fall (October-November) 2011 and winter (January-February) 2012. The surveys were conducted along tracklines oriented perpendicular to the shoreline and spaced 20-30 km apart. The total survey effort varied during each survey due to weather conditions, but ranged between 13,500 – 15,600 km. Each of these surveys was conducted using a two-team approach to develop estimates of visibility bias using the independent observer approach with Distance analysis (Laake and Borchers 2004). A model for the probability of detection on the trackline as a function of sighting conditions (sea state, glare, water color, etc.) was developed using data across all 4 surveys. This model was then applied to detection probability functions specific to each survey to account for the probability of detection as a function of distance from the trackline and additional environmental covariates. A bootstrap resampling approach was used to estimate the variance of the estimates. The survey data were post-stratified into spatial boundaries corresponding to the defined boundaries of bottlenose dolphin stocks within the surveyed area. The abundance estimates for the Northern Coastal Stock of bottlenose dolphins were based upon tracklines and sightings in waters from the shoreline to the 20-m isobath and between the Mississippi River Delta and 84°W longitude, including waters of northern Chandeleur Sound. The seasonal abundance estimates for this stock were: spring – 15,831 (CV=0.38), summer – 6,792 (CV=0.28), fall – 2,384 (CV=0.38) and winter – 2,384 (CV=0.31). Due to the uncertainty in stock movements and apparent seasonal variability in the abundance of the stock, a weighted average of these seasonal estimates was taken where the weighting was the inverse of the CV. This approach weights estimates with higher precision more heavily in the final weighted mean. The resulting weighted mean and best estimate of abundance for the Northern Coastal Stock of bottlenose dolphins was 7,185 (CV=0.21).

Previous abundance estimates for the Northern and Eastern Coastal Stocks were derived from aerial surveys conducted during 17 July to 8 August 2007. Survey effort covered waters from the shoreline to 200m depth and was stratified such that the majority of effort was expended in the 0-20m depth range of the coastal stocks. The survey team consisted of an observer stationed at each of two forward bubble windows and a third observer stationed at a belly window that monitored the trackline. Surveys were typically flown during favorable sighting conditions at Beaufort sea state less than or equal to 3 (surface winds <10 knots). Abundance estimates were derived using Distance analysis including environmental covariates that had a significant influence on sighting probability (Buckland *et al.* 2001), but these estimates were not corrected for  $g(0)$  and are thus negatively biased. The resulting abundance estimate for the Northern Coastal Stock was 2,473 (CV=0.25).

Table 1. Summary of recent abundance estimates for the Northern Coastal Stock of bottlenose dolphins. Month, year and area covered during each abundance survey, and resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).			
Month/Year	Area	$N_{best}$	CV
July-Aug 2007	shoreline to 20 m, Northern Coastal Stock waters (Mississippi River Delta to 84°W longitude)	2,473	0.25

Spring, summer and fall 2011, winter 2012	shoreline to 20 m, Northern Coastal Stock waters (Mississippi River Delta to 84°W longitude)	7,185	0.21
---	--	-------	------

### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for the Northern Coastal Stock of bottlenose dolphins is 7,185 (CV=0.21). The minimum population estimate for the Northern Coastal Stock is 6,044 bottlenose dolphins.

### Current Population Trend

There are insufficient data to determine population trends for this stock. The abundance estimates for summer 2007 and summer 2012 are 2,473 (CV=0.25) and 6,792 (CV=0.28), respectively.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are not known for this stock. The maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

### POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate and a recovery factor (Wade and Angliss 1997). The minimum population size is 6,044. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.5 because the stock is of unknown status. PBR for the northern Gulf of Mexico Northern Coastal Stock of bottlenose dolphins is 60.

### ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual human-caused mortality and serious injury of the Northern Coastal Stock of bottlenose dolphins during 2008-2012 is unknown. Two mortalities were documented involving the Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line) fishery; however, it is not possible to estimate the total number of interactions or mortalities associated with the hook and line fishery since there is no systematic observer program.

### New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998; Andersen *et al.* 2008; NOAA 2012). NMFS defines serious injury as an “*injury that is more likely than not to result in mortality*”. Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

### Fisheries Information

The commercial fisheries which potentially could interact with the Northern Coastal Stock in the northern Gulf of Mexico are the Category II Southeastern U.S. Atlantic, Gulf of Mexico shrimp trawl, Southeastern U.S. Atlantic, Gulf of Mexico stone crab trap/pot, and Gulf of Mexico menhaden purse seine, and the Category III Gulf of Mexico blue crab trap/pot and Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line) fisheries (Appendix III).

### Hook and Line Fisheries

During 2008-2012, 2 mortalities involving hook and line gear entanglement or ingestion were documented for the Northern Coastal Stock. The mortalities occurred in 2012 and 2011. The mortalities were included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 [for 2008-2011 data] and 15 April 2013 [for 2012 data]) and are included in the stranding totals presented in Table 2.

### **Shrimp Trawl Fishery**

During 2008-2012, no interactions between bottlenose dolphins of the Northern Coastal Stock and the shrimp trawl fishery were observed. A voluntary observer program for the shrimp trawl fishery began in 1992 and became mandatory in 2007. A total of 5 bottlenose dolphin mortalities were observed in the shrimp trawl fishery during 2003, 2007, 2008, 2010 and 2011, and 1 serious injury was observed during 2012. The 2003 mortality occurred off the coast of Alabama and could have belonged to the Northern Coastal Stock or a bay, sound and estuary stock (Mobile Bay, Bonsecour Bay Stock or Mississippi Sound, Lake Borgne, Bay Boudreau Stock). During 1992-2007 the observer program recorded an additional 6 unidentified dolphins caught in a lazy line or turtle excluder device, and 1 of these animals, a mortality in 2001, belonged to the Northern Coastal Stock. The observer report indicated the animal may have already been decomposed, but this could not be confirmed because there was no necropsy.

### **Blue and Stone Crab Trap/Pot Fisheries**

There have been no reported mortalities or serious injuries involving trap/pot gear for the Northern Coastal Stock to date. However, mortalities and serious injuries have been reported for the Eastern Coastal Stock, Western Coastal Stock, and bay, sound and estuary stocks. Since there is no systematic observer program, it is not possible to estimate the total number of interactions or mortalities associated with crab traps/pots.

### **Menhaden Purse Seine Fishery**

During 2008-2012, no interactions between the Northern Coastal Stock and the menhaden purse seine fishery were documented. There is currently no observer program for the Gulf of Mexico menhaden purse seine fishery; however, recent interactions with bottlenose dolphins have been reported via two sources. First, during 2011, a pilot observer program operated from May through September, and observers documented 3 dolphins trapped within purse seine nets (within waters of the Western Coastal Stock and Mississippi Sound, Lake Borgne, Bay Boudreau Stock). All 3 were released alive without serious injury (Maze-Foley and Garrison in prep). Second, through the Marine Mammal Authorization Program (MMAP), there have been 13 self-reported incidental takes (all mortalities) of bottlenose dolphins in northern Gulf of Mexico coastal and estuarine waters by the menhaden purse seine fishery during 2000-2012. These takes likely affected the following stocks: Western Coastal Stock; Northern Coastal Stock; Mississippi Sound, Lake Borgne, Bay Boudreau Stock; and Mississippi River Delta Stock. Specific self-reported takes under the MMAP that might be attributed to the Northern Coastal Stock are as follows: one take of a single bottlenose dolphin was reported in Louisiana waters during 2001 that likely belonged to Mississippi River Delta Stock or Northern Coastal Stock; and during 2000, there was one reported take of a single bottlenose dolphin in Louisiana waters that likely belonged to Mississippi River Delta Stock or Northern Coastal Stock.

The menhaden purse seine fishery was observed to take 9 bottlenose dolphins (3 fatally) between 1992 and 1995 (NMFS unpublished data). During that period, there were 1,366 sets observed out of 26,097 total sets, which if extrapolated for all years suggests that as many as 172 bottlenose dolphins could have been taken in this fishery with up to 57 animals killed.

Without an ongoing observer program it is not possible to obtain statistically reliable information for this fishery on the number of sets annually, the incidental take and mortality rates, and the communities from which bottlenose dolphins are being taken.

### **Strandings**

A total of 90 bottlenose dolphins were found stranded in Northern Coastal Stock waters of the Gulf of Mexico from 2008 through 2012 (Table 2; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 and 15 April 2013). Evidence of human interactions (e.g., gear entanglement, mutilation, gunshot wounds) was detected for 5 of these dolphins. Bottlenose dolphins are known to become entangled in, or ingest recreational and commercial fishing gear (Wells and Scott 1994; Gorzelany 1998; Wells *et al.* 1998; Wells *et al.* 2008), and some are struck by vessels (Wells and Scott 1997; Wells *et al.* 2008).

There are a number of difficulties associated with the interpretation of stranding data. It is possible that some or all of the stranded dolphins may have been from a nearby bay, sound and estuary stock; however, the proportion of stranded dolphins belonging to another stock cannot be determined because of the difficulty of determining from where the stranded carcass originated. Stranding data probably underestimate the extent of human-related mortality and serious injury because not all of the dolphins that die or are seriously injured due to human interactions wash ashore, nor will all of those that do wash ashore necessarily show signs of fishery-interaction or other human interactions. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of human interaction, and the condition of the carcass if badly decomposed can inhibit the

interpretation of cause of death.

Since 1990, there have been 13 bottlenose dolphin die-offs or Unusual Mortality Events (UMEs) in the northern Gulf of Mexico, and 7 of these have occurred within the boundaries of the Northern Coastal Stock and may have affected the stock. 1) From January through May 1990, a total of 367 bottlenose dolphins stranded in the northern Gulf of Mexico. Overall this represented a two-fold increase in the prior maximum recorded strandings for the same period, but in some locations (i.e., Alabama) strandings were 10 times the average number. The cause of the 1990 mortality event could not be determined (Hansen 1992). 2) In 1993-1994 an UME of bottlenose dolphins likely caused by morbillivirus started in the Florida Panhandle and spread west with most of the mortalities occurring in Texas (Lipscomb 1993; Lipscomb *et al.* 1994). From February through April 1994, 220 bottlenose dolphins were found dead on Texas beaches, of which 67 occurred in a single 10-day period. 3) In 1996 an UME was declared for bottlenose dolphins in Mississippi when 27 bottlenose dolphins stranded during November and December. The cause was not determined, but a *Karenia brevis* (red tide) bloom was suspected to be responsible. 4) Between August 1999 and May 2000, 152 bottlenose dolphins died coincident with *K. brevis* blooms and fish kills in the Florida Panhandle (additional strandings included 3 Atlantic spotted dolphins, *Stenella frontalis*, 1 Risso's dolphin, *Grampus griseus*, 2 Blainville's beaked whales, *Mesoplodon densirostris*, and 4 unidentified dolphins). 5) In March and April 2004, in another Florida Panhandle UME possibly related to *K. brevis* blooms, 105 bottlenose dolphins and 2 unidentified dolphins stranded dead (NMFS 2004). Although there was no indication of a *K. brevis* bloom at the time, high levels of brevetoxin were found in the stomach contents of the stranded dolphins (Flewelling *et al.* 2005). 6) A separate UME was declared in the Florida Panhandle after elevated numbers of dolphin strandings occurred in association with a *K. brevis* bloom in September 2005. Dolphin strandings remained elevated through the spring of 2006 and brevetoxin was again detected in the tissues of some of the stranded dolphins. Between September 2005 and April 2006 when the event was officially declared over, a total of 90 bottlenose dolphin strandings occurred (plus strandings of 3 unidentified dolphins). 7) An UME was declared for cetaceans in the northern Gulf of Mexico beginning 1 February 2010; and, as of 2013, the event is still ongoing. It includes cetaceans that stranded prior to the Deepwater Horizon oil spill (see "Habitat Issues" below), during the spill, and after. During 2010, 14 animals from this stock were considered to be part of the UME, during 2011, 40 animals, and during 2012, 16 animals.

Table 2. Bottlenose dolphin strandings occurring in Northern Coastal Stock waters of the northern Gulf of Mexico from 2008 to 2012, as well as number of strandings for which evidence of human interaction (HI) was detected and number of strandings for which it could not be determined (CBD) if there was evidence of HI. Data are from the NOAA National Marine Mammal Health and Stranding Response Database (unpublished data, accessed 13 September 2012 [for 2008-2011 data] and 15 April 2013 [for 2012 data]). Please note HI does not necessarily mean the interaction caused the animal's death.

Stock	Category	2008	2009	2010	2011	2012	Total
Northern Coastal Stock	Total Stranded	7	8	18 <sup>a</sup>	40 <sup>b</sup>	17 <sup>c</sup>	90
	Human Interaction						
	---Yes	0	1	1	1 <sup>d</sup>	2 <sup>e</sup>	5
	---No	1	3	3	0	4	11
	---CBD	6	4	14	39	11	74

<sup>a</sup> This total includes 14 strandings that are part of the ongoing UME in the northern Gulf of Mexico.

<sup>b</sup> All of these strandings were part of the ongoing UME in the northern Gulf of Mexico.

<sup>c</sup> This total includes 16 strandings that are part of the ongoing UME in the northern Gulf of Mexico.

<sup>d</sup> This was an entanglement interaction (mortality) with recreational hook and line gear.

<sup>e</sup> Includes 1 entanglement interaction (mortality) with hook and line gear.

### Other Mortality

The problem of dolphin depredation of fishing gear is increasing in the Gulf of Mexico. There have been 4 recent cases of fishermen illegally "taking" dolphins due to dolphin depredation of recreational and commercial fishing gear. One recent case of a shrimp fisherman illegally "taking" a dolphin in Mississippi Sound occurred during summer 2012. In December 2013 the fisherman was convicted under the MMPA for knowingly shooting a dolphin with a shotgun while shrimping. A commercial fisherman was indicted in November 2008 for throwing pipe bombs at dolphins off Panama City, Florida, and charged in March 2009 for "taking" dolphins with an explosive device. In 2006 a charter boat fishing captain was charged under the MMPA for shooting at a dolphin that was

swimming around his catch in the Gulf of Mexico, off Panama City, Florida. In 2007 a second charter fishing boat captain was fined under the MMPA for shooting at a bottlenose dolphin that was attempting to remove a fish from his line in the Gulf of Mexico, off Orange Beach, Alabama. Feeding or provisioning of wild bottlenose dolphins has been documented in Florida, particularly near Panama City Beach in the Panhandle (Samuels and Bejder 2004) and south of Sarasota Bay (Cunningham-Smith *et al.* 2006; Powell and Wells 2011), and also in Texas near Corpus Christi (Bryant 1994). Feeding wild dolphins is defined under the MMPA as a form of 'take' because it can alter their natural behavior and increase their risk of injury or death. Nevertheless, a high rate of provisioning was observed near Panama City Beach in 1998 (Samuels and Bejder 2004), and provisioning has been observed south of Sarasota Bay since 1990 (Cunningham-Smith *et al.* 2006; Powell and Wells 2011). There are emerging questions regarding potential linkages between provisioning and depredation of recreational fishing gear and associated entanglement and ingestion of gear, which is increasing through much of Florida. During 2006, an estimated 2% of the long-term resident dolphins of Sarasota Bay died from ingestion of recreational fishing gear (Powell and Wells 2011).

Swimming with wild bottlenose dolphins has also been documented in Florida in Key West (Samuels and Engleby 2007) and near Panama City Beach (Samuels and Bejder 2004). Near Panama City Beach, Samuels and Bejder (2004) concluded that dolphins were amenable to swimmers due to illegal provisioning. Swimming with wild dolphins may cause harassment, and harassment is illegal under the MMPA.

## **HABITAT ISSUES**

The Deepwater Horizon (DWH) MC252 drilling platform, located approximately 50 miles southeast of the Mississippi River Delta in waters about 1500m deep, exploded on 20 April 2010. The rig sank, and over 87 days ~4.9 million barrels of oil were discharged from the wellhead until it was capped on 15 July 2010 (McNutt *et al.* 2012). During the response effort dispersants were applied extensively at the seafloor and at the sea surface (Lehr *et al.* 2010; OSAT 2010). In-situ burning, or controlled burning of oil at the surface, was also used extensively as a response tool (Lehr *et al.* 2010). The oil, dispersant and burn residue compounds present ecological concerns. The magnitude of this oil spill was unprecedented in U.S. history, causing impacts to wildlife, natural habitats and human communities along coastal areas from western Louisiana to the Florida Panhandle (NOAA 2011). It could be years before the entire scope of damage is ascertained (NOAA 2011).

Given the trajectory of the surface oil during the spill and the documented oiling of shoreline (Michel *et al.* 2013), it is likely the Northern Coastal Stock of bottlenose dolphins was exposed to oil during the event. A substantial number of beaches and wetlands along the Louisiana coast experienced heavy or moderate oiling (OSAT-2 2011; Michel *et al.* 2013). The heaviest oiling in Louisiana occurred west of the Mississippi River on the Mississippi Delta and in Barataria and Terrebonne Bays, and to the east of the river on the Chandeleur Islands. Some heavy to moderate oiling occurred on Alabama and Florida beaches, with the heaviest stretch occurring from Dauphin Island, Alabama, to Gulf Breeze, Florida. Light to trace oil was reported along the majority of Mississippi's mainland coast, from Gulf Breeze to Panama City, Florida, and outside of Atchafalaya and Vermilion Bays in western Louisiana. Heavy to light oiling occurred on Mississippi's barrier islands (Michel *et al.* 2013).

Shortly after the oil spill, the Natural Resource Damage Assessment (NRDA) process was initiated under the Oil Pollution Act of 1990. A variety of NRDA research studies are being conducted to determine potential impacts of the spill on marine mammals. These studies have focused on identifying the type, magnitude, severity, length and impact of oil exposure to oceanic, continental shelf, coastal and estuarine marine mammals. The research is ongoing. For coastal and estuarine dolphins, the NOAA-led efforts include: active surveillance to detect stranded animals in remote locations; aerial surveys to document the distribution, abundance, species and exposure of marine mammals and sea turtles relative to oil from DWH spill; assessment of sublethal and chronic health impacts on coastal and estuarine bottlenose dolphins in Barataria Bay, Louisiana, and a reference site in Sarasota Bay, Florida; and assessment of injuries to dolphin stocks in Barataria Bay and Chandeleur Sound, Louisiana, Mississippi Sound, and as a reference site, St. Joseph Bay, Florida.

Dolphins were observed with tar balls attached to them and seen swimming through oil slicks close to shore and inland bays. The effects of oil exposure on marine mammals depend on a number of factors including the type and mixture of chemicals involved, the amount, frequency and duration of exposure, the route of exposure (inhaled, ingested, absorbed, or external) and biomedical risk factors of the particular animal (Geraci 1990). In general, direct external contact with petroleum compounds or dispersants with skin may cause skin irritation, chemical burns and infections. Inhalation of volatile petroleum compounds or dispersants may irritate or injure the respiratory tract, which could lead to pneumonia or inflammation. Ingestion of petroleum compounds may cause injury to the gastrointestinal tract, which could affect an animal's ability to digest or absorb food. Absorption of petroleum compounds or dispersants may damage kidney, liver and brain function in addition to causing immune suppression

and anemia. Long term chronic effects such as lowered reproductive success and decreased survival may occur (Geraci 1990).

The nearshore habitat occupied by the 3 coastal stocks is adjacent to areas of high human population and in some areas, such as Tampa Bay, Florida, Galveston, Texas, and Mobile, Alabama, is highly industrialized. Concentrations of anthropogenic chemicals such as PCBs and DDT and its metabolites vary from site to site, and can reach levels of concern for bottlenose dolphin health and reproduction in the southeastern U.S. (Schwacke *et al.* 2002). PCB concentrations in 3 stranded dolphins sampled from the Eastern Coastal Stock area ranged from 16-46µg/g wet weight. Two stranded dolphins from the Northern Coastal Stock area had the highest levels of DDT derivatives of any of the bottlenose dolphin liver samples analyzed in conjunction with a 1990 mortality investigation conducted by NMFS (Varanasi *et al.* 1992). The significance of these findings is unclear, but there is some evidence that increased exposure to anthropogenic compounds may reduce immune function in bottlenose dolphins (Lahvis *et al.* 1995), or impact reproduction through increased first-born calf mortality (Wells *et al.* 2005). Concentrations of chlorinated hydrocarbons and metals were relatively low in most of the bottlenose dolphins examined in conjunction with an anomalous mortality event in Texas bays in 1990; however, some had concentrations at levels of possible toxicological concern (Varanasi *et al.* 1992). Agricultural runoff following periods of high rainfall in 1992 was implicated in a high level of bottlenose dolphin mortalities in Matagorda Bay, which is adjacent to the Western Coastal Stock area (NMFS unpublished data).

The Mississippi River, which drains about two-thirds of the continental U.S., flows into the north-central Gulf of Mexico and deposits its nutrient load which is linked to the formation of one of the world's largest areas of seasonal hypoxia (Rabalais *et al.* 1999). This area is located in Louisiana coastal waters west of the Mississippi River delta. How it affects bottlenose dolphins is not known.

#### **STATUS OF STOCK**

The bottlenose dolphin is not listed as threatened or endangered under the Endangered Species Act. However, because an UME of unprecedented size and duration (began 1 February 2010 and is ongoing) has impacted the Northern Coastal Stock area, NMFS considers this to be a strategic stock under the MMPA. Total U.S. fishery-related mortality and serious injury for this stock is not known and there is insufficient information available to determine whether the total fishery-related mortality and serious injury is insignificant and approaching the zero mortality and serious injury rate. The status of this stock relative to OSP in the Gulf of Mexico EEZ is unknown. There are insufficient data to determine the population trends for this stock.

#### **REFERENCES CITED**

- Andersen, M.S., K.A. Forney, T.V.N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley and L. Engleby 2008. Differentiating serious and non-serious injury of marine mammals: report of the serious injury technical workshop, 10-13 September 2007, Seattle, WA. NOAA Tech. Memo. NMFS-OPR-39 94 pp.
- Angliss, R.P. and D.P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: Report of the serious injury workshop, 1-2 April 1997, Silver Spring, MD. NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background and a summary of the 1995 Assessments. NOAA Tech. Memo. NMFS-OPR-6, 73 pp.
- Beier, A.G. 2001. Occurrence, distribution, and movement patterns of outer coastline bottlenose dolphins off Galveston, Texas. Master's thesis from Texas A&M University. 97 pp.
- Blaylock, R.A. and W. Hoggard. 1994. Preliminary estimates of bottlenose dolphin abundance in southern U.S. Atlantic and Gulf of Mexico continental shelf waters. NOAA Tech. Memo NMFS-SEFSC-356, 10 pp.
- Bryant, L. 1994. Report to Congress on results of feeding wild dolphins: 1989-1994. National Marine Fisheries Service, Office of Protected Resources, 23 pp.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D L. Borchers and L. Thomas. 2001. Introduction to distance sampling: Estimating abundance of biological populations. Oxford University Press, Oxford, UK. 432 pp.
- Cunningham-Smith, P., D.E. Colbert, R.S. Wells and T. Speakman. 2006. Evaluation of human interactions with a wild bottlenose dolphin (*Tursiops truncatus*) near Sarasota Bay, Florida, and efforts to curtail the interactions. *Aquat. Mamm.* 32(3):346-356.
- Fazioli, K.L., S. Hofmann and R.S. Wells. 2006. Use of Gulf of Mexico coastal waters by distinct assemblages of bottlenose dolphins (*Tursiops truncatus*). *Aquat. Mamm.* 32(2): 212-222.

- Flewelling, L.J., J.P. Naar, J.P. Abbott, D.G. Baden, N.B. Barros, G.D. Bossart, M.D. Bottein, D.G. Hammond, E.M. Haubold, C.A. Heil, M.S. Henry, H.M. Jacocks, T.A. Leighfield, R.H. Pierce, T.D. Pitchford, S.A. Rommel, P.S. Scott, K.A. Steidinger, E.W. Truby, F.M. Van Dolah and J.H. Landsberg. 2005. Red tides and marine mammal mortalities: Unexpected brevetoxin vectors may account for deaths long after or remote from an algal bloom. *Nature* 435: 755-756.
- Fulling, G.L., K.D. Mullin and C.W. Hubard. 2003. Abundance and distribution of cetaceans in outer continental shelf waters of the U.S. Gulf of Mexico. *Fish. Bull.* 101: 923-932.
- Geraci, J.R. 1990. Physiologic and toxic effects on cetaceans. Pages 167-197 in: J.R. Geraci and D.J. St. Aubin (eds.) *Sea mammals and oil: Confronting the risks*. Academic Press, New York. 259 pp.
- Gorzelany, J.F. 1998. Unusual deaths of two free-ranging Atlantic bottlenose dolphins (*Tursiops truncatus*) related to ingestion of recreational fishing gear. *Mar. Mamm. Sci.* 14(3): 614-617.
- Hansen, L.J. (ed.). 1992. Report on investigation of 1990 Gulf of Mexico bottlenose dolphin strandings. NOAA-NMFS-SEFSC Contribution MIA-92/93-21. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Irvine, A.B., M.D. Scott, R.S. Wells and J.H. Kaufmann. 1981. Movements and activities of the Atlantic bottlenose dolphin, *Tursiops truncatus*, near Sarasota, Florida. *Fish. Bull. U.S.* 79: 671-688.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. pp. 108-189 In: S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake and L. Thomas (eds.) *Advanced distance sampling*. Oxford University Press, New York. 434 pp.
- Lahvis, G.P., R.S. Wells, D.W. Kuehl, J.L. Stewart, H.L. Rhinehart and C.S. Via. 1995. Decreased lymphocyte responses in free-ranging bottlenose dolphins (*Tursiops truncatus*) are associated with increased concentrations of PCB's and DDT in peripheral blood. *Environ. Health Perspect.* 103: 67-72.
- Lehr, B., S. Bristol and A. Possolo, eds. 2010. Oil budget calculator: Deepwater Horizon. Technical documentation. Prepared by the Federal Interagency Solutions Group, Oil Budget Calculator Science and Engineering Team for the National Incident Command. Available from: [http://www.restorethegulf.gov/sites/default/files/documents/pdf/OilBudgetCalc\\_Full\\_HQ-Print\\_111110.pdf](http://www.restorethegulf.gov/sites/default/files/documents/pdf/OilBudgetCalc_Full_HQ-Print_111110.pdf)
- Lipscomb, T.P. 1993. Some answers to questions about morbillivirus. pp. 4-5. In: R.A. Blaylock, B. Mase and D.K. Odell (eds.) *Strandings*, Vol. 2, No. 3, SEFSC Miami Laboratory, Miami, Florida, 7 pp.
- Lipscomb, T.P., S. Kennedy, D. Moffet and B.K. Ford. 1994. Morbilliviral disease in an Atlantic bottlenose dolphin (*Tursiops truncatus*) from the Gulf of Mexico. *J. Wildl. Dis.* 30(4): 572-576.
- Lynn, S.K. and B. Würsig. 2002. Summer movement patterns of bottlenose dolphins in a Texas bay. *G. Mex. Sci.* 20(1): 25-37.
- Maze-Foley, K. and L.P. Garrison. in prep. Preliminary serious injury determinations for small cetaceans off the southeast U.S. coast, 2007-2011.
- McNutt, M.K., R. Camilli, T.J. Crone, G.D. Guthrie, P.A. Hsieh, T.B. Ryerson, O. Savas and F. Shaffer. 2012. Review of flow rate estimates of the *Deepwater Horizon* oil spill. *P. Natl. Acad. Sci. USA* 109 (50): 20260-20267.
- Michel, J., E.H. Owens, S. Zengel, A. Graham, Z. Nixon, T. Allard, W. Holton, P.D. Reimer, A. Lamarche, M. White, N. Rutherford, C. Childs, G. Mauseth, G. Challenger and E. Taylor. 2013. Extent and degree of shoreline oiling: *Deepwater Horizon* oil spill, Gulf of Mexico, USA. *PLOS ONE* 8(6): e65087.
- Mullin, K.D., R.R. Lohofener, W. Hoggard, C.L. Roden and C.M. Rogers. 1990. Abundance of bottlenose dolphins, *Tursiops truncatus*, in the coastal Gulf of Mexico. *Northeast Gulf Sci.* 11(2): 113-122.
- NMFS. 2004. Interim report on the bottlenose dolphin (*Tursiops truncatus*) unusual mortality event along the Panhandle of Florida, March-April 2004. 35 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- NOAA. 2011. Public scoping for preparation of a programmatic environmental impact statement for the Deepwater Horizon BP Oil Spill. Available from: <http://www.gulfspillrestoration.noaa.gov/wp-content/uploads/2011/04/Public-DWH-PEIS-Scoping-Review-Document1.pdf>
- NOAA. 2012. Federal Register 77:3233. National policy for distinguishing serious from non-serious injuries of marine mammals. Available from: <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>
- Operational Science Advisory Team (OSAT). 2010. Summary report for sub-sea and sub-surface oil and dispersant detection: Sampling and monitoring. Prepared for P.F. Zukunft, RADM, U.S. Coast Guard, Federal On-Scene Coordinator, Deepwater Horizon MC252, December 17, 2010. Available from: [http://www.restorethegulf.gov/sites/default/files/documents/pdf/OSAT\\_Report\\_FINAL\\_17DEC.pdf](http://www.restorethegulf.gov/sites/default/files/documents/pdf/OSAT_Report_FINAL_17DEC.pdf)
- Operational Science Advisory Team (OSAT-2). 2011. Summary report for fate and effects of remnant oil remaining in the beach environment. Annex B: Spatial oil distribution. Available from:

- <http://www.restorethegulf.gov/release/2011/03/01/osat-2-fate-and-effects-oil-beaches>
- Powell, J.R. and R.S. Wells. 2011. Recreational fishing depredation and associated behaviors involving common bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Mar. Mamm. Sci.* 27(1): 111-129.
- Rabalais, N.N., R.E. Turner and W.J. Wiseman, Jr. 1999. Hypoxia in the northern Gulf of Mexico: Linkage with the Mississippi River. Pages 297-322 *in*: H. Kumpf, K. Steidinger and K. Sherman (eds.) *The Gulf of Mexico large marine ecosystem*. Blackwell Science, Oxford, UK.
- Samuels, A. and L. Bejder. 2004. Chronic interactions between humans and free-ranging bottlenose dolphins near Panama City Beach, Florida, USA. *J. Cetacean Res. Manage.* 6: 69-77.
- Samuels, A. and L. Engleby. 2007. Evaluating impacts of tourism on free-ranging bottlenose dolphins near Key West, Florida. Final Technical Report, PWD Proposal # 2003-18. Available from: Dolphin Ecology Project, 727 11th Avenue South, St. Petersburg, FL 33701.
- Schwacke, L.H., E.O. Voit, L.J. Hansen, R.S. Wells, G.B. Mitchum, A.A. Hohn and P.A. Fair. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the Southeast United States coast. *Environ. Toxicol. Chem.* 21: 2752-2764.
- Scott, G.P. 1990. Management-oriented research on bottlenose dolphins by the Southeast Fisheries Center. pp. 623-639 *in*: S. Leatherwood and R.R. Reeves (eds.) *The bottlenose dolphin*. Academic Press, San Diego, CA. 653 pp.
- Sellas, A.B., R.S. Wells and P.E. Rosel. 2005. Mitochondrial and nuclear DNA analyses reveal fine scale geographic structure in bottlenose dolphins (*Tursiops truncatus*) in the Gulf of Mexico. *Conserv. Genet.* 6: 715-728.
- Varanasi, U., K.L. Tilbury, D.W. Brown, M.M. Krahn, C.A. Wigren, R.C. Clark and S.L. Chan. 1992. pp. 56-86. *In*: L.J. Hansen (ed.) *Report on investigation of 1990 Gulf of Mexico bottlenose dolphin strandings*, Southeast Fisheries Science Center Contribution MIA-92/93-21, 219 pp.
- Vollmer, N.L. 2011. Population structure of common bottlenose dolphins in coastal and offshore waters of the Gulf of Mexico revealed by genetic and environmental analyses. Ph.D. Dissertation from University of Louisiana at Lafayette. 420 pp.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, Seattle, WA. NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Wells, R.S. 1986. Structural aspects of dolphin societies. Ph.D. dissertation from University of California, Santa Cruz. 234 pp.
- Wells, R.S. and M.D. Scott. 1994. Incidence of gear entanglement for resident inshore bottlenose dolphins near Sarasota, Florida. Page 629 *in*: W.F. Perrin, G.P. Donovan and J. Barlow (eds.) *Gillnets and cetaceans*. *Rep. Int. Whal. Commn., Special Issue* 15.
- Wells, R.S. and M.D. Scott. 1997. Seasonal incidence of boat strikes on bottlenose dolphins near Sarasota, Florida. *Mar. Mamm. Sci.* 13(3): 475-480.
- Wells, R.S., S. Hofmann and T.L. Moors. 1998. Entanglement and mortality of bottlenose dolphins, *Tursiops truncatus*, in recreational fishing gear in Florida. *Fish. Bull.* 96(3): 647-650.
- Wells, R.S., V. Tornero, A. Borrell, A. Aguilar, T.K. Rowles, H.L. Rhinehart, S. Hofmann, W.M. Jarman, A.A. Hohn and J.C. Sweeney. 2005. Integrating life history and reproductive success data to examine potential relationships with organochlorine compounds for bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Sci. Total Environ.* 349: 106-119.
- Wells, R.S., J.B. Allen, S. Hoffman, K. Bassos-Hull, D.A. Fauquier, N.B. Barros, R.E. DeLynn, G. Sutton, V. Socha and M.D. Scott. 2008. Consequences of injuries on survival and reproduction of common bottlenose dolphins (*Tursiops truncatus*) along the west coast of Florida. *Mar. Mamm. Sci.* 24: 774-794.