

HARP SEAL (*Pagophilus groenlandicus*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The harp seal occurs throughout much of the North Atlantic and Arctic Oceans (Ronald and Healey 1981; Lavigne and Kovacs 1988). The world's harp seal population is divided into three separate stocks, each identified with a specific pupping site on the pack ice (Lavigne and Kovacs 1988; Bonner 1990). The largest stock is located off eastern Canada and is divided into two breeding herds. The Front herd breeds off the coast of Newfoundland and Labrador, and the Gulf herd breeds near the Magdalen Islands in the middle of the Gulf of St. Lawrence (Sergeant 1965; Lavigne and Kovacs 1988). The second stock breeds on the West Ice off eastern Greenland (Lavigne and Kovacs 1988), and the third stock breeds on the ice in the White Sea off the coast of Russia. The Front/Gulf stock is equivalent to the western North Atlantic stock. Perry *et al.* (2000) found no significant genetic differentiation between the two Northwest Atlantic whelping areas, though the authors pointed out some uncertainty surrounding that finding due to small sample sizes.

Harp seals are highly migratory (Sergeant 1965; Stenson and Sjare 1997). Breeding occurs at different times for each stock between late-February and April. Adults then assemble on suitable pack ice to undergo the annual molt. The migration then continues north to Arctic summer feeding grounds. In late September, after a summer of feeding, nearly all adults and some of the immature animals of the western North Atlantic stock migrate southward along the Labrador coast, usually reaching the entrance to the Gulf of St. Lawrence by early winter. There they split into two groups, one moving into the Gulf and the other remaining off the coast of Newfoundland. The southern limit of the harp seal's habitat extends into the U.S. Atlantic Exclusive Economic Zone (EEZ) during winter and spring.

Since the early 1990s, numbers of sightings and strandings have been increasing off the east coast of the United States from Maine to New Jersey (Katona *et al.* 1993; Rubinstein 1994; Stevick and Fernald 1998; McAlpine 1999; Lacoste and Stenson 2000; Soulen *et al.* 2013). These appearances usually occur in January-May (Harris *et al.* 2002), when the western North Atlantic stock of harp seals is at its most southern point of migration. Concomitantly, a southward shift in winter distribution off Newfoundland was observed during the mid-1990s, which was attributed to abnormal environmental conditions (Lacoste and Stenson 2000).

POPULATION SIZE

Abundance estimates for the western North Atlantic stock are available which use a variety of methods including




Figure 1: From: *Technical Briefing on the Harp Seal Hunt in Atlantic Canada*

http://www.dfo-po.gc.ca/misc/seal_briefing_e.htm

aerial surveys and mark-recapture (Table 1). These methods involve surveying the whelping concentrations and estimating total population adult numbers from pup production. Roff and Bowen (1983) developed an estimation model to provide a more precise estimate of total abundance. This technique incorporates recent pregnancy rates and estimates of age-specific hunting mortality (CAFSAC 1992). This model has subsequently been updated in Shelton *et al.* (1992, 1996), Stenson (1993), Warren *et al.* (1997), and Hammill and Stenson (2011) to consider struck and loss animals, mortality related to poor ice conditions, and variable reproductive rates. A population model was used to examine changes in the size of the population from 1952-2014 (Hammill *et al.* 2014). The model was fit to 12 estimates of pup production from 1952 to 2012, and to annual estimates of age-specific pregnancy rates between 1954 and 2013. Total population size in 2012 was estimated to be 7,445,000 (95% CI: 6.1 to 8.8 million), and projected to be 7,411,000 (95% CI: 6.1 to 8.7 million) in 2014. The population appears to be relatively stable (Hammill *et al.* 2015), though pup production has become highly variable among years (Stenson *et al.* 2014). A pup survey conducted in March 2017 will provide updated abundance estimates.

Uncertainties not accounted for include variations in reproductive rates as well as changes in mortality due to varying ice conditions.

Table 1. Summary of abundance estimates for western North Atlantic harp seals in Canadian waters. Year and area covered during each abundance survey, resulting abundance estimate (N_{best}) and confidence interval (CI).

Month/Year	Area	N_{best}	CI
2012	Front and Gulf	7.4 million	(95% CI 6.1–8.8 million)
2014 ^a	Front and Gulf	7.4 million	(95% CI 6.1–8.7 million)

^a The 2014 abundance estimate is based on model projections from the 2012 survey

Minimum population estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by (Wade and Angliss 1997). The best estimate of abundance for western North Atlantic harp seals, based on the last 2012 survey, is 7.4 million (CV=0.09, 95% CI 6.1-8.8 million; Hammill *et al.* 2014). The minimum population is 6.9 million. Data are insufficient to calculate the minimum population estimate for U.S. waters due to low sighting rates.

Current population trend

Harp seal pup production in the 1950s was estimated at 645,000, but had decreased to 225,000 by 1970 (Sergeant 1975). Estimated production then began to increase and continued to increase through the late 1990s, reaching 998,000 (CV=0.10) in 1999 (Stenson *et al.* 2003). Estimated pup production in 2008 was 1,630,300 (CV=6.8%), but decreased to 790,000 (SE=69,700, CV=8.8%) in 2012 (Stenson *et al.* 2014). This estimate is approximately half of the estimated number of pups born in 2008, likely due to lower reproductive rates in 2012 compared to 2008 (Stenson *et al.* 2014). Uncertainties in fecundity rates as well as uncertainties in ice conditions (which could impact harp seals' body condition and breeding success) have potentially large impacts on population trends.

The status of the population in U.S. waters is unknown. Recent increases in strandings may not be indicative of population size.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock due to limited understanding of stock specific life history parameters in U.S. waters. Therefore, for purposes of this assessment, the maximum net productivity rate was assumed to be 0.12. This value is based on theoretical modeling showing that pinniped populations may not grow at rates much greater than 12% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size in U.S. waters is unknown. The maximum productivity rate is 0.12, the default value for pinnipeds. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative

to optimum sustainable population (OSP) was set at 1.0 the population is increasing. PBR for the western North Atlantic harp seal in U.S. waters is unknown. The PBR for the stock in U.S. waters is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

For the period 2013–2017 the total estimated annual human caused mortality and serious injury to harp seals was 232,422. This is derived from three components: 1) 65 harp seals (CV=0.21) from the observed U.S. fisheries (Table 2a); 2) an average of 2 stranded seals from 2013-2017 that showed signs of non-fishing human interaction; and 3) an average catch of 232,355 seals from 2013-2017 by Canada and Greenland, including bycatch in the lumpfish fishery (Table 2b). Uncertainties in bycatch estimates are small compared to the magnitude of commercial and subsistence harvest in Canada. A potential source of unquantified human-caused mortality is the mortality associated with poor ice conditions due to climate change.

Fishery Information

U.S.

Detailed fishery information is reported in the Appendix III.

Northeast Sink Gillnet:

During 2013–2017, 34 mortalities were observed in the northeast sink gillnet fishery (Hatch and Orphanides 2014; 2015; 2016, Orphanides 2019, 2020). There were no observed injuries of harp seals in the Northeast region during 2013–2017 to assess using new serious injury criteria.

See Table 2a for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

Table 2a. Summary of the incidental mortality of harp seal (*Pagophilus groenlandicus*) by commercial fishery including the years sampled (Years), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

Fishery	Years	Data Type ^a	Observer Coverage ^e ^b	Observed Serious Injury ^c	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Mortality
Northeast Sink Gillnet	2013	Obs. Data, Weighout, Logbooks	0.11	0	2	0	22	22	0.75	65 (0.21)
	2014		0.18	0	9	0	57	57	0.42	
	2015		0.14	0	12	0	119	119	0.34	
	2016		0.10	0	5	0	85	85	0.50	
	2017		0.12	0	6	0	44	44	0.37	
TOTAL	-	-	-	-	-	-	-	-	65 (0.21)	

a. Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Observer Program. The Northeast Fisheries Observer Program collects landings data (Weighout) and total landings are used as a measure of total effort for the sink gillnet fishery. Mandatory logbook (Logbook) data are used to determine the spatial distribution of fishing effort in the Northeast sink gillnet fishery.

b. The observer coverages for the Northeast sink gillnet fishery and the mid-Atlantic coastal sink gillnet fisheries are ratios based on tons of fish landed. North Atlantic bottom trawl fishery coverages are ratios based on trips.

c. Serious injuries were evaluated for the 2013–2017 period and include both at-sea monitor and traditional observer data (Josephson *et al.* 2019).

Other Mortality

U.S.

From 2013 to 2017, 194 harp seal stranding mortalities were reported (Table 3; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 23 October 2018). Eleven (5.6%) of the mortalities during this five-year period showed signs of human interaction (2 in 2013, 4 in 2014, 2 in 2015, 1 in 2016 and 2 in 2017), 1 of which with some sign of fishery interaction (2013). Harris and Gupta (2006) analyzed NMFS 1996-2002 stranding data and suggested that the distribution of harp seal strandings in the Gulf of Maine was consistent with the species’ seasonal migratory patterns in this region.

CANADA

Harp seals have been commercially hunted since the mid-1800s in the Canadian Atlantic (Stenson 1993). Between 2003 and 2010 the harp seal total allowable catch (TAC) in Canada ranged from 270,000 to 330,000 (ICES 2016). In 2011 the TAC was raised to 400,000 and since then, has remained at this level each year. The TAC includes allocations for aboriginal harvesters (6,840), development of new products (20,000), and personal use (2,000). There is no specific allocation or quotas for catches in Arctic Canada. Commercial catches in Canada have remained below 80,000 since 2009 (Table 2b).

Table 2b. Summary of the Canadian directed catch and bycatch mortality of Northwest Atlantic harp seal (*Pagophilus groenlandicus*) by year.

Fishery	2013	2014	2015	2016	2017 ^f	Average
Commercial catches ^a	90,703	54,830	35,304	66,865	66,865	62,913
Struck and lost ^b	86,970	66,946	81,609	83,268	83,268	75,699
Greenland subsistence catch ^c	80,102	62,147	78,749	78,749	78,749	75,699
Canadian Arctic ^d	1,000	1,000	1,000	1,000	1,000	1,000
Newfoundland lumpfish ^e	12,330	12,330	12,330	12,330	12,330	12,330
Total	271,105	197,253	208,992	242,212	242,212	232,355

a. ICES 2016

b. Animals that are killed but not recovered and reported. Values include seals from both Canada and Greenland (ICES 2016).

c. ICES 2016. Catches in 2015 and 2016 are an average from 2005-2014

d. ICES 2016.

e. Estimates of bycatch levels in the last decade are not available and so the average annual level during the previous decade (12,330) has been assumed (DFO 2014)

f. 2017 statistics are not available. 2016 numbers are reported for 2016 and 2017.

Table 3. Harp seal (*Pagophilus groenlandicus*) stranding mortalities ^a along the U.S. Atlantic coast (2013–2017) with subtotals of animals recorded as pups in parentheses.

State	2013	2014	2015	2016	2017	Total
Maine	2	2 (1)	1	4	3	12
New Hampshire	1	0	0	2	0	3
Massachusetts	6 (1)	28	17	19 (1)	13 (1)	83
Rhode Island	1	9	4	3	4	21
Connecticut	0	0	0	1	1	2
New York	9	18	12	1	7	47
New Jersey	2	1	3	1	0	7
Delaware	1	0	0	0	0	1
Maryland	0	0	1	0	0	2
Virginia	1	9	4	1	1	6

State	2013	2014	2015	2016	2017	Total
North Carolina	2	1	2	2 (1)	2 (1)	9
Total	23	68	44	34	32	194
Unspecified seals (all states)	25	38	31	13	86	193

a. Mortalities include animals found dead and animals that were euthanized, died during handling, or died in the transfer to, or upon arrival at, rehab facilities.

STATUS OF STOCK

Harp seals are not listed as threatened or endangered under the Endangered Species Act and the western North Atlantic stock is not considered strategic under the Marine Mammal Protection Act. The level of human-caused mortality and serious injury in the U.S. Atlantic EEZ is low relative to the total stock size. The status of the harp seal stock, relative to OSP, in the U.S. Atlantic EEZ is unknown, but the stock's abundance appears to have stabilized. The total U.S. fishery-related mortality and serious injury for this stock is very low relative to the stock size and can be considered insignificant and approaching zero mortality and serious injury rate. Based on the low levels of uncertainties described in the above sections, it expected these uncertainties will have little effect on the status of this stock.

REFERENCES CITED

- Angliss, R.P. and D.P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: Report of the serious injury workshop, 1–2 April 1997, Silver Spring, MD. NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Bonner, W.N. 1990. The natural history of seals. Facts on File, New York. 196 pp.
- CAFSAC 1992. Update on population estimates of harp seal in the Northwest Atlantic. Canadian Atlantic Fisheries Scientific Advisory Committee.
- DFO 2014. Updated status of Northwest Atlantic harp seals, (*Pagophilus groenlandicus*). Department of Fisheries and Oceans. DFO Can. Sci. Advis. Sec. Sci. Rep. 2014/011. 15 pp.
- Hammill, M.O. and Stenson, G.B. 2011. Estimating abundance of Northwest Atlantic harp seals, examining the impact of density dependence. DFO Can. Sci. Advis. Sec. Res. Doc. 2011/011.
- Hammill, M.O., Stenson, G.B., Mosnier A., and Doniol-Valcroze, T. 2014. Abundance estimates of Northwest Atlantic harp seals and management advice for 2014. DFO Can. Sci. Advis. Sec. Res. Doc. 2014/022. v + 33 p.
- Hammill, M.O, Stenson, G.B., Doniol-Valcroze, T., Mosnier, A. 2015. Conservation of northwest Atlantic harp seals: Past success, future uncertainty? Biol. Cons. 192:181–191.
- Harris, D.E. and S. Gupta. 2006. GIS-based analysis of ice-breeding seal strandings in the Gulf of Maine. Northeast. Nat. 13: 403–420.
- Harris, D.E., B. Lelli and G. Jakush. 2002. Harp seal records from the southern Gulf of Maine: 1997–2001. Northeast. Nat. 9(3): 331–340.
- Hatch, J.M. and C.D. Orphanides. 2015. Estimates of cetacean and pinniped bycatch in the 2013 New England sink and mid-Atlantic gillnet fisheries. Northeast Fish. Sci. Cent. Ref. Doc.15-15. 33 pp.
- Hatch, J.M. and C.D. Orphanides. 2016. Estimates of cetacean and pinniped bycatch in the 2014 New England sink and mid-Atlantic gillnet fisheries. Northeast Fish Sci Cent Ref Doc.16-05. 22 pp.
- ICES 2016. Report of the ICES/NAFO/NAMMCO Working Group on Harp and Hooded Seals (WGHARP), 26-30 September 2016, ICES HQ, Copenhagen, Denmark. ICES CM 2016/ACOM:21.85 pp.
- Josephson, E. Wenzel F, M.C. Lyssikatos. 2019. Serious injury determinations for small cetaceans and pinnipeds caught in commercial fisheries off the northeast U.S. coast, 2013-2017. Northeast Fish. Sci. Cent. Ref. Doc. 19-17. 29 pp.
- Katona, S.K., V. Rough and D.T. Richardson. 1993. A field guide to whales, porpoises, and seals from Cape Cod to Newfoundland. Smithsonian Institution Press, Washington, DC. 316 pp.

- Lacoste, K.N. and G.B. Stenson. 2000. Winter distribution of harp seals (*Phoca groenlandica*) off eastern Newfoundland and southern Labrador. *Polar Biol.* 23: 805–811.
- Lavigne, D.M. and K.M. Kovacs. 1988. Harps and hoods: Ice breeding seals of the Northwest Atlantic. University of Waterloo Press, Waterloo, Ontario, Canada. 174 pp.
- Lyssikatos, M.C., S. Chavez-Rosales and J. Hatch. 2020. Estimates of cetacean and pinniped bycatch in Northeast and mid-Atlantic bottom trawl fisheries, 2013-2017. *Northeast Fish. Sci. Cent. Ref. Doc.* 20-04. 11 pp.
- McAlpine, D.F. 1999. Increase in extralimital occurrences of ice-breeding seals in the northern Gulf of Maine region: more seals or fewer fish. *Mar. Mamm. Sci.* 15:906911.
- Orphanides, C.D. and Hatch J. 2017. Estimates of cetacean and pinniped bycatch in the 2015 New England sink and mid-Atlantic Gillnet fisheries. *Northeast Fish Sci Cent Ref Doc.* 17-18. 21 pp.
- Orphanides, C.D. 2019. Estimates of cetacean and pinniped bycatch in the 2016 New England sink and mid-Atlantic Gillnet fisheries. *Northeast Fish Sci Cent Ref Doc.* 19-05. 12 pp.
- Orphanides, C.D. 2020. Estimates of cetacean and pinniped bycatch in the 2017 New England sink and mid-Atlantic Gillnet fisheries. *Northeast Fish Sci Cent Ref Doc.* 20-03. 16 pp. Available at: <https://www.fisheries.noaa.gov/resource/publication-database/marine-mammal-mortality-and-serious-injury-reports>
- Perry E.A., G. B. Stenson, S.E. Bartlett, W.S. Davidson and S.M. Carr. 2000. DNA sequence identifies genetically distinguishable populations of harp seals (*Pagophilus groenlandicus*) in the Northwest and Northeast Atlantic. *Mar. Biol.* 137:53–58.
- Roff, D.A. and W.D. Bowen. 1983. Population dynamics and management of the Northwest Atlantic harp seal. *Can. J. Fish. Aquat. Sci* 40: 919–932.
- Ronald, K. and P.J. Healey. 1981. Harp Seal. Pages 55–87 in: S. H. Ridgway and R. J. Harrison, (eds.) *Handbook of marine mammals, Vol. 2: Seals.* Academic Press, New York.
- Rubinstein, B. 1994. An apparent shift in distribution of ice seals, *Phoca groenlandica*, *Cystophora cristata*, and *Phoca hispida*, toward the east coast of the United States. MA thesis. Department of Biology. Boston, MA, Boston University.
- Sergeant, D.E. 1965. Migrations of harp seal *Pagophilus groenlandicus* (Erxleben) in the Northwest Atlantic. *J. Fish. Res. Board Can* 22: 433–464.
- Sergeant, D.E. 1975. Estimating numbers of harp seals. *Rapp. P.-v. Reun. Cons. int Explor. Mer.* 169:274–280.
- Shelton, P.A., N.G. Caddigan and G.B. Stenson. 1992. Model estimates of harp seal population trajectories in the Northwest Atlantic. *CAFSAC Res. Doc.* 92/89. 23 pp.
- Shelton, P.A., G.B. Stenson, B. Sjare and W.G. Warren. 1996. Model estimates of harp seal numbers-at-age for the Northwest Atlantic. *NAFO Sci. Coun. Studies* 26:1–14.
- Soulen B.K., K. Cammen, T.F. Schultz, and D.W. Johnston. 2013. Factors affecting harp seal (*Pagophilus groenlandicus*) strandings in the northwest Atlantic *PlosOne* <http://dx.doi.org/10.1371/journal.pone.0068779>
- Stenson, G.B. 1993. The status of pinnipeds in the Newfoundland region. *NAFO SCR Doc.* 93/34.
- Stenson, G.B. and B. Sjare. 1997. Seasonal distribution of harp seals, *Phoca groenlandica*, in the Northwest Atlantic. *ICES C.M.* 1997/CC:10 (Biology and Behavior II). 23 pp.
- Stenson, G.B., L.P. Rivest, M.O. Hammill, J. F. Gosselin and B. Sjare 2003. Estimating pup production of harp seals, *Pagophilus groenlandicus*, in the Northwest Atlantic. *Mar. Mamm. Sci.* 19: 141–160.
- Stenson, G. B. 2008. Recent catches of harp seals (*Pagophilus groenlandicus*) in the Northwest Atlantic DFO. *Can. Sci. Advisory Sec. Res. Doc.* 2008/080.
- Stenson, G.B., M.O. Hammill, J.W. Lawson, and J.F. Gosselin 2014. Estimating pup production of northwest Atlantic harp seals, *Pagophilus groenlandicus*, in 2012. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2014/057.v + 43 pp.
- Stevick, P.T. and T.W. Fernald. 1998. Increase in extralimital records of harp seals in Maine. *Northeast. Nat.* 5: 75–82.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Warren, W.G., P.A. Shelton and G.B. Stenson. 1997. Quantifying some of the major sources of uncertainty associated with estimates of harp seal prey consumption. Part 1: Uncertainty in the estimates of harp seal population size. *J. Northwest Atl. Fish. Sci.* 22:289–302.