

COMMON BOTTLENOSE DOLPHIN (*Tursiops truncatus truncatus*) Mississippi Sound, Lake Borgne, Bay Boudreau Stock

NOTE – NMFS is in the process of writing individual stock assessment reports for each of the 31 bay, sound and estuary stocks of common bottlenose dolphins in the Gulf of Mexico. Until this effort is completed and 31 individual reports are available, some of the basic information presented in this report will also be included in the report: “Northern Gulf of Mexico Bay, Sound and Estuary Stocks”.

STOCK DEFINITION AND GEOGRAPHIC RANGE

Common bottlenose dolphins are distributed throughout the bays, sounds and estuaries of the northern Gulf of Mexico (Mullin 1988). Long-term (year-round, multi-year) residency by at least some individuals has been reported from nearly every site where photographic identification (photo-ID) or tagging studies have been conducted in the Gulf of Mexico (e.g., Irvine and Wells 1972; Shane 1977; Gruber 1981; Irvine *et al.* 1981; Wells 1986; Wells *et al.* 1987; Scott *et al.* 1990; Shane 1990; Wells 1991; Bräger 1993; Bräger *et al.* 1994; Fertl 1994; Wells *et al.* 1996a,b; Wells *et al.* 1997; Weller 1998; Maze and Würsig 1999; Lynn and Würsig 2002; Wells 2003; Hubard *et al.* 2004; Irwin and Würsig 2004; Shane 2004; Balmer *et al.* 2008; Urian *et al.* 2009; Bassos-Hull *et al.* 2013). In many cases,

residents predominantly use the bay, sound or estuary waters, with limited movements

through passes to the Gulf of Mexico (Shane 1977; Shane 1990; Gruber 1981; Irvine *et al.* 1981; Shane 1990; Maze and Würsig 1999; Lynn and Würsig 2002; Fazioli *et al.* 2006; Bassos-Hull *et al.* 2013). Early studies indicating

year-round residency to bays in both the eastern and western Gulf of Mexico led to the delineation of

33 bay, sound and estuary stocks, including Mississippi Sound, Lake Borgne, Bay Boudreau, with the first stock assessment reports published in 1995.

More recently, genetic data also support the concept of relatively discrete, demographically independent bay, sound and estuary stocks (Duffield and Wells 2002; Sellas *et al.* 2005). Sellas *et al.* (2005) examined population subdivision among Sarasota Bay, Tampa Bay, and Charlotte Harbor, Florida; Matagorda Bay, Texas; and the coastal Gulf of Mexico (1-12 km offshore) from just outside Tampa Bay to the south end of Lemon Bay, and found evidence of significant population structure among all areas on the basis of both mitochondrial DNA control region sequence data and 9 nuclear microsatellite loci. The Sellas *et al.* (2005) findings support the identification of bay, sound and estuary populations distinct from those occurring in adjacent Gulf coastal waters. Differences in reproductive seasonality from site to site also suggest genetic-based distinctions among areas (Urian *et al.* 1996). Photo-ID and genetic data from several inshore areas of the southeastern United States also support the existence of resident estuarine animals and a differentiation between animals biopsied along the Atlantic coast and those biopsied




Figure 1. Geographic extent of the Mississippi Sound, Lake Borgne, Bay Boudreau Stock, located on the coasts of Alabama, Mississippi and Louisiana. Dark circles indicate sightings of common bottlenose dolphins during aerial surveys conducted in spring, summer and fall of 2011 and in winter of 2012.

within estuarine systems at the same latitude (Caldwell 2001; Gubbins 2002; Zolman 2002; Mazzoil *et al.* 2005; Litz 2007; Rosel *et al.* 2009; NMFS unpublished).

The Mississippi Sound, Lake Borgne, Bay Boudreau Stock area (Figure 1) is complex with an estimated surface area of 3,711 km² (Scott *et al.* 1989). Mississippi Sound itself has a surface area of about 2,100 km² (Eleuterius 1978a,b) and is bounded by Mobile Bay in the east, Lake Borgne in the west, and the opening to Bay Boudreau in the southwest. It is bordered to the north by the mainlands of Louisiana, Mississippi and Alabama and to the south by six barrier islands: Cat, West Ship, East Ship, Horn, Petit Bois and Dauphin Islands (Eleuterius 1978b), and in the extreme west, Louisiana marshes. Mississippi Sound is an open embayment with large passes between the barrier islands allowing broad access to the Gulf of Mexico, including two dredged shipping channels. Average depth at mean low water is 2.98 m, and tides are diurnal with an average range of 0.57 m (Eleuterius 1978b). Sea surface temperature ranges seasonally from 9°C to 32°C and salinity from 0 to 33 ppt from winter to summer, respectively (Christmas 1973). The bottom type is soft substrate consisting of mud and/or sand (Moncreiff 2007). Lake Borgne and Bay Boudreau are part of the Pontchartrain Basin and are remnants of the Saint Bernard lobe of Mississippi River Delta that existed until about 2000 years ago when the Mississippi River changed course (Roberts 1997; Penland *et al.* 2013). Lake Borgne has an average depth of 3 m and an average salinity of 7 ppt (USEPA 1999). Bay Boudreau is a large shallow complex in the Saint Bernard marshes and consists of marshes, bayou, shallow bays and points (Penland *et al.* 2013).

The Mississippi Sound, Lake Borgne, Bay Boudreau Stock area (“MS Sound Region”) configuration is in part a result of the management of the live-capture fishery for bottlenose dolphins (Scott 1990). Mississippi Sound was once the site of the largest live-capture fishery of bottlenose dolphins in North America (Reeves and Leatherwood 1984). Between 1973 and 1988, of the 533 bottlenose dolphins removed from Southeastern U.S. waters, 202 were removed from Mississippi Sound and adjacent waters (Scott 1990). In 1989, the Alliance of Marine Mammal Parks and Aquariums declared a self-imposed moratorium on the capture of bottlenose dolphins in the Gulf of Mexico (Corkeron 2009).

Passage of the Marine Mammal Protection Act in 1972 and the concomitant need to manage the live-capture fishery for bottlenose dolphins was the impetus for much of the earliest bottlenose dolphin research in the MS Sound Region. This work focused on estimating the abundance of bottlenose dolphins (see below) and, to a lesser extent, on stock structure research primarily to provide live-capture quota recommendations (Scott 1990). To gather baseline biological data and study dolphin ranging patterns, 57 bottlenose dolphins were captured from Mississippi Sound, freeze-branded and released from 1982-1983 (Solangi and Dukes 1983; Lohoefer *et al.* 1990a). Re-sighting efforts for these dolphins conducted from 1982-1985 by Lohoefer *et al.* (1990a) suggested at least some individual dolphins exhibited fidelity for specific areas within Mississippi Sound.

The first dedicated photo-ID effort in the area undertaken by Hubard *et al.* (2004) during 1995-1996 established a working photo-ID catalog for Mississippi Sound. Photo-ID data suggested that some individual dolphins, seen multiple times, displayed spatial and temporal patterns of site fidelity, and some dolphins showed preferences to different habitats, particularly barrier islands, channels or mainland coasts (Hubard *et al.* 2004). Some individuals were seen in the same seasons both years, while others were seen in multiple seasons with a gap during winter months (Hubard *et al.* 2004). During photo-ID/line transect surveys in 1995 and 1996, several animals photographed in 1991 (Mullin and Hoggard 1992a,b) were re-sighted (Hubard *et al.* 2004). Also, two dolphins freeze branded during the live capture performed by Solangi and Dukes (1983) were re-sighted by Hubard *et al.* (2004).

Mackey (2010) also examined site fidelity as well as residency patterns of bottlenose dolphins in a portion of Mississippi Sound using photo-ID data. During 2004-2007, Mackey (2010) primarily followed dolphins near and on both the Gulf and sound sides the barrier islands and along the Gulfport Shipping Channel and identified three different residency patterns. Of the 687 dolphins identified in those surveys, 71 (10%) were classified as year-round residents, 109 (16%) as seasonal residents, and 498 (73.5%) as transients. These patterns may not be representative of the MS Sound Region. Dolphins sighted near the barrier islands adjacent to or within the range of the Northern Coastal Stock of bottlenose dolphins may have a higher probability of being transient. Outside of the ship channel, a small proportion of the dolphins sighted by Mackey (2010) were from the interior two-thirds of Mississippi Sound (adjacent to the mainland) where dolphins may have quite different residency patterns. Mackey (2010) also identified two animals that were freeze-branded during the live captures 20 years earlier (Solangi and Dukes 1983). Both Mackey (2010) and Hubard *et al.* (2004) noted low re-sighting rates of dolphins with a high percentage of dolphins seen only on one occasion. Both studies also suggested dolphins move out of the Sound into deeper Gulf of Mexico waters during winter months (Hubard *et al.* 2004; Mackey 2010). Definitive conclusions on bottlenose dolphin site fidelity and residency patterns in the MS Sound Region are difficult to make based on available research. Establishing residency patterns in the MS Sound Region using photo-ID studies that cover large study areas (e.g., Hubard *et al.* 2004) will be difficult because of the large number of dolphins that inhabit the area and its

open geography. Nevertheless, studies to date indicate that, similar to other Gulf of Mexico areas, some individuals are long-term inhabitants of the MS Sound Region. The current stock boundary does not include any coastal waters outside of the barrier islands. Further research is needed to determine the degree to which dolphins of this stock utilize nearshore coastal waters outside the MS Sound Region. The stock boundaries are subject to change upon further study of dolphin residency patterns in estuarine waters of Alabama, Mississippi and Louisiana. Information on the use of coastal waters will be important when considering exposure to coastal fisheries as estuarine animals that make use of nearshore coastal waters would be at risk of entanglement in fishing gear while moving along the coast. Ongoing NOAA photo-ID surveys initiated in 2010, as well as data from tracking of 19 bottlenose dolphins tagged with satellite-linked transmitters in and around Mississippi Sound in July 2013, will address some of these issues as the data become available.

POPULATION SIZE

The best available abundance estimate for the Mississippi Sound, Lake Borgne, Bay Boudreau Stock of bottlenose dolphins is 901 (CV=0.63) based on a winter 2012 aerial survey.

Recent surveys and abundance estimates

The Southeast Fisheries Science Center conducted aerial surveys of continental shelf waters (shoreline to 200 m depth) along the U.S. Gulf of Mexico coast from the Florida Keys to the Texas/Mexico border during spring (March-April) 2011, summer (July-August) 2011, fall (October-November) 2011 and winter (January-February) 2012. The surveys were conducted along tracklines oriented perpendicular to the shoreline and spaced 20-30 km apart. The total survey effort varied during each survey due to weather conditions, but ranged between 13,500 – 15,600 km. Each of these surveys was conducted using a two-team approach to develop estimates of visibility bias using the independent observer approach with Distance analysis (Laake and Borchers 2004). A model for the probability of detection on the trackline as a function of sighting conditions (seas state, glare, water color, etc.) was developed using data across all four surveys. This model was then applied to detection probability functions specific to each survey to account for the probability of detection as a function of distance from the trackline and additional environmental covariates. A bootstrap resampling approach was used to estimate the variance of the estimates. The survey data were post-stratified into spatial boundaries corresponding to the defined boundaries of bottlenose dolphin stocks within the surveyed area. The abundance estimates for the Mississippi Sound, Lake Borgne, Bay Boudreau Stock of bottlenose dolphins were based upon tracklines and sightings in waters along the Alabama, Mississippi and Louisiana coasts inside of the barrier islands. The surveys did not include tracklines in Lake Borgne, but the estimated density was extrapolated to include the entire stock area. The seasonal abundance estimates for this stock were: spring – 2,395 (CV=0.42), summer – 1,709 (CV= 0.59), fall – 1,140 (CV=0.41) and winter – 900 (CV=0.63). As with other bay, sound and estuary stocks, it is possible that there is movement of transient animals from coastal waters into the MS Sound region on a seasonal basis. In order to assure that the abundance estimate for the stock reflects primarily resident animals, the lowest seasonal estimate (winter) was used to determine N_{best} for this stock. The resulting best estimate of abundance for the Mississippi Sound, Lake Borgne, Bay Boudreau Stock of bottlenose dolphins was 900 (CV=0.63).

Earlier abundance estimates

Aerial and small boat surveys conducted in the MS Sound Region covered different portions of the region and yielded a wide range of abundance estimates for bottlenose dolphins. Because of the differences in techniques and areas surveyed, it is very difficult to compare results. Aerial strip transect surveys conducted by Leatherwood *et al.* (1978) compared aerial survey techniques for bottlenose dolphins, but the study also produced population estimates for bottlenose dolphins in Mississippi Sound and the adjacent Gulf of Mexico (to about 10 km south of the barrier islands) of $1,342 \pm 847$ in 1974 and 879 ± 368 in 1975. Thompson (1982) surveyed central Mississippi Sound (“off Pascagoula”) in 1980 using aerial line-transect sampling methods, and abundance estimates ranged from 93 dolphins (SE=22) in December 1980 to 140 dolphins (SE=86) in September 1980. While line-transect is a rigorous and repeatable survey method, this study produced negatively biased estimates of density and abundance (Thompson 1982) due to the fact that the strip of transect directly under the aircraft was not observed. Scott *et al.* (1989) attempted to correct this bias by utilizing an aircraft with a glass bubble nose and placing an observer in it to observe the track-line at all times. Their estimates for the MS Sound Region ranged from 205 in winter to 858 in summer. (Abundances for Mississippi Sound only ranged from 136 dolphins in winter to 719 dolphins in summer.) Boat-based mark-recapture surveys using dolphins freeze-branded during a previous live-capture study were performed by Lohofener *et al.* (1990a) to assess the impacts of removing 30 dolphins from the population for captivity. The pre-removal estimate was 2,392 dolphins, and the post-removal estimate was 7,052 dolphins (Lohofener *et al.*

1990a), but these were probably not accurate estimates, as too many assumptions of mark-recapture analysis were likely violated in this study (Lohoefer *et al.* 1990a). Boat-based line-transect abundance surveys of Mississippi Sound (about 55% of the MS Sound Region) were carried out by Lohoefer *et al.* (1990b) in 1984 and 1985, yielding much higher abundance estimates than aerial strip- or line-transect surveys and suggesting a seasonal shift in bottlenose dolphin abundance. For the entire Sound, abundance estimates were 2,400 and 500 dolphins for summer and winter, respectively. Another series of line-transect aerial surveys were performed in fall of 1992 by Blaylock and Hoggard (1994), where the abundance was reported as 1,401 for the MS Sound Region. The two most recent abundance estimates from Mississippi Sound were boat-based line-transect surveys and only covered a portion of Mississippi Sound. Hubard *et al.* (2004) surveyed an area bounded by the western end of Horn Island and the eastern end of Petit Bois Island that was roughly one-quarter the size of the entire Sound. Again, abundances were found to fluctuate seasonally with higher abundances observed in summer months in 1995 (584 dolphins) and 1996 (555 dolphins) versus winter 1995-1996 months (268 dolphins). Miller *et al.* (2013) reported abundance estimates for a study area in eastern Mississippi Sound roughly 2,104 km² in size that included areas up to 15 km south of the barrier islands. Abundance estimates were 2,255 dolphins in summer 2007 and 1,413 dolphins in winter 2007-2008 (Miller *et al.* 2013).

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distributed abundance estimate as specified by Wade and Angliss (1997). The best estimate of abundance for this stock of bottlenose dolphins is 901 (CV=0.63). The minimum population estimate for the MS Sound Region is 551 bottlenose dolphins.

Current Population Trend

There are insufficient data to determine the population trends for this stock.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. The maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of the minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size of bottlenose dolphins in the MS Sound Region is 551. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.5 because this stock is of unknown status. PBR for the Mississippi Sound, Lake Borgne, Bay Boudreau Stock of bottlenose dolphins is 5.6.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual human-caused mortality and serious injury of bottlenose dolphins in the MS Sound Region during 2008-2012 is unknown. During 2008-2012, 2 mortalities were documented in the Gulf of Mexico menhaden purse seine fishery, 3 mortalities were documented involving the Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line) fishery, and 1 mortality was documented in the Gulf of Mexico blue crab trap/pot fishery. In addition, 1 mortality was documented in a research gillnet, and 1 mortality occurred incidental to sea turtle relocation trawling. It is not possible to estimate the total number of mortalities or serious injuries associated with menhaden purse seine, hook and line, or blue crab trap/pot fisheries since there are no systematic observer programs for those fisheries.

New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998; Andersen *et al.* 2008; NOAA 2012). NMFS defines serious injury as an “injury that is more likely than not to result in mortality”. Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

Fishery Information

The commercial fisheries that potentially could interact with this stock are the Category II Southeastern U.S. Atlantic, Gulf of Mexico shrimp trawl, Gulf of Mexico menhaden purse seine and the Category III Gulf of Mexico blue crab trap/pot and Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line) fisheries (Appendix III).

Menhaden Purse Seine Fishery

During 2008-2012, there were 2 mortalities and 1 animal released alive without serious injury documented within waters of the MS Sound Region involving the menhaden purse seine fishery.

There is currently no observer program for the Gulf of Mexico menhaden purse seine fishery; however, recent incidental takes have been reported via two sources. First, during 2011, a pilot observer program operated from May through September, and observers documented 3 dolphins trapped within purse seine nets. All 3 were released alive without serious injury (Maze-Foley and Garrison in prep). Two of the 3 dolphins were trapped within a single purse seine within waters of the Western Coastal Stock, and the third animal was trapped in waters of the MS Sound Region. Second, through the Marine Mammal Authorization Program (MMAP), there have been 13 self-reported incidental takes (all mortalities) of bottlenose dolphins in northern Gulf of Mexico coastal and estuarine waters by the menhaden purse seine fishery. These takes likely affected the following stocks: Western Coastal Stock; Northern Coastal Stock; Mississippi Sound, Lake Borgne, Bay Boudreau Stock; and Mississippi River Delta Stock. Specific self-reported takes under the MMAP likely involving the MS Sound Region are as follows: two dolphins were reported taken in a single purse seine during 2012 in waters of Mississippi Sound; one take of a single unidentified dolphin was reported during 2002 in waters of Mississippi Sound; and during 2000, 3 bottlenose dolphins were reported taken in a single purse seine in waters of Mississippi Sound.

Without an ongoing observer program it is not possible to obtain statistically reliable information for this fishery on the number of sets annually, the incidental take and mortality rates, and the communities from which bottlenose dolphins are being taken.

Hook and Line Fisheries

During 2008-2012 there were 3 mortalities for which hook and line gear entanglement or ingestion were documented. Two mortalities occurred during 2011 and 1 during 2012. These mortalities were included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 [for 2008-2011 data] and 15 April 2013 [for 2012 data]) and are included in the totals presented in Table 1.

Blue Crab Trap/Pot Fishery

During 2008-2012 there was 1 documented mortality of a bottlenose dolphin in commercial blue crab trap/pot gear. The mortality occurred during 2011 and was included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 and 15 April 2013) and in the totals presented in Table 1. There is no systematic observer coverage of crab trap/pot fisheries, so it is not possible to quantify total mortality.

Shrimp Trawl Fishery

During 2008-2012 there were no documented interactions with the commercial shrimp trawl fishery within the MS Sound Region; however, it should be noted that observer coverage of the shrimp trawl fishery does not extend into bay, sound and estuary waters.

Other Mortality

From 2008 to 2012, 306 bottlenose dolphins were reported stranded within the MS Sound Region (Table 1; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, 13 September 2012 and 15 April 2013). Evidence of human interactions was detected for 18 stranded dolphins. Human interactions were from numerous sources, including 3 entanglements with hook and line gear, 1 entanglement with commercial blue crab trap/pot gear, 1 incidental take in a research gillnet, 1 incidental take during turtle relocation trawling, 2 gunshot wounds, and 2 animals that were visibly oiled (see Table 1). Stranding data probably underestimate the extent of human-caused mortality and serious injury because not all of the marine mammals that die or are seriously injured in human interactions are discovered, reported or investigated, nor will all of those that are found necessarily show signs of entanglement or other human interaction. Finally, the level of technical expertise among stranding

network personnel varies widely as does the ability to recognize signs of human interactions.

The MS Sound Region has been affected by several bottlenose dolphin die-offs or Unusual Mortality Events (UMEs). From January through May 1990, a total of 367 bottlenose dolphins stranded in the northern Gulf of Mexico including Mississippi. Overall this represented a two-fold increase in the prior maximum recorded number of strandings for the same period, but in some locations (i.e., Alabama) strandings were 10 times the average number. The cause of the 1990 mortality event could not be determined (Hansen 1992). In 1996 a UME was declared for bottlenose dolphins in Mississippi when 27 bottlenose dolphins stranded during November and December. The cause was not determined, but a *Karenia brevis* (red tide) bloom was suspected to be responsible. A UME was declared for cetaceans in the northern Gulf of Mexico beginning 1 February 2010; and, as of 2013, the event is still ongoing. It includes cetaceans that stranded prior to the Deepwater Horizon oil spill (see “Habitat Issues” below), during the spill, and after. During 2010, 92 stranded dolphins from this stock were considered to be part of the UME; during 2011, 115 dolphins, and during 2012, 45 dolphins.

One mortality was documented in 2011 in the MS Sound Region as a result of an entanglement in a research gillnet. This mortality was included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 and 15 April 2013) and in the totals presented in Table 1.

As part of its annual coastal dredging program, the Army Corps of Engineers conducts sea turtle relocation trawling during hopper dredging as a protective measure for marine turtles. One bottlenose dolphin mortality was documented during 2011 in MS Sound Region incidental to relocation trawling activities. This mortality was included in the stranding database (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 13 September 2012 and 15 April 2013) and in the totals presented in Table 1.

The problem of dolphin depredation of fishing gear is increasing in Gulf of Mexico coastal and estuary waters and illegal feeding or provisioning of wild bottlenose dolphins has been documented in Florida and Texas (Bryant 1994; Samuels and Bejder 2004; Cunningham-Smith *et al.* 2006; Powell and Wells 2011). There are emerging questions regarding potential linkages between provisioning and depredation of recreational fishing gear and associated entanglement and ingestion of gear. To date there are no records of provisioning for this stock area. However, one recent case of a shrimp fisherman illegally “taking” a dolphin in Mississippi Sound occurred during summer 2012. In December 2013 the fisherman was convicted under the MMPA for knowingly shooting a dolphin with a shotgun while shrimping.

Table 1. Common bottlenose dolphin strandings occurring in the Mississippi Sound, Lake Borgne, Bay Boudreau Stock area from 2008 to 2012, as well as number of strandings for which evidence of human interaction was detected and number of strandings for which it could not be determined (CBD) if there was evidence of human interaction (HI). Data are from the NOAA National Marine Mammal Health and Stranding Response Database (unpublished data, accessed 13 September 2012 [for 2008-2011 data] and 15 April 2013 [for 2012 data]). Please note human interaction does not necessarily mean the interaction caused the animal’s death.

Stock	Category	2008	2009	2010	2011	2012	Total
Mississippi Sound, Lake Borgne, Bay Boudreau Stock	Total Stranded	16	36	93 ^a	116 ^b	45 ^c	306
	Human Interaction						
	---Yes	0	1	7 ^d	7 ^e	3 ^f	18
	---No	2	3	2	5	1	13
	---CBD	14	32	84	104	41	275

^a 92 strandings were part of the ongoing UME event in the northern Gulf of Mexico.

^b 115 strandings were part of the ongoing UME event in the northern Gulf of Mexico.

^c All 45 strandings were part of the ongoing UME event in the northern Gulf of Mexico.

^d Includes 2 strandings that were visibly oiled.

^e Includes 2 entanglement interactions (mortalities) with hook and line fishing gear, 1 entanglement interaction (mortality) with commercial blue crab trap/pot gear, 1 mortality incidental to sea turtle relocation trawling, and 1 entanglement interaction (mortality) with a research gillnet.

^f Includes 1 entanglement interaction (mortality) with hook and line fishing gear and 2 stranded animals (mortalities) with gunshot wounds.

HABITAT ISSUES

The Deepwater Horizon (DWH) MC252 drilling platform, located approximately 50 miles southeast of the Mississippi River Delta in waters about 1500m deep, exploded on 20 April 2010. The rig sank, and over 87 days ~4.9 million barrels of oil were discharged from the wellhead until it was capped on 15 July 2010 (McNutt *et al.* 2012). During the response effort dispersants were applied extensively at the seafloor and at the sea surface (Lehr *et al.* 2010; OSAT 2010). In-situ burning, or controlled burning of oil at the surface, was also used extensively as a response tool (Lehr *et al.* 2010). The oil, dispersant and burn residue compounds present ecological concerns. The magnitude of this oil spill was unprecedented in U.S. history, causing impacts to wildlife, natural habitats and human communities along coastal areas from western Louisiana to the Florida Panhandle (NOAA 2011). It could be years before the entire scope of damage is ascertained (NOAA 2011).

Given the trajectory of the surface oil during the spill and the documented oiling of shoreline (Michel *et al.* 2013), it is likely the Mississippi Sound, Lake Borgne, Bay Boudreau Stock of bottlenose dolphins was exposed to oil during the event. Light to trace oil was reported along the majority of Mississippi's mainland coast, from Gulf Breeze to Panama City, Florida, and outside of Atchafalaya and Vermilion Bays in western Louisiana. Heavy to light oiling occurred on Mississippi's barrier islands (Michel *et al.* 2013). A substantial number of beaches and wetlands along the Louisiana coast experienced heavy or moderate oiling (OSAT-2 2011; Michel *et al.* 2013). The heaviest oiling in Louisiana occurred west of the Mississippi River on the Mississippi Delta and in Barataria and Terrebonne Bays, and to the east of the river on the Chandeleur Islands. Some heavy to moderate oiling occurred on Alabama and Florida beaches, with the heaviest stretch occurring from Dauphin Island, Alabama, to Gulf Breeze, Florida.

Shortly after the oil spill, the Natural Resource Damage Assessment (NRDA) process was initiated under the Oil Pollution Act of 1990. A variety of NRDA research studies are being conducted to determine potential impacts of the spill on marine mammals. These studies have focused on identifying the type, magnitude, severity, length and impact of oil exposure to oceanic, continental shelf, coastal and estuarine marine mammals. The research is ongoing. For coastal and estuarine dolphins, the NOAA-led efforts include: active surveillance to detect stranded animals in remote locations; aerial surveys to document the distribution, abundance, species and exposure of marine mammals and sea turtles relative to oil from DWH spill; assessment of sublethal and chronic health impacts on coastal and estuarine bottlenose dolphins in Barataria Bay, Louisiana, and a reference site in Sarasota Bay, Florida; and assessment of injuries to dolphin stocks in Barataria Bay and Chandeleur Sound, Louisiana, Mississippi Sound, and as a reference site, St. Joseph Bay, Florida.

Coastal dolphins have been observed with tar balls attached to them and seen swimming through oil slicks close to shore and inland bays. The effects of oil exposure on marine mammals depend on a number of factors including the type and mixture of chemicals involved, the amount, frequency and duration of exposure, the route of exposure (inhaled, ingested, absorbed, or external) and biomedical risk factors of the particular animal (Geraci 1990). In general, direct external contact with petroleum compounds or dispersants with skin may cause skin irritation, chemical burns and infections. Inhalation of volatile petroleum compounds or dispersants may irritate or injure the respiratory tract, which could lead to pneumonia or inflammation. Ingestion of petroleum compounds may cause injury to the gastrointestinal tract, which could affect an animal's ability to digest or absorb food. Absorption of petroleum compounds or dispersants may damage kidney, liver and brain function in addition to causing immune suppression and anemia. Long term chronic effects such as lowered reproductive success and decreased survival may occur (Geraci 1990).

Besides oil exposure, another habitat concern for the MS Sound Region is environmental contaminants. Persistent organic pollutant (PCBs, chlordanes, mirex, DDTs, HCB and dieldrin) and polybrominated diphenyl ether concentrations were determined from bottlenose dolphin blubber samples from 14 locations, including Mississippi Sound, along the U.S. Atlantic and Gulf coasts and Bermuda (Kucklick *et al.* 2011). Dolphins from both rural and urban estuarine and coastal waters were sampled. Dolphins sampled from Mississippi Sound had relatively high concentrations of some pollutants, like PBDEs, HCB, mirex and DDTs, and more intermediate concentrations of dieldrin, PCBs and chlordanes, when compared to dolphins sampled from the other 13 locations (Kucklick *et al.* 2011).

The presence of vessels may impact bottlenose dolphin behavior in bays, sounds and estuaries. Miller *et al.* (2008) investigated the immediate responses of bottlenose dolphins to "high-speed personal watercraft" (i.e., boats) in Mississippi Sound. They found an immediate impact on dolphin behavior demonstrated by an increase in traveling behavior and dive duration, and a decrease in feeding behavior for non-traveling groups. The findings suggested dolphins attempted to avoid high-speed personal watercraft. It is unclear whether repeated short-term effects will result in long-term consequences like reduced health and viability of dolphins. Further studies are needed to determine the impacts throughout the Gulf of Mexico.

STATUS OF STOCK

Bottlenose dolphins are not listed as threatened or endangered under the Endangered Species Act. However, because an UME of unprecedented size and duration (began 1 February 2010 and is ongoing) has impacted the northern Gulf of Mexico, including the Mississippi Sound, Lake Borgne, Bay Boudreau Stock, NMFS considers this stock to be strategic under the MMPA. The total human-caused mortality and serious injury for this stock is unknown and there is insufficient information available to determine whether the total fishery-related mortality and serious injury for this stock is insignificant and approaching zero mortality and serious injury rate. The status of this stock relative to OSP is unknown. There are insufficient data to determine population trends for this stock.

REFERENCES CITED

- Andersen, M.S., K.A. Forney, T.V.N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley and L. Engleby. 2008. Differentiating serious and non-serious injury of marine mammals: report of the serious injury technical workshop, 10-13 September 2007, Seattle, WA. NOAA Tech. Memo. NMFS-OPR-39. 94 pp.
- Angliss, R.P. and D.P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: Report of the serious injury workshop, 1-2 April 1997, Silver Spring, MD. NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Balmer, B.C., R.S. Wells, S.M. Nowacek, D.P. Nowacek, L.H. Schwacke, W.A. McLellan, F.S. Scharf, T.K. Rowles, L.J. Hansen, T.R. Spradlin and D.A. Pabst. 2008. Seasonal abundance and distribution patterns of common bottlenose dolphins (*Tursiops truncatus*) near St. Joseph Bay, Florida, USA. *J. Cetacean Res. Manage.* 10(2): 157-167.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Bassos-Hull, K., R.M. Perrtree, C.C. Shepard, S. Schilling, A.A. Barleycorn, J.B. Allen, B.C. Balmer, W.E. Pine and R.S. Wells. 2013. Long-term site fidelity and seasonal abundance estimates of common bottlenose dolphins (*Tursiops truncatus*) along the southwest coast of Florida and responses to natural perturbations. *J. Cetacean Res. Manage.* 13(1):19-30.
- Blaylock, R.A. and W. Hoggard. 1994. Preliminary estimates of bottlenose dolphin abundance in southern U.S. Atlantic and Gulf of Mexico continental shelf waters. NOAA Tech. Memo. NMFS-SEFSC-356, 10 pp.
- Bräger, S. 1993. Diurnal and seasonal behavior patterns of bottlenose dolphins (*Tursiops truncatus*). *Mar. Mamm. Sci.* 9: 434-440.
- Bräger, S., B. Würsig, A. Acevedo and T. Henningsen. 1994. Association patterns of bottlenose dolphins (*Tursiops truncatus*) in Galveston Bay, Texas. *J. Mamm.* 75(2): 431-437.
- Bryant, L. 1994. Report to Congress on results of feeding wild dolphins: 1989-1994. National Marine Fisheries Service, Office of Protected Resources, 23 pp.
- Corkeron, P. 2009. Captivity. Pp 183-188 in W.F. Perrin, B. Würsig, J.G. M. Thewissen (eds), *The encyclopedia of marine mammals*. Academic Press.
- Caldwell, M. 2001. Social and genetic structure of bottlenose dolphin (*Tursiops truncatus*) in Jacksonville, Florida. Ph.D. dissertation from University of Miami. 143 pp.
- Christmas, J.Y. 1973. Phase I: Area description. Pages 1-71 in: J.Y. Christmas, (ed.) *Cooperative Gulf of Mexico estuarine inventory and study*, Mississippi. Gulf Coast Research Laboratory, Ocean Springs, MS.
- Cunningham-Smith, P., D.E. Colbert, R.S. Wells and T. Speakman. 2006. Evaluation of human interactions with a wild bottlenose dolphin (*Tursiops truncatus*) near Sarasota Bay, Florida, and efforts to curtail the interactions. *Aquat. Mamm.* 32(3): 346-356.
- Duffield, D.A. and R.S. Wells. 2002. The molecular profile of a resident community of bottlenose dolphins, *Tursiops truncatus*. Pages 3-11 in: C. J. Pfeiffer, (ed.) *Cell and Molecular Biology of Marine Mammals*. Krieger Publishing, Melbourne, FL.
- Eleuterius, C.K. 1978a. Geographical definition of Mississippi Sound. *Gulf Research Reports* 6:179-181.
- Eleuterius, C.K. 1978b. Classification of Mississippi Sound as to estuary hydrological type. *Gulf Research Reports* 6:185-187.
- Fazioli, K.L., S. Hofmann and R.S. Wells 2006. Use of Gulf of Mexico coastal waters by distinct assemblages of bottlenose dolphins (*Tursiops truncatus*). *Aquat. Mamm.* 32(2): 212-222.
- Fertl, D.C. 1994. Occurrence patterns and behavior of bottlenose dolphins (*Tursiops truncatus*) in the Galveston ship channel. *Texas J. Sci.* 46: 299-317.

- Geraci, J.R. 1990. Physiologic and toxic effects on cetaceans. Pages 167-197 in: J. R. Geraci and D. J. St. Aubin (eds.) Sea mammals and oil: Confronting the risks. Academic Press, New York. 259 pp.
- Gruber, J.A. 1981. Ecology of the Atlantic bottlenosed dolphin (*Tursiops truncatus*) in the Pass Cavallo area of Matagorda Bay, Texas. M. Sc. thesis from Texas A&M University, College Station. 182 pp.
- Gubbins, C. 2002. Association patterns of resident bottlenose dolphins (*Tursiops truncatus*) in a South Carolina estuary. *Aquat. Mamm.* 28: 24-31.
- Hansen, L.J. (ed.) 1992. Report on investigation of 1990 Gulf of Mexico bottlenose dolphin strandings. NOAA-NMFS-SEFSC Contribution MIA-92/93-21. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Hubard, C.W., K. Maze-Foley, K.D. Mullin and W.W. Schroeder 2004. Seasonal abundance and site fidelity of bottlenose dolphins (*Tursiops truncatus*) in Mississippi Sound. *Aquat. Mamm.* 30: 299-310.
- Irvine, A.B., M.D. Scott, R.S. Wells and J.H. Kaufmann. 1981. Movements and activities of the Atlantic bottlenose dolphin, *Tursiops truncatus*, near Sarasota, Florida. *Fish. Bull.* 79: 671-688.
- Irvine, B. and R.S. Wells. 1972. Results of attempts to tag Atlantic bottlenose dolphins (*Tursiops truncatus*). *Cetology* 13: 1-5.
- Irwin, L.J. and B. Würsig. 2004. A small resident community of bottlenose dolphins, *Tursiops truncatus*, in Texas: Monitoring recommendations. *G. Mex. Sci.* 22(1): 13-21.
- Kucklick, J., L. Schwacke, R. Wells, A. Hohn, A. Guichard, J. Yordy, L. Hansen, E. Zolman, R. Wilson, J. Litz, D. Nowacek, T. Rowles, R. Pugh, B. Balmer, C. Sinclair and P. Rosel. 2011. Bottlenose dolphins as indicators of persistent organic pollutants in the western North Atlantic Ocean and northern Gulf of Mexico. *Environ. Sci. Technol.* 45: 4270-4277.
- Laake, J.L. and D.L. Borchers 2004. Methods for incomplete detection at distance zero. Pages 108-189 in: S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake and L. Thomas (eds.) *Advanced distance sampling*. Oxford University Press, New York. 434 pp.
- Leatherwood, S., J.R. Gilbert and D.G. Chapman. 1978. An evaluation of some techniques for aerial censuses of bottlenosed dolphins. *Journal of Wildlife Management* 42(2):239-250.
- Lehr, B., S. Bristol and A. Possolo, eds. 2010. Oil budget calculator: Deepwater Horizon. Technical documentation. Prepared by the Federal Interagency Solutions Group, Oil Budget Calculator Science and Engineering Team for the National Incident Command. Available from: http://www.restorethegulf.gov/sites/default/files/documents/pdf/OilBudgetCalc_Full_HQ-Print_111110.pdf
- Litz, J.A. 2007. Social structure, genetic structure, and persistent organohalogen pollutants in bottlenose dolphins (*Tursiops truncatus*) in Biscayne Bay, Florida. Ph.D. dissertation from University of Miami. 140 pp.
- Lohofener, R., W. Hoggard, R. Ford and J. Benigno. 1990a. Studies of Mississippi Sound bottlenose dolphins: Assessing the effects of the removal of 30 bottlenose dolphins from Mississippi Sound. Part 1(of 2) of the Final report to the U.S. Marine Mammal Commission in partial fulfillment of interagency agreement MM2910909-2. Available from: National Marine Fisheries Service, P.O. Drawer 1207, Pascagoula, MS 39568.
- Lohofener, R., W. Hoggard, K. Mullin, R. Ford and J. Benigno. 1990b. Studies of Mississippi Sound bottlenose dolphins: Estimates of bottlenose dolphin density in Mississippi Sound from small boat surveys. Part 2 (of 2) of the Final report the U.S. Marine Mammal Commission in partial fulfillment of interagency agreement MM2910909-2. Available from: National Marine Fisheries Service, P.O. Drawer 1207, Pascagoula, MS 39568.
- Lynn, S.K. and B. Würsig 2002. Summer movement patterns of bottlenose dolphins in a Texas bay. *G. Mex. Sci.* 20(1): 25-37.
- Mackey, A.D. 2010. Site fidelity and association patterns of bottlenose dolphins (*Tursiops truncatus*) in the Mississippi Sound. M.A. thesis. The University of Southern Mississippi, Hattiesburg. 144 pp.
- Maze, K.S. and B. Würsig 1999. Bottlenose dolphins of San Luis Pass, Texas: Occurrence patterns, site fidelity, and habitat use. *Aquat. Mamm.* 25: 91-103.
- Maze-Foley, K. and L.P. Garrison. in prep. Preliminary serious injury determinations for small cetaceans off the southeast U.S. coast, 2007-2011.
- Mazzoil, M., S.D. McCulloch and R.H. Defran. 2005. Observations on the site fidelity of bottlenose dolphins (*Tursiops truncatus*) in the Indian River Lagoon, Florida. *Fla. Sci.* 68: 217-226.
- McNutt, M.K., R. Camilli, T.J. Crone, G.D. Guthrie, P.A. Hsieh, T.B. Ryerson, O. Savas and F. Shaffer. 2012. Review of flow rate estimates of the *Deepwater Horizon* oil spill. *P. Natl. Acad. Sci. USA* 109 (50): 20260-20267.
- Michel, J., E.H. Owens, S. Zengel, A. Graham, Z. Nixon, T. Allard, W. Holton, P.D. Reimer, A. Lamarche, M.

- White, N. Rutherford, C. Childs, G. Mauseth, G. Challenger and E. Taylor. 2013. Extent and degree of shoreline oiling: *Deepwater Horizon* oil spill, Gulf of Mexico, USA. PLOS ONE 8(6): e65087.
- Miller, L.J., A.D. Mackey, M. Solangi and S.A. Kuczaj II. 2013. Population abundance and habitat utilization of bottlenose dolphins in the Mississippi Sound. *Aquatic Conserv: Mar. Freshw. Ecosyst.* 23: 145-151.
- Miller, L.J., M. Solangi and S.A. Kuczaj, II. 2008. Immediate response of Atlantic bottlenose dolphins to high-speed personal watercraft in the Mississippi Sound. *J. Mar. Biol. Assoc. U.K.* 88(6): 1139-1143.
- Moncreiff, C.A. 2007. Mississippi Sound and the Gulf Islands. Pages 76-85 *in*: L. Handley, D. Altsman and R. DeMay, (eds.) *Seagrass status and trends in the northern Gulf of Mexico: 1940-2002*. USGS Scientific Investigations Report 2006-5287.
- Mullin, K.D. 1988. Comparative seasonal abundance and ecology of bottlenose dolphins (*Tursiops truncatus*) in three habitats of the north-central Gulf of Mexico. Ph.D. thesis. Mississippi State University, Starkville. 135 pp.
- Mullin, K.D. and W. Hoggard. 1992a. Low-level monitoring of the abundance of bottlenose dolphins in Mississippi Sound: Progress Report #1 (1991 surveys). Available from: National Marine Fisheries Service, P.O. Drawer 1207, Pascagoula, MS 39568.
- Mullin, K.D. and W. Hoggard. 1992b. Low-level monitoring of the abundance of bottlenose dolphins in Mississippi Sound: Progress Report #2 (1992 surveys). Available from: National Marine Fisheries Service, P.O. Drawer 1207, Pascagoula, MS 39568.
- NOAA. 2011. Public scoping for preparation of a programmatic environmental impact statement for the Deepwater Horizon BP Oil Spill. Available from: <http://www.gulfspillrestoration.noaa.gov/wp-content/uploads/2011/04/Public-DWH-PEIS-Scoping-Review-Document1.pdf>
- NOAA. 2012. Federal Register 77:3233. National policy for distinguishing serious from non-serious injuries of marine mammals. Available from: <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>
- Operational Science Advisory Team (OSAT). 2010. Summary report for sub-sea and sub-surface oil and dispersant detection: Sampling and monitoring. Prepared for P. F. Zukunft, RADM, U.S. Coast Guard, Federal On-Scene Coordinator, Deepwater Horizon MC252, December 17, 2010. Available from: http://www.restorethegulf.gov/sites/default/files/documents/pdf/OSAT_Report_FINAL_17DEC.pdf
- Operational Science Advisory Team (OSAT-2). 2011. Summary report for fate and effects of remnant oil remaining in the beach environment. Annex B: Spatial oil distribution. Available from: <http://www.restorethegulf.gov/release/2011/03/01/osat-2-fate-and-effects-oil-beaches>
- Penland, S., A. Beall and J. Kindinger (eds). 2013. Environmental atlas of the Lake Pontchartrain Basin. USGS Open File Report 02-206. <http://pubs.usgs.gov/of/2002/of02-206/>.
- Powell, J.R. and R.S. Wells. 2011. Recreational fishing depredation and associated behaviors involving common bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Mar. Mamm. Sci.* 27(1): 111-129.
- Reeves, R.R. and S. Leatherwood. 1984. Live-capture fisheries for cetaceans in USA and Canadian waters, 1973-1982. *Rep. Int. Whal. Comm.* 34: 497-507.
- Roberts, H.H. 1997. Dynamic changes of the Holocene Mississippi River delta plain: The delta cycle. *J. Coastal Res.* 13:605-627.
- Rosel, P.E., L. Hansen and A.A. Hohn. 2009. Restricted dispersal in a continuously distributed marine species: common bottlenose dolphins *Tursiops truncatus* in coastal waters of the western North Atlantic. *Molec. Ecol.* 18: 5030-5045.
- Samuels, A. and L. Bejder. 2004. Chronic interactions between humans and free-ranging bottlenose dolphins near Panama City Beach, Florida, USA. *J. Cetacean Res. Manage.* 6: 69-77.
- Scott, G.P. 1990. Management-oriented research on bottlenose dolphins in the Southeast Fisheries Center. Pages 623-639 *in*: S. Leatherwood and R.R. Reeves, (eds.) *The bottlenose dolphin*. Academic Press, San Diego, CA.
- Scott, G.P., D.M. Burn, L.J. Hansen and R.E. Owen. 1989. Estimates of bottlenose dolphin abundance in the Gulf of Mexico from regional aerial surveys. CRD 88/89-07. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Scott, M.D., R.S. Wells and A.B. Irvine 1990. A long-term study of bottlenose dolphins on the west coast of Florida. Pages 235-244 *in*: S. Leatherwood and R.R. Reeves, (eds.) *The bottlenose dolphin*. Academic Press, San Diego, CA.
- Sellas, A.B., R.S. Wells and P.E. Rosel 2005. Mitochondrial and nuclear DNA analyses reveal fine scale geographic structure in bottlenose dolphins (*Tursiops truncatus*) in the Gulf of Mexico. *Conserv. Genet.* 6(5): 715-728.
- Shane, S.H. 1977. The population biology of the Atlantic bottlenose dolphin, *Tursiops truncatus*, in the Aransas Pass area of Texas. M. Sc. thesis from Texas A&M University, College Station. 238 pp.

- Shane, S.H. 1990. Behavior and ecology of the bottlenose dolphin at Sanibel Island, Florida. Pages 245-265 in: S. Leatherwood and R.R. Reeves, (eds.) The bottlenose dolphin. Academic Press, San Diego, CA.
- Shane, S.H. 2004. Residence patterns, group characteristics, and association patterns of bottlenose dolphins near Sanibel Island, Florida. *G. Mex. Sci.* 22(1): 1-12.
- Solangi, M.S. and G.E. Dukes. 1983. Atlantic bottlenose dolphin, *Tursiops truncatus*, herd studies in the Mississippi Sound, USA: Capture, freeze marking and biological sampling. Final Report, NMFS (Contract No. NA82-GA-C-00023). Available from: National Marine Fisheries Service, P.O. Drawer 1207, Pascagoula, MS 39568.
- Thompson, N.B. 1982. Estimates of abundance of *Tursiops truncatus*, the bottlenose dolphin in: St. Joseph-Apalachicola Bays, Florida; Mississippi Sound, Mississippi; and the Aransas-Copano-San Antonio Bay complex, Texas. NOAA/NMFS/SEFC/Mami Laboratory Fishery Analysis Division Technical Report.
- Urian, K.W., D.A. Duffield, A.J. Read, R.S. Wells and D.D. Shell. 1996. Seasonality of reproduction in bottlenose dolphins, *Tursiops truncatus*. *J. Mamm.* 77: 394-403.
- Urian, K.W., S. Hofmann, R.S. Wells and A.J. Read. 2009. Fine-scale population structure of bottlenose dolphins (*Tursiops truncatus*) in Tampa Bay, Florida. *Mar. Mamm. Sci.* 25(9): 619-638.
- USEPA. 1999. Ecological Condition of Estuaries in the Gulf of Mexico. EPA 620-R-98-004. U.S. Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Gulf Ecology Division, Gulf Breeze, Florida.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Weller, D.W. 1998. Global and regional variation in the biology and behavior of bottlenose dolphins. Ph. D. thesis from Texas A&M University, College Station. 142 pp.
- Wells, R.S. 1986. Population structure of bottlenose dolphins: Behavioral studies along the central west coast of Florida. Contract report to NMFS, SEFSC. Contract No. 45-WCNF-5-00366. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149. 58 pp.
- Wells, R.S. 1991. The role of long-term study in understanding the social structure of a bottlenose dolphin community. Pages 199-225 in: K. Pryor and K.S. Norris, (eds.) Dolphin societies: Discoveries and puzzles. University of California Press, Berkeley.
- Wells, R.S. 2003. Dolphin social complexity: Lessons from long-term study and life history. Pages 32-56 in: F.B.M. de Waal and P.L. Tyack, (eds.) Animal social complexity: Intelligence, culture, and individualized societies. Harvard University Press, Cambridge, MA.
- Wells, R.S., M.K. Bassos, K.W. Urian, W.J. Carr and M.D. Scott. 1996a. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Charlotte Harbor, Florida: 1990-1994. NOAA Tech. Memo. NMFS-SEFSC-384. 36 pp.
- Wells, R.S., M.K. Bassos, K.W. Urian, S.H. Shane, E.C.G. Owen, C.F. Weiss, W.J. Carr and M.D. Scott. 1997. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Pine Island Sound, Florida: 1996. Contract report to National Marine Fisheries Service, Southeast Fisheries Center Contribution No. 40-WCNF601958. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Wells, R.S., M.D. Scott and A.B. Irvine. 1987. The social structure of free ranging bottlenose dolphins. Pages 247-305 in: H. Genoways, (ed.) Current Mammalogy, Vol. 1. Plenum Press, New York.
- Wells, R.S., K.W. Urian, A.J. Read, M.K. Bassos, W.J. Carr and M.D. Scott. 1996b. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Tampa Bay, Florida: 1988-1993. NOAA Tech. Memo. NMFS-SEFSC-385. 25 pp.
- Zolman, E.S. 2002. Residence patterns of bottlenose dolphins (*Tursiops truncatus*) in the Stono River estuary, Charleston County, South Carolina, U.S.A. *Mar. Mamm. Sci.* 18: 879-892.