

HARBOR SEAL (*Phoca vitulina richardsi*): Gulf of Alaska Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Harbor seals inhabit coastal and estuarine waters off Baja California, north along the western coasts of the United States, British Columbia, and Southeast Alaska, west through the Gulf of Alaska and Aleutian Islands, and in the Bering Sea northward to Cape Newenham and the Pribilof Islands. They haul out on rocks, reefs, beaches, and drifting glacial ice, and feed in marine, estuarine, and occasionally fresh waters. Harbor seals generally are non-migratory, with local movements associated with such factors as tides, weather, season, food availability, and reproduction (Scheffer and Slipp 1944; Fisher 1952; Bigg 1969, 1981). The results of recent satellite tagging studies in Southeast Alaska, Prince William Sound, and Kodiak are also consistent with the conclusion that harbor seals are non-migratory (Frost et al. 1996, Swain et al. 1996). However, some long-distance movements of tagged animals in Alaska have been recorded (Pitcher and McAllister 1981, Frost et al. 1996). Strong fidelity of individuals for haulout sites in June and August also has been reported, although these studies considered only limited areas during a relatively short period of time (Pitcher and Calkins 1979, Pitcher and McAllister 1981).

The following information was considered in classifying stock structure based on the Dizon et al. (1992) phylogeographic approach: 1) Distributional data: geographic distribution continuous, natal dispersal characteristics unknown, breeding dispersal is presumed to be very limited, year-round site fidelity observed, seasonal movements greater than 300 km rare (Harvey 1987) except in western Alaska (Hoover-Miller 1994); 2) Population response data: substantial differences in population dynamics between Southeast Alaska and the rest of Alaska, and presumed differences between Gulf of Alaska and Bering Sea (Hoover 1988, Hoover-Miller 1994, Withrow and Loughlin 1996); 3) Phenotypic data: clinal variation in body size and color phase (Shaughnessy and Fay 1977, Kelly 1981); 4) Genotypic data: undetermined for Alaska, mitochondrial DNA analyses currently underway. Preliminary genetic data indicate substantial variation in mtDNA suggesting at least two genetically distinct stocks in Alaska (Westlake and O’Corry-Crowe 1997). However, until additional samples are analyzed the Alaska Scientific Review Group (SRG) recommended using the same stock boundaries as in the Stock Assessment Reports for 1996 (Hill et al. 1997).

The Alaska SRG concluded that the scientific data available to support three distinct biological stocks (i.e., genetically isolated populations) were equivocal. However, the Alaska SRG recommended that the available data were sufficient to justify the establishment of three management units for harbor seals in Alaska (DeMaster 1996). Further, the SRG recommended that, unlike the stock structure reported in Small and DeMaster (1995), animals in the Aleutian Islands should be included in the same management unit as animals in the Gulf of Alaska. As noted above, this recommendation has been adopted by NMFS with the caveat that management units and stocks are equivalent for the purposes of managing incidental take under section 118 of the Marine Mammal Protection Act (Wade and Angliss 1997). Therefore, based primarily on the significant population decline of seals in the Gulf of Alaska, the possible decline in the Bering Sea, and the stable population in Southeast Alaska (see Current Population Trend section in the respective harbor seal report for details), three separate stocks are recognized in Alaska waters: 1) the Southeast Alaska stock - occurring from the Alaska/British Columbia border to Cape Suckling, Alaska (144°W), 2) the Gulf of Alaska




Figure 8. Approximate distribution of harbor seals in Alaska waters (shaded area).

stock - occurring from Cape Suckling to Unimak Pass, including animals throughout the Aleutian Islands, and 3) the Bering Sea stock - including all waters north of Unimak Pass (Fig. 8). Information concerning the three harbor seal stocks recognized along the West Coast of the continental United States can be found in the Stock Assessment Reports for the Pacific Region.

POPULATION SIZE

Extensive photographic aerial surveys of harbor seals from the Gulf of Alaska stock were conducted during 1994 and 1996. The Aleutian Islands were surveyed from 29 August to 8 September of 1994 (Withrow and Loughlin 1995a). Between 25 August and 3 September of 1996 the south side of the Alaska Peninsula, Cook Inlet, Kenai Peninsula, Kodiak Archipelago, and Copper River Delta were surveyed (Withrow and Loughlin 1997). All known harbor seal haulout sites in each area were surveyed, and reconnaissance surveys were flown prior to photographic surveys to establish the location of additional sites. Aerial surveys were flown within 2 hours on either side of low tide, based on the assumption that at locations affected by tides, harbor seals haul out in greatest numbers at and around the time of low tide (Pitcher and Calkins 1979, Calambokidis et al. 1987). One to seven repetitive photographic counts were obtained for each major haulout site within each study area. Coefficients of variation (CV) were determined for multiple surveys and found to be <0.19 in all cases. This method of estimating abundance and its CV assumes that during the survey period no migration occurred between sites and that there was no trend in the number of animals ashore. The number of seals moving between areas was assumed to be small considering each area's large geographic size, though a small number of seals may have been counted twice or not at all.

During summer of 1996, two different aerial surveys covered portions of Prince William Sound. During August 17-26 surveys of trend route A in Prince William Sound resulted in an adjusted mean count of 984 (CV=0.045) seals (Frost et al. 1997). Between August 27 and September 6 surveys of trend route B, excluding Columbia Bay (a tidewater glacial haulout system), in Prince William Sound resulted in a mean count of 1,261 (CV=0.044) seals (unpubl. data, J. Burns, Living Resources Inc., P. O. Box 83570, Fairbanks, AK, 99708). During the route B surveys, the count data from Columbia Bay were considered unreliable due to difficult ice conditions and the widely scattered distribution of seals. Instead, a reasonable minimum estimate for the number of harbor seals using Columbia Bay at the time of the surveys (1,000 seals) will be added below (see Minimum Population Estimate section). Combining the counts from trend routes A and B results in a mean count of 2,245 (CV=0.032) harbor seals in Prince William Sound, excluding Columbia Bay.

Due to the extreme difficulty in censusing harbor seals during the 1994 Aleutian Islands survey, it is recommended that the maximum count of 3,437 be used for an abundance estimate for that region (Withrow and Loughlin 1995a). The coefficient of variation for the mean count (CV=0.059) should be used for the 1994 survey data because an estimate for the CV is not available for the maximum count. The mean count for the 1996 surveys was 16,013 (CV=0.025) harbor seals, with the following mean counts for the major survey areas: Copper River Delta 3,174 (CV=0.078); Prince William Sound 2,245; Kenai Peninsula 713 (CV=0.072); Cook Inlet 2,244 (CV=0.105); Kodiak Archipelago 4,437 (CV=0.035); and the south side of the Alaska Peninsula 3,200 (CV=0.034). Therefore, for the Gulf of Alaska stock of harbor seals, the total combined count from the 1994 and 1996 aerial surveys was 19,450 (CV=0.023) animals.

Data collected from 36 tagged harbor seals in Southeast Alaska during 1994 resulted in a correction factor of 1.74 (CV=0.068) to account for animals in the water which are thus missed during the aerial surveys (Withrow and Loughlin 1995b). In 1995, 25 harbor seals were tagged at a sand bar haulout near Cordova, AK (note: within the Gulf of Alaska). The haulout behavior of these seals was monitored from August 12 to 23, and a correction factor of 1.50 (CV=0.047) was developed for the 1995 aerial survey in this area (Withrow and Loughlin 1996). Although much of the haulout substrate in the Gulf of Alaska area is rocky, the 1.50 CF (correction factor) from 1995 is considered to be the best available and most conservative CF for the 1996 survey data because the data used to estimate the CF were 1) collected in the survey area, 2) collected during a comparable low-tide survey window, and 3) collected more closely to the peak haul out time period (i.e., CF data collected from 12 August to 23 August versus the survey data from 23 August to 9 September). The Southeast Alaska correction factor of 1.74 was not employed for this stock because the data used to calculate the CF were 1) not collected from the Gulf of Alaska area and 2) collected to some extent after the survey period was completed (i.e., CF data from SE Alaska were collected from 1 September to 11 September)(Alaska SRG, see DeMaster 1996). Therefore, using the Gulf of Alaska correction factor results in an abundance estimate of 29,175 ($19,450 \times 1.50$, CV=0.052) for the Gulf of Alaska stock of harbor seals.

The next round of aerial surveys to assess the abundance of this stock will occur during the summers of 1999 (Aleutian Islands) and 2001 (Gulf of Alaska). Preliminary results of these surveys will be available in autumn of the respective survey year.

Minimum Population Estimate

The minimum population estimate (N_{MIN}) for this stock is calculated using Equation 1 from the PBR Guidelines (Wade and Angliss 1997): $N_{\text{MIN}} = N/\exp(0.842 \times [\ln(1 + [\text{CV}(N)]^2)]^{1/2})$. Using the population estimate (N) of 29,175 and its associated $\text{CV}(N)$ of 0.052, N_{MIN} for this stock of harbor seals is 27,917. Including the minimum population estimate for Columbia Bay (1,000 animals) results in an N_{MIN} of 28,917 harbor seals for the Gulf of Alaska stock.

Current Population Trend

The population trend in the Aleutian Islands is unclear because the 1994 survey was the most complete census to date for that region. Previous harbor seal counts in that area are not comparable to the 1994 data because they were conducted incidental to surveys designed to assess other species (i.e., sea otters or Steller sea lions). However, a subset of the 1994 survey in the eastern Aleutian Islands indicated a count of 1,600 in an area that had counts of approximately 1,000-2,500 seals during 1975-77 (Small 1996).

In Prince William Sound, harbor seal numbers declined by 57% from 1984 to 1992 (Pitcher 1989, Frost and Lowry 1993). The decline began before the 1989 *Exxon Valdez* oil spill, was greatest in the year of the spill, and may have lessened thereafter. Between 1989 and 1995 aerial survey counts of 25 haulout sites in Prince William Sound (trend route A) showed significant declines in the number of seals during the molt (19%) and during pupping (31%) (Frost et al. 1996). Adjusted molt period counts for 1996 were 15% lower than the 1995 counts, indicating that harbor seal numbers in Prince William Sound have not yet recovered from the spill or whatever was causing the decline and that the long-term decline has not ended (Frost et al. 1997).

A steady decrease in numbers of harbor seals has been reported throughout the Kodiak Archipelago from the mid-1970s to the 1990s. On southwestern Tugidak Island, formally one of the largest concentrations of harbor seals in the world, counts declined 85% from 1976 (6,919) to 1988 (1,014) (Pitcher 1990). More recently, the Tugidak Island count has increased from 769 in 1992 to 1,420 in 1996 (Small 1996, Withrow and Loughlin 1997), although this still only represents a fraction of its historical size. The population around Kodiak Island, based on an aerial photographic route established in 1992, is estimated to have increased at 7.2% annually from 1992-96 (Small et al. 1997). Despite some positive signs of growth in certain areas, the overall Gulf of Alaska stock size remains small compared to its size in the 1970s and 1980s.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Reliable rates of maximum net productivity have not been estimated for the Gulf of Alaska or Bering Sea harbor seal stock. Population growth rates were estimated at 6% and 8% between 1991 and 1992 in Oregon and Washington, respectively (Huber et al. 1994). Harbor seals have been protected in British Columbia since 1970, and the population has responded with an annual rate of increase of approximately 12.5% since 1973 (Olesiuk et al. 1990). However, until additional data become available from which more reliable estimates of population growth can be determined, it is recommended that the pinniped maximum theoretical net productivity rate (R_{MAX}) of 12% be employed for this stock (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 re-authorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $\text{PBR} = N_{\text{MIN}} \times 0.5R_{\text{MAX}} \times F_{\text{R}}$. The recovery factor (F_{R}) for this stock is 0.5, the value for pinniped stocks with unknown status (Wade and Angliss 1997). Thus, for the Gulf of Alaska stock of harbor seals, $\text{PBR} = 868$ animals ($28,917 \times 0.06 \times 0.5$).

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fisheries Information

Three different commercial fisheries operating within the range of the Gulf of Alaska stock of harbor seals were monitored for incidental take by fishery observers during 1990-96: Gulf of Alaska groundfish trawl, longline, and pot fisheries. For the fisheries with observed takes, the range of observer coverage over the 7-year period, as well as the annual observed and estimated mortalities are presented in Table 7a. The mean annual (total) mortality rate was 0.4 (CV=1.0) for the Gulf of Alaska groundfish trawl fishery and was 0.2 (CV=1.0) Gulf of Alaska pot fishery. The harbor seal taken in the pot fishery in 1995 (7% observer coverage) occurred during an unmonitored haul and therefore could not be used to estimate mortality for the entire fishery. Therefore, 1 mortality was used as both the observed mortality and estimated mortality in 1995 for that fishery, and should be considered a minimum estimate.

Table 7a. Summary of incidental mortality of harbor seals (Gulf of Alaska stock) due to commercial fisheries from 1990 through 1996 and calculation of the mean annual mortality rate. Mean annual mortality in brackets represents a minimum estimate from self-reported fisheries information or stranding data. Data from 1992 to 1996 (or the most recent 5 years of available data) are used in the mortality calculation when more than 5 years of data are provided for a particular fishery. n/a indicates that data are not available.

Fishery name	Years	Data type	Range of observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Gulf of Alaska (GOA) groundfish trawl	90-96	obs data	33-55%	0, 1, 1, 0, 0, 0, 0	0, 3, 2, 0, 0, 0, 0	0.4 (CV=1.0)
GOA finfish pot	90-96	obs data	5-13%	0, 0, 0, 0, 0, 1, 0	0, 0, 0, 0, 0, 1, 0	0.2 (CV=1.0)
Prince William Sound salmon drift gillnet	90-91	obs data	4-5%	2, 1	36, 12	24 (CV=0.50)
Alaska Peninsula/Aleutian Islands salmon drift gillnet	90	obs data	4%	0	0	0
Observer program total						24.6 (CV=0.49)
				Reported mortalities		
Cook Inlet salmon set gillnet	90-96	self reports	n/a	6, 0, 1, 0, n/a, n/a, n/a	n/a	[≥ 1.75]
Prince William Sound set gillnet	90-96	self reports	n/a	0, 0, 0, 1, n/a, n/a, n/a	n/a	[≥ 0.25]
Kodiak salmon set gillnet	90-96	self reports	n/a	3, 0, 0, 0, n/a, n/a, n/a	n/a	[≥ 0.75]
Alaska salmon purse seine (except for Southeast)	90-96	self reports	n/a	0, 0, 0, 2, n/a, n/a, n/a	n/a	[≥ 0.5]
Alaska Peninsula/Aleutian Islands salmon drift gillnet	90-96	self reports	n/a	9, 2, 12, 5, n/a, n/a, n/a	n/a	[≥ 7.0]
unknown Gulf of Alaska fishery	92-96	strand data	n/a	0, 0, 0, 0, 1	n/a	[≥ 0.2]
Minimum total annual mortality						≥ 35.05 (CV=0.49)

In the Prince William Sound salmon drift gillnet fishery, observers recorded 2 incidental mortalities of harbor seals in 1990 (Wynne et al. 1991), and 1 in 1991 (Wynne et al. 1992). The extrapolated kill estimates were 36 (95% CI 2-74) in 1990 and 12 (95% CI 1-44) in 1991, resulting in a mean kill rate of 24 (CV=0.5) animals per year for this fishery. In 1990, observers boarded 300 (57.3%) of the 524 vessels that fished in the Prince William Sound salmon drift gillnet fishery, monitoring a total of 3,166 sets, or roughly 4% of the estimated number of sets made by the fleet. In 1991, observers boarded 531 (86.9%) of the 611 registered vessels and monitored a total of 5,875 sets, or roughly 5% of the estimated sets made by the fleet. The estimated mortality rate of harbor seals based on the 1990 and 1991 observed mortalities for this fishery is 0.0002 kills per set. Fisher self-reports of harbor seal mortalities due to this fishery detail 19, 4, 7, 24, and 0 mortalities in 1990, 1991, 1992, 1993, and 1996, respectively. The extrapolated (estimated) mortality from the 1990-91 observer program (24 seals per year) accounts for these mortalities, so they do not appear in Table 7a. Combining the estimates from the groundfish trawl and pot fisheries presented above ($0.4 + 0.2 = 0.6$) with the estimate from the Prince William Sound salmon drift gillnet fishery (24) results in an estimated annual incidental kill rate in observed fisheries of 24.6 (CV=0.49) harbor seals per year from this stock. It should be noted that in 1990, observers also boarded 59 (38.3%) of the 154 vessels participating in the Alaska Peninsula/Aleutian Island salmon drift gillnet fishery, monitoring a total of 373 sets, or roughly 4% of the estimated number of sets made by the fleet (Wynne et al. 1991). Although no interaction with harbor seals was recorded by observers in 1990, due in part to the low level of observer coverage, mortalities did occur as recorded in fisher self-reports (see Table 7a).

An additional source of information on the number of harbor seals killed or injured incidental to commercial fishery operations is the self-reported fisheries information required of vessel operators by the MMPA. During the period between 1990 and 1996, fisher self-reports from 5 unobserved fisheries (see Table 7a) resulted in an annual mean of 10.25 mortalities from interactions with commercial fishing gear. However, because logbook records (fisher self-reports required during 1990-94) are most likely negatively biased (Credle et al. 1994), these are considered to be minimum estimates. These totals are based on all available self-reported fisheries information for Gulf of Alaska fisheries, except the Prince William Sound salmon drift gillnet fishery and the Gulf of Alaska groundfish trawl and pot fisheries for which observer data were presented above. In 1990, fisher self-reports from the Cook Inlet set and drift gillnet fisheries were combined. As a result, some of the harbor seal mortalities reported in 1990 may have occurred in the drift net fishery. Self-reported fisheries data are incomplete for 1994, not available for 1995, and considered unreliable for 1996 (see Appendix 4).

Strandings of harbor seals entangled in fishing gear or with injuries caused by interactions with gear are another source of mortality data. During the 5-year period from 1992 to 1996 the only fishery-related harbor seal stranding was reported in June of 1996 on Middleton Island. The entanglement could not be attributed to a particular fishery and as a result has been included in Table 7a as occurring in an unknown fishery. Fishery-related strandings during 1992-96 result in an estimated annual mortality of 0.2 harbor seals from this stock. This estimate is considered a minimum because not all entangled animals strand and not all stranded animals are found or reported.

The estimated minimum annual mortality rate incidental to commercial fisheries is 36 (rounded up), based on observer data (24.6) and self-reported fisheries information (10.25) or stranding data (0.2) where observer data were not available. However, a reliable estimate of the mortality rate incidental to commercial fisheries is currently unavailable because of the absence of observer placements in several fisheries mentioned above.

Subsistence/Native Harvest Information

The 1992-96 subsistence harvest of harbor seals in Alaska was estimated by the Alaska Department of Fish and Game, under contract with the NMFS (Table 7b: Wolfe and Mishler 1993, 1994, 1995, 1996, 1997). In each year, data were collected through systematic interviews with hunters and users of marine mammals in approximately 2,100 households in about 60 coastal communities within the geographic range of the harbor seal in Alaska. Between 1992-96, interviews were conducted in approximately 29 communities that lie within the range of the Gulf of Alaska harbor seal stock. The statewide total subsistence take of harbor seals in 1992 was estimated at 2,888 (95% CI 2,320-3,741), with 2,535 harvested and 353 struck and lost. The total subsistence take in 1993 was estimated at 2,736 (95% CI 2,334-3,471), with 2,365 harvested and 371 struck and lost. The total subsistence take in 1994 was estimated at 2,621 (95% CI 2,110-3,457), with 2,313 harvested and 308 struck and lost. The total subsistence take in 1995 was estimated at 2,742 (95% CI 2,184-3,679), with 2,499 harvested and 243 struck and lost. The total subsistence take in 1996 was estimated at 2,741 (95% CI 2,378-3,479), with 2,415 harvested and 327 struck and lost.

Table 7b provides a summary of the subsistence harvest information for the Gulf of Alaska stock. The mean annual subsistence take from this stock of harbor seals, including struck and lost, over the 3-year period from 1994 to 1996 was 791 animals. The reported average age-specific kill of the harvest from the Gulf of Alaska stock since 1992 was 58% adults, 27% juveniles, 2% pups, and 13% of unknown age. The reported average sex-specific kill of the harvest was 44% males, 18% females, and 38% of unknown sex.

Table 7b. Summary of the subsistence harvest data for the Gulf of Alaska stock of harbor seals, 1992-96.

Year	Estimated total number taken	Percentage of statewide total	Number harvested	Number struck and lost
1992	967	33.7%	884	83
1993	914	33.5%	812	102
1994	913	34.9%	819	94
1995	724	26.4%	683	41
1996	735	26.8%	679	56
Mean annual take (1994-96)	791			

Other Mortality

Illegal intentional killing of harbor seals occurs, but the magnitude of this mortality is unknown (Note: the 1994 Amendments to the MMPA made intentional lethal take of any marine mammal illegal except where imminently necessary to protect human life).

STATUS OF STOCK

Sustainable harvest levels for this stock will be determined from the analysis of information gathered through the cooperative management process, and will reflect the degree of uncertainty associated with the information obtained for this stock. Efforts were initiated in 1995 and 1996 to develop a cooperative approach for management of this stock; however, a final agreement has not been approved to date.

Harbor seals are not listed as “depleted” under the MMPA or listed as “threatened” or “endangered” under the Endangered Species Act. A reliable estimate of the annual rate of mortality incidental to commercial fisheries is unavailable. Therefore, it is unknown whether the kill rate due to commercial fishing is insignificant. At present, annual fishery-related mortality levels less than 87 animals per year (i.e., 10% of PBR) can be considered insignificant and approaching zero mortality and serious injury rate. Based on currently available data, the estimated annual level of total human-caused mortality is 827 (36 + 791) harbor seals which does not exceed the PBR (868) for this stock. Until additional information on mortality incidental to commercial fisheries becomes available, the Gulf of Alaska stock of harbor seals is not classified as strategic. This classification is consistent with the recommendations of the Alaska SRG (DeMaster 1998). The status of this stock relative to its Optimum Sustainable Population size is unknown.

REFERENCES

- Bigg, M. A. 1969. The harbour seal in British Columbia. Fish. Res. Bd. Can. Bull. 172. 33 pp.
- Bigg, M. A. 1981. Harbour seal, *Phoca vitulina*, Linnaeus, 1758 and *Phoca largha*, Pallas, 1811. Pp. 1-27, In S. H. Ridgway and R. J. Harrison (eds.), Handbook of Marine Mammals, vol.2: Seals. Academic Press, New York.
- Calambokidis, J., B. L. Taylor, S. D. Carter, G. H. Steiger, P. K. Dawson, and L. D. Antrim. 1987. Distribution and haul out behavior of harbor seals in Glacier Bay, Alaska. Can. J. Zool. 65:1391-1396.
- Credle, V. R., D. P. DeMaster, M. M. Merklein, M. B. Hanson, W. A. Karp, and S. M. Fitzgerald (eds.). 1994. NMFS observer programs: minutes and recommendations from a workshop held in Galveston, Texas, November 10-11, 1993. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-94-1, 96 pp.

- DeMaster, D. P. 1996. Minutes from the 11-13 September 1996 meeting of the Alaska Scientific Review Group, Anchorage, Alaska. 20 pp. + appendices. (available upon request - D. P. DeMaster, National Marine Mammal Laboratory, 7600 Sand Point Way, NE, Seattle, WA 98115).
- DeMaster, D. P. 1998. Minutes from sixth meeting of the Alaska Scientific Review Group, 21-23 October 1997, Seattle, Washington. 40 pp. (available upon request - D. P. DeMaster, National Marine Mammal Laboratory, 7600 Sand Point Way, NE, Seattle, WA 98115).
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Fisher, H. D. 1952. The status of the harbour seal in British Columbia, with particular reference to the Skeena River. *Fish. Res. Bd. Can. Bull.* 93. 58 pp.
- Frost, K. F., and L. F. Lowry. 1993. Assessment of injury to harbor seals in Prince William Sound, Alaska, and adjacent areas following the *Exxon Valdez* oil spill. State-Federal Natural Resource Damage Assessment, Marine Mammals Study No.5. 95 pp.
- Frost, K. F., L. F. Lowry, R. J. Small, and S. J. Iverson. 1996. Monitoring, habitat use, and trophic interactions of harbor seals in Prince William Sound. *Exxon Valdez* Oil Spill Restoration Project Annual Report (Project # 95064), Alaska Dep. of Fish and Game, Division of Wildlife Conservation. Fairbanks, AK. 131 pp.
- Frost, K. F., L. F. Lowry, J. M. Ver Hoef, and S. J. Iverson. 1997. Monitoring, habitat use, and trophic interactions of harbor seals in Prince William Sound, Alaska. *Exxon Valdez* Oil Spill Restoration Project Annual Report (Project # 96064), Alaska Dep. of Fish and Game, Division of Wildlife Conservation. Fairbanks, AK. 56 pp.
- Harvey, J. T. 1987. Population dynamics, annual food consumption, movements, and diving behavior of harbor seals, *Phoca vitulina*, in Oregon. Ph.D. dissertation, Oregon State Univ., Corvallis, OR. 177 pp.
- Hill, P. S., D. P. DeMaster, and R. J. Small. 1997. Alaska Marine Mammal Stock Assessments, 1996. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-78, 150 pp.
- Hoover, A. A. 1988. Harbor seal. Pp. 125-157, *In* J. W. Lentfer (ed.), Selected marine mammals of Alaska: Species accounts with research and management recommendations. Marine Mammal Commission, Washington, D.C.
- Hoover-Miller, A. A. 1994. Harbor seal (*Phoca vitulina*) biology and management in Alaska. Marine Mammal Commission, Washington D.C., Contract #T75134749.
- Huber, H., S. Jeffries, R. Brown, and R. DeLong. 1994. Harbor Seal Stock Assessment in Washington and Oregon 1993. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Kelly, B. P. 1981. Pelage polymorphism in Pacific harbor seals. *Can. J. Zool.* 59:1212-1219.
- Olesiuk, P. F., M. A. Bigg, and G. M. Ellis. 1990. Recent trends in the abundance of harbour seals, *Phoca vitulina*, in British Columbia. *Can. J. Fish. and Aquat. Sci.* 47:992-1003.
- Pitcher, K. W. 1989. Harbor seal trend count surveys in southern Alaska, 1988. Final report Contract MM4465852-1 to U.S. Marine Mammal Commission, Washington, D.C. 15 pp.
- Pitcher, K. W. 1990. Major decline in number of harbor seals, *Phoca vitulina richardsi*, on Tugidak Island, Gulf of Alaska. *Mar. Mammal Sci.* 6:121-134.
- Pitcher, K. W., and D. G. Calkins. 1979. Biology of the harbor seal (*Phoca vitulina richardsi*) in the Gulf of Alaska. U.S. Dep. Commer., NOAA, OCSEAP Final Rep. 19(1983):231-310.
- Pitcher, K. W., and D. C. McAllister. 1981. Movements and haul out behavior of radio-tagged harbor seals, *Phoca vitulina*. *Can. Field Nat.* 95:292-297.
- Scheffer, V. B., and J. W. Slipp. 1944. The harbor seal in Washington state. *Amer. Midl. Nat.* 32:373-416.
- Shaughnessy, P. D., and F. H. Fay. 1977. A review of the taxonomy and nomenclature of North Pacific harbour seals. *J. Zool. (Lond.)*. 182:385-419.
- Small, R. J. 1996. Population assessment of harbor seals in Alaska: report of a workshop held in Fairbanks, Alaska, November 14-16, 1995. 36 pp.
- Small, R. J., and D. P. DeMaster. 1995. Alaska marine mammal stock assessments 1995. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-57, 93 pp.
- Small, R. J., G. W. Pendelton, and K. M. Wynne. 1997. Harbor seal population trends in the Ketchikan, Sitka, and Kodiak Island areas of Alaska. Pp. 7-32 *In* Annual Report: Harbor seal investigations in Alaska. NOAA Grant NA57FX0367. Alaska Dep. of Fish and Game, Division of Wildlife Conservation. Anchorage, AK.

- Swain, U., J. Lewis, G. Pendelton, and K. Pitcher. 1996. Movements, haulout, and diving behavior of harbor seals in southeast Alaska and Kodiak Island. Pp. 59-144 *In* Annual Report: Harbor seal investigations in Alaska. NOAA Grant NA57FX0367. Alaska Dep. of Fish and Game, Division of Wildlife Conservation. Douglas, AK.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Westlake, R. L., and G. O'Corry-Crowe. 1997. Genetic investigation of Alaskan harbor seal stock structure using mtDNA. Pp. 205-234 *In* Annual Report: Harbor seal investigations in Alaska. NOAA Grant NA57FX0367. Alaska Dep. of Fish and Game, Division of Wildlife Conservation. Anchorage, AK.
- Withrow, D. E., and T. R. Loughlin. 1995a. Abundance and distribution of harbor seals (*Phoca vitulina richardsi*) along the Aleutian Islands during 1994. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Withrow, D. E., and T. R. Loughlin. 1995b. Haulout behavior and method to estimate the proportion of harbor seals missed during molt census surveys in Alaska. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Withrow, D. E., and T. R. Loughlin. 1996. Haulout behavior and a correction factor estimate for the proportion of harbor seals missed during molt census surveys near Cordova, Alaska. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Withrow, D. E., and T. R. Loughlin. 1997. Abundance and distribution of harbor seals (*Phoca vitulina richardsi*) along the south side of the Alaska Peninsula, Shumagin Islands, Coon Inlet, Kenai Peninsula, and the Kodiak Archipelago in 1996. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Wolfe, R. J., and C. Mishler. 1993. The subsistence harvest of harbor seal and sea lion by Alaska natives in 1992. Final report for year one, subsistence study and monitor system (no. 50ABNF20055). Prepared for the NMFS by Alaska Dep. Fish and Game, Juneau, Alaska, 94 pp. + appendices.
- Wolfe, R. J., and C. Mishler. 1994. The subsistence harvest of harbor seal and sea lion by Alaska natives in 1993. Final report for year two, subsistence study and monitor system (no. 50ABNF20055). Prepared for the NMFS by Alaska Dep. Fish and Game, Juneau, Alaska, 60 pp. + appendices.
- Wolfe, R. J., and C. Mishler. 1995. The subsistence harvest of harbor seal and sea lion by Alaska natives in 1994. Final report for year three, subsistence study and monitor system (no. 50ABNF20055). Prepared for NMFS by Alaska Dept. Fish and Game, Juneau, Alaska, 69 pp. + appendices.
- Wolfe, R. J., and C. Mishler. 1996. The subsistence harvest of harbor seal and sea lion by Alaska natives in 1995. Final report for year four, subsistence study and monitor system (no. 50ABNF400080). Prepared for NMFS by Alaska Dept. Fish and Game, Juneau, Alaska, 69 pp. + appendices.
- Wolfe, R. J., and C. Mishler. 1997. The subsistence harvest of harbor seal and sea lion by Alaska natives in 1996. Technical Paper 241. Draft Final report for year five, subsistence study and monitor system (no. 50ABNF400080). Prepared for NMFS by Alaska Dept. Fish and Game, Juneau, Alaska, 70 pp. + appendices.
- Wynne, K. M., D. Hicks, and N. Munro. 1991. 1990 salmon gillnet fisheries observer programs in Prince William Sound and South Unimak Alaska. Annual Rept. NMFS/NOAA Contract 50ABNF000036. 65 pp. NMFS, Alaska Region, Office of Marine Mammals, P.O. Box 21668, Juneau, AK 99802.
- Wynne, K. M., D. Hicks, and N. Munro. 1992. 1991 Marine mammal observer program for the salmon driftnet fishery of Prince William Sound Alaska. Annual Rept. NMFS/NOAA Contract 50ABNF000036. 53 pp. NMFS, Alaska Region, Office of Marine Mammals, P.O. Box 21668, Juneau, AK 99802.