

MINKE WHALE (*Balaenoptera acutorostrata*): Alaska Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE:

In the North Pacific, minke whales occur from the Bering and Chukchi Seas south to near the Equator (Leatherwood et al. 1982). The following information was considered in classifying stock structure according to the Dizon et al. (1992) phylogeographic approach: 1) Distributional data: geographic distribution continuous, 2) Population response data: unknown; 3) Phenotypic data: unknown; and 4) Genotypic data: unknown. Based on this limited information, the International Whaling Commission (IWC) recognizes three stocks of minke whales in the North Pacific: one in the Sea of Japan/East China Sea, one in the rest of the western Pacific west of 180°N, and one in the “remainder” of the Pacific (Donovan 1991). The “remainder” stock designation reflects the lack of exploitation in the eastern Pacific and does not indicate that only one population exists in this area (Donovan 1991).

In the “remainder” area, minke whales are relatively common in the Bering and Chukchi Seas and in the inshore waters of the Gulf of Alaska (Mizroch 1992), but are not considered abundant in any other part of the eastern Pacific (Leatherwood et al. 1982, Brueggeman et al. 1990). Minke whales are known to penetrate loose ice during the summer, and some individuals venture north of the Bering Strait (Leatherwood et al. 1982). A recent survey in the central Bering Sea in July-August 1999 resulted in 20 on-effort sightings of minke whales, most of which occurred along the upper slope in waters 100-200 m deep (Moore et al., in review). In the northern part of their range minke whales are believed to be migratory, whereas they appear to establish home ranges in the inland waters of Washington and along central California (Dorsey et al. 1990). Because the “resident” minke whales from California to Washington appear behaviorally distinct from migratory whales farther north, minke whales in Alaska are considered a separate stock from minke whales in California, Oregon, and Washington. Accordingly, two stocks of minke whales are recognized in U. S. waters: 1) Alaska, and 2) California/Washington/Oregon (Fig. 34). The California/ Oregon/Washington minke whale stock is reported separately in the Stock Assessment Reports for the Pacific Region.




Figure 34. Approximate distribution of minke whales in the eastern North Pacific (shaded area).

POPULATION SIZE

No estimates have been made for the number of minke whales in the entire North Pacific. However, some information is now available on the numbers of minke whales in the Bering Sea. A visual survey for cetaceans was conducted in the central Bering Sea in July-August 1999 in cooperation with research on commercial fisheries (Moore et al., in review). The survey included 6,043 kilometers of tracklines, most of which were west of St. Matthew Island, north of the 200 m bathymetric contour, and south of the U.S./Russia Convention line. Results of this cetacean survey provide an estimated abundance of 936 minke whales (95% CI 473-1,852; CV = 0.35) in the central Bering Sea during the summer. This estimate cannot be used as an estimate of the entire Alaska stock of minke whales because only a portion of the stock's range was surveyed.

Minimum Population

At this time, it is not possible to produce a reliable estimate of minimum abundance for this stock, as current estimates of abundance are not available.

Current Population Trend

There are no data on trends in minke whale abundance in Alaska waters.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

There are no estimates of the growth rate of minke whale populations in the North Pacific (Best 1993). Hence, until additional data become available, it is recommended that the cetacean maximum net productivity rate (R_{MAX}) of 4% be employed for this stock (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) for this stock is calculated as the product of minimum population size, 0.5 maximum net productivity, and a recovery factor. Given the status of this stock is unknown, the appropriate recovery factor is 0.5 (Wade and Angliss 1997). However, because an estimate of minimum abundance is not available, it is not possible to estimate a PBR for the Alaska minke whale stock at this time.

ANNUAL HUMAN-CAUSED MORTALITY

Fishery Information

Six different commercial fisheries operating in Alaska waters within the range of the Alaska minke whale stock were monitored for incidental take by NMFS observers during 1990-99: Bering Sea (and Aleutian Islands) groundfish trawl, longline, and pot fisheries, and Gulf of Alaska groundfish trawl, longline, and pot fisheries. No minke whale mortalities were observed for any of these fisheries. In 1989, one minke whale mortality (extrapolated to 2 mortalities) was observed in the Bering Sea/Gulf of Alaska joint-venture groundfish trawl fishery, the predecessor to the current Alaska groundfish trawl fishery.

In the past, minke whales have been caught in both coastal set gillnets and offshore drift gillnets (Small and DeMaster 1995). Logbook data are available for part of 1989-1994, after which incidental mortality reporting requirements were modified. Under the new system, logbooks are no longer required; instead, fishers provide self-reports. Data for the 1994-95 phase-in period are fragmentary. After 1995, the level of reporting dropped dramatically, such that the records are considered incomplete and estimates of mortality based on them represent minimums (see Appendix 7 for details). There have been no logbook reports or self-reports of minke whales seriously injured or killed incidental to any fishery in Alaska.

The estimated annual mortality rate incidental to commercial fisheries is zero. Therefore, the annual human-caused mortality level is considered to be insignificant and approaching a zero mortality and serious injury rate.

Subsistence/Native Harvest Information

No minke whales were ever taken by the modern shore-based whale fishery in the eastern North Pacific which lasted from 1905 to 1971 (Rice 1974). Subsistence takes of minke whales by Alaska Natives are rare, but have been known to occur. Only seven minke whales are reported to have been taken for subsistence by Alaska Natives between 1930 and 1987 (C. Allison, pers. comm., International Whaling Commission, The Red House, Station Road, Histon, Cambridge, UK). The most recent harvest (2 whales) in Alaska occurred in 1989 (Anonymous 1991). Based on this information, the annual subsistence take averaged zero minke whales during the 3-year period from 1993 to 1995.

STATUS OF STOCK

Minke whales are not listed as “depleted” under the MMPA or listed as “threatened” or “endangered” under the Endangered Species Act. The greatest uncertainty regarding the status of the Alaska minke whale stock has to do with the uncertainty pertaining to the stock structure of this species in the eastern North Pacific. Because minke whales are considered common in the waters off Alaska and because the number of human-related removals is currently thought to be minimal, this stock is not considered a strategic stock. Reliable estimates of the minimum population size, population trends, PBR, and status of the stock relative to OSP are currently not available.

CITATIONS

- Anon. 1991. International Whaling Commission Report. Rep. Int. Whal. Comm. 41:1-2.
- Best, P. B. 1993. Increase rates in severely depleted stocks of baleen whales. ICES J. Mar. Sci. 50:169-186.
- Brueggeman, J. J., G. A. Green, K. C. Balcomb, C. E. Bowlby, R. A. Grotefendt, K. T. Briggs, M. L. Bonnell, R. G. Ford, D. H. Varoujean, D. Heinemann, and D. G. Chapman. 1990. Oregon-Washington marine mammal and seabird survey: Information synthesis and hypothesis formulation. U.S. Dep. Interior, Outer Continental Shelf Study, Minerals Management Service 89-0030.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. Conserv. Biol. 6:24-36.
- Donovan, G. P. 1991. A review of IWC stock boundaries. Rept. Int. Whal. Comm. (Special Issue 13):39-68.
- Dorsey, E. M., S. J. Stern, A. R. Hoelzel, and J. Jacobsen. 1990. Minke whale (*Balaenoptera acutorostrata*) from the west coast of North America: individual recognition and small scale site fidelity. Rept. Int. Whal. Comm. (Special Issue 12):357-368.
- Leatherwood, S., R. R. Reeves, W. F. Perrin, and W. E. Evans. 1982. Whales, dolphins, and porpoises of the eastern North Pacific and adjacent Arctic waters: A guide to their identification. U.S. Dep. Commer., NOAA Tech. Rept. NMFS Circular 444. 245 pp.
- Mizroch, S. A. 1992. Distribution of minke whales in the North Pacific based on sightings and catch data. Unpubl. doc. submitted to the Int. Whal. Comm. (SC/43/Mi36). 37 pp.
- Moore, S. E., J. M. Waite, L. L. Mazzuca, and R. C. Hobbs. Provisional estimates of mysticete whale abundance on the central Bering Sea shelf. In review.
- Rice, D. W. 1974. Whales and whale research in the eastern North Pacific. Pp. 170-195, *In* W. E. Schevill (ed.), The whale problem: A status report. Harvard Press, Cambridge, MA.
- Small, R. J., and D. P. DeMaster. 1995. Alaska marine mammal stock assessments 1995. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-57, 93 pp.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.