

SPERM WHALE (*Physeter macrocephalus*): North Pacific Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The sperm whale is one of the most widely distributed of any marine mammal species, perhaps only exceeded by the killer whale (Rice 1989). They feed primarily on medium-sized to large-sized squids but also take substantial quantities of large demersal and mesopelagic sharks, skates, and fishes (Rice 1989). In the North Pacific, sperm whales are distributed widely (Fig. 31), with the northernmost boundary extending from Cape Navarin (62°N) to the Pribilof Islands (Omura 1955). Females and young sperm whales usually remain in tropical and temperate waters year-round, while males are thought to move north in the summer to feed in the Gulf of Alaska, Bering Sea, and waters around the Aleutian Islands (Kasuya and Miyashita 1988). Sightings surveys conducted by NMML in the summer months between 2001 and 2006 have found sperm whales to be the most frequently sighted large cetacean in the coastal waters around the central and western Aleutian Islands (NMML unpublished data). Acoustic surveys detected the presence of sperm whales year-round in the Gulf of Alaska although they appear to be more common in summer than in winter (Mellinger et al. 2004). These seasonal detections are consistent with the hypothesis that sperm whales migrate to higher latitudes in summer and migrate to lower latitudes in winter (Whitehead and Arnborn 1987).

Discovery Mark data from the days of commercial whaling (270 recoveries) show extensive movements from U.S. and Canadian coastal waters into the Gulf of Alaska and Bering Sea (Omura and Ohsumi 1964; Ivashin and Rovnin 1967; Ohsumi and Masaki 1975; Wada 1980; Kasuya and Miyashita 1988. Rice (AFSC-NMML, retired, pers. comm.) marked 176 sperm whales during U.S. cruises from 1962-1970, mostly between 32° and 36° N off the California coast. Seven of those marked whales were recovered and the recovery locations ranged from offshore California, Oregon, British Columbia waters to the western Gulf of Alaska. A whale marked by Canadian researchers moved from near Vancouver Island, British Columbia to the Aleutian Islands near Adak. A whale marked by Japanese researchers moved from the Bering Sea just north of the Aleutians to waters off Vancouver Island, British Columbia. Based on these data, there appears to be movements along the U.S. west coast into the Gulf of Alaska and Bering Sea/Aleutian Islands region (BSAI).

The following information was considered in classifying stock structure based on the Dizon et al. (1992) phylogeographic approach: 1) Distributional data: geographic distribution continuous though data indicate three “somewhat” discrete population centers (i.e., Hawaii, west coast of the continental United States, and Alaska); 2) Population response data: unknown; 3) Phenotypic data: unknown; and 4) Genotypic data: unknown. For management purposes, the International Whaling Commission (IWC) recognizes two management units of sperm whales in the North Pacific (eastern and western). However, the IWC has not reviewed its sperm whale stock boundaries in recent years (Donovan 1991). Based on this limited information, and lacking additional data concerning population structure, sperm whales of the eastern North Pacific have been divided into three separate stocks as dictated by the U. S. waters in which they are found: 1) Alaska (North Pacific stock), 2) California/Oregon/Washington, and 3) Hawaii. The California/Oregon/Washington and Hawaii sperm whale stocks are reported separately in the Stock Assessment Reports for the Pacific Region.




Figure 31. Approximate distribution of sperm whales in the north Pacific includes deep waters south of 62°N to the equator.

POPULATION SIZE

Current and historic estimates for the abundance of sperm whales in the North Pacific are considered unreliable. Therefore, caution should be exercised in interpreting published estimates of abundance. The abundance of sperm whales in the North Pacific was reported to be 1,260,000 prior to exploitation, which by the late 1970s was estimated to have been reduced to 930,000 whales (Rice 1989). Confidence intervals for these estimates were not provided. These estimates include whales from the California/Oregon/Washington stock, for which a separate abundance estimate is currently available (see Stock Assessment Reports for the Pacific Region).

Although Kato and Miyashita (1998) believe their estimate to be upwardly biased, their preliminary analysis indicates 102,112 (CV = 0.155) sperm whales in the western North Pacific. The number of sperm whales of the North Pacific occurring within Alaska waters is unknown. As the data used in estimating the abundance of sperm whales in the entire North Pacific are over 8 years old at this time and there are no available estimates for numbers of sperm whales in Alaska waters, a reliable estimate of abundance for the North Pacific stock is not available.

Minimum Population Estimate

At this time, it is not possible to produce a reliable estimate of minimum abundance for this stock, as a current estimate of abundance is not available.

Current Population Trend

Reliable information on trends in abundance for this stock is currently not available (Braham 1992).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

A reliable estimate of the maximum net productivity rate is not currently available for the North Pacific stock of sperm whale. Hence, until additional data become available, it is recommended that the cetacean maximum net productivity rate (R_{MAX}) of 4% be employed for this stock at this time (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $PBR = N_{MIN} H 0.5R_{MAX} H F_R$. The recovery factor (F_R) for this stock is 0.1, the value for cetacean stocks which are classified as endangered (Wade and Angliss 1997). However, because a reliable estimate of minimum abundance N_{MIN} is currently not available, the PBR for this stock is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fisheries Information

In previous stock assessments, there were six commercial fisheries monitored by federal observers in Alaska that could have had incidental serious injuries or mortalities of sperm whales. In 2004, the definitions of these commercial fisheries were changed to reflect target species: these new definitions have resulted in the identification of 22 observed fisheries in the Gulf of Alaska and Bering Sea that use trawl, longline, or pot gear (69 FR 70094, 2 December 2004). Of these, there is at least one fishery that has incurred incidental serious injuries or mortalities of sperm whales, the Gulf of Alaska sablefish longline fishery. However, since 2001 there have been no observed mortalities in federally observed Alaska fisheries and therefore the minimum estimated annual mortality rate incidental to U.S. commercial fisheries is zero (Perez unpubl. ms.).

Subsistence/Native Harvest Information

Sperm whales have never been reported to be taken by subsistence hunters (Rice 1989).

Other Mortality

Sperm whales were the dominant species killed by the commercial whaling industry as it developed in the North Pacific in the years after the second World War (Mizroch and Rice 2006). Between 1946 and 1967, most of the sperm whales were caught in waters near Japan and in the Bering Sea/Aleutian Islands (BSAI) region. The BSAI catches were dominated by males. After 1967, whalers moved out of the BSAI region and began to catch even larger numbers of sperm whales further south in the North Pacific between 30° and 50° N (Mizroch and Rice 2006, Figs. 7-9). The reported catch of sperm whales taken by commercial whalers operating in the North Pacific

between 1925 and 1987 was 260,285 sperm whales, of which, 258,829 were taken between 1946 and 1987 (International Whaling Commission, BIWS catch data, February 2003 version, unpublished). This value underestimates the actual kill in the North Pacific as a result of under-reporting by U.S.S.R. pelagic whaling operations. Brownell et al. (2000) estimated that the U.S.S.R. under-reported catches during 1949-71 by as much as 60%. In addition, new information suggests that Japanese land-based whaling operations also under-reported sperm whale catches during the post-World War II era (Kasuya 1999). The last year that the U.S.S.R reported catches of sperm whales was in 1979 and the last year that Japan reported catches was in 1987 (International Whaling Commission, BIWS catch data, February 2003 version, unpublished).

Other Issues

NMFS observers aboard longline vessels targeting both sablefish and halibut have documented sperm whales feeding off longline gear in the Gulf of Alaska (Hill and Mitchell 1998, Hill et al., 1999; Perez 2006). Fishery observers recorded several instances during 1995-97 in which sperm whales were deterred by fishermen (i.e., yelling at the whales or throwing seal bombs in the water).

Annual longline surveys have been recording sperm whale predation on catch since 1998 (Sigler et al. 2003). Between 1989 and 2003, sperm whale predation on catch has occurred at 38 of the surveyed stations: all events were located in the Gulf of Alaska and none were located in the Bering Sea. The sablefish catch at the stations where predation occurs is lower than at those stations where no predation occurred. Undamaged catches may also occur when sperm whales are present; in these cases, sperm whales apparently feed off the discard. Observer records document that predation on catch is widespread in the Gulf of Alaska (Perez 2006).

STATUS OF STOCK

Sperm whales are listed as “endangered” under the Endangered Species Act of 1973, and therefore designated as “depleted” under the MMPA. As a result, this stock is classified as a strategic stock. However, on the basis of total abundance, current distribution, and regulatory measures that are currently in place, it is unlikely that this stock is in danger of extinction (Braham 1992). Reliable estimates of the minimum population, population trends, PBR, and status of the stock relative to its Optimum Sustainable Population size are currently not available, although the estimated annual rate of human-caused mortality and serious injury seems minimal for this stock. Because the PBR is unknown, the level of annual U.S. commercial fishery-related mortality that can be considered insignificant and approaching zero mortality and serious injury rate is unknown. There are no known habitat issues that are of particular concern for this stock.

CITATIONS

- Braham, H. 1992. Endangered whales: Status update. Working document presented at A Workshop on the Status of California Cetacean Stocks (SOCCS/14). 35 pp. + tables. (Available upon request - Alaska Fisheries Science Center, 7600 Sand Point Way, NE, Seattle, WA 98115).
- Brownell, R. L., A. V. Yablokov, and V. A. Zemsky. 2000. USSR pelagic catches of North Pacific sperm whales, 1949-1979: conservation implications. Pp. 123-130 *In* A. V. Yablokov and V. A. Zemsky (eds.), Soviet whaling data (1949-1979). Center for Russian Environmental Policy, Marine Mammal Council, Moscow, Russia.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Donovan, G. P. 1991. A review of IWC stock boundaries. *Rept. Int. Whal. Comm. (Special Issue 13):*39-68.
- Hill, P. S., and E. Mitchell. 1998. Sperm whale interactions with longline vessels in Alaska waters during 1997. Unpubl. doc. Submitted to *Fish. Bull., U.S.* (Available upon request - P. S. Hill, Alaska Fisheries Science Center, 7600 Sand Point Way, NE, Seattle, WA 98115).
- Hill, P. S., J. L. Laake, and E. Mitchell. 1999. Results of a pilot program to document interactions between sperm whales and longline vessels in Alaska waters. *U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-108*, 42 pp.
- Ivashin M. V., and A. A. Rovnin. 1967. Some results of the Soviet whale marking in the waters of the North Pacific. *Norsk Hvalfangst-tidende* 56:123-135.
- Kasuya, T. 1999. Examination of the reliability of catch statistics in the Japanese coastal sperm whale fishery. *J. Cetacean Res. Manage.* 1: 109-122.
- Kasuya T., and T. Miyashita. 1988. Distribution of sperm whale stocks in the North Pacific. *Sci. Rep. Whales Res. Inst.* 39: 31-75.

- Kato, H., and T. Miyashita. 1998. Current status of North Pacific sperm whales and its preliminary abundance estimates. Unpubl. report submitted to Int. Whal. Comm. (SC/50/CAWS/52). 6 pp.
- Mellinger, D. K., Stafford, K. M., and Fox, C. G. 2004. Seasonal occurrence of sperm whale (*Physeter macrocephalus*) sounds in the Gulf of Alaska, 1999-2001. *Mar. Mammal Sci.* 20(1):48-62.
- Mizroch S. A., and D. W. Rice. 2006. Have North Pacific killer whales switched prey species in response to depletion of the great whale populations? *Mar. Ecol. Prog. Ser.* 310:235-246.
- Ohsumi S., and Y. Masaki. 1975. Japanese whale marking in the North Pacific, 1963-1972. *Bull. Far Seas Fish. Res. Lab.* 12:171-219.
- Omura, H. 1955. Whales in the northern part of the North Pacific. *Nor. Hvalfangst-Tidende* 44(6):323-345.
- Omura H., and S. Ohsumi. 1964. A review of Japanese whale marking in the North Pacific to the end of 1962, with some information on marking in the Antarctic. *Norsk Hvalfangst-tidende* 53:90-112.
- Perez, M. A. 2006. Analysis of marine mammal bycatch data from the trawl, longline, and pot groundfish fisheries of Alaska, 1998-2004, defined by geographic area, gear type, and target groundfish catch species. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-167.
- Perez, M. A. Unpubl. ms. Bycatch of marine mammals by the groundfish fisheries in the U.S. EEZ of Alaska, 2005. 67 pp. Available NMML-AFSC.
- Rice, D. W. 1989. Sperm whale, *Physeter macrocephalus*. Pp. 177-233 *In* S. H. Ridgway and R. Harrison (eds.), *Handbook of Marine Mammals. Vol. 4. River Dolphins and the Larger Toothed Whales.* Academic Press, New York.
- Sigler, M.F., C. R. Lunsford, J. T. Fujioka, and S. A. Lowe. 2003. Alaska Sablefish Assessment for 2004. *In* Stock Assessment and Fishery Evaluation Report for the Groundfish Fisheries of the Bering Sea/Aleutian Islands Regions. North Pac. Fish. Mgmt. Council., Anchorage, AK, Section 3:223-292.
- Wada S. 1980. On the genetic uniformity of the North Pacific sperm whale. *Reports of the Int. Whal. Comm. Special Issue* 2:205-211.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Whitehead, H. and Arnborn, T. 1987. Social organization of sperm whale off the Galapagos Island, February-April 1985. *Can. J. Zool.* 65(4):913-919.