

## STELLER SEA LION (*Eumetopias jubatus*): Western U. S. Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

Steller sea lions range along the North Pacific Rim from northern Japan to California (Loughlin et al. 1984), with centers of abundance and distribution in the Gulf of Alaska and Aleutian Islands (Fig. 3). The species is not known to migrate, but individuals disperse widely outside of the breeding season (late May-early July), thus potentially intermixing with animals from other areas (Sease and York 2003). Despite the wide-ranging movements of juveniles and adult males in particular, exchange between rookeries by breeding adult females and males (other than between adjoining rookeries) is low (NMFS 1995).


Loughlin (1997) considered the following information when classifying stock structure based on the phylogeographic approach of Dizon et al. (1992): 1) Distributional data: geographic distribution continuous, yet a high degree of natal site fidelity and low (<10%) exchange rate of breeding animals between rookeries; 2) Population response data: substantial differences in population dynamics (York et al. 1996); 3) Phenotypic data: unknown; and 4) Genotypic data: substantial differences in mitochondrial DNA (Bickham et al. 1996). Based on this

information, two separate stocks of Steller sea lions were recognized within U. S. waters: an eastern U. S. stock, which includes animals east of Cape Suckling, Alaska (144°W), and a western U. S. stock, which includes animals at and west of Cape Suckling (Loughlin 1997, Fig. 1).

Steller sea lions that breed in Asia have been considered part of the western stock. While Steller sea lions seasonally inhabit coastal waters of Japan in the winter, breeding rookeries are currently only located in Russia (Burkanov and Loughlin 2005). Analyses of genetic data differ in their interpretation of separation between Asian and Alaskan sea lions. Based on analysis of mitochondrial DNA, Baker et al. (2005) found evidence of a genetic split between the Commander Islands (Russia) and Kamchatka that would include Commander Island sea lions within the western U.S. stock and animals west of there in an Asian stock. However, Hoffman et al. (2006) did not support an Asian/western stock split based on their analysis of nuclear microsatellite markers indicating high rates of male gene flow. All genetic analyses confirm a strong separation between western and eastern stocks, and there may be sufficient morphological differentiation to support elevating the two recognized stocks to subspecies (Phillips et al. 2011).

### POPULATION SIZE

The most recent comprehensive estimate (pups and non-pups) of abundance of the western stock of Steller sea lions in Alaska is based on aerial surveys of non-pups in June-July 2008 (Fritz et al. 2008a) and aerial and ground-based pup counts in June and July of 2009 and 2010 (DeMaster 2009; Fritz and Gelatt 2011). Data from these surveys represent actual counts of pups and non-pups at all rookeries and major haulout sites. During the 2008 aerial survey, a total of 31,245 non-pups was counted at 275 rookeries and haulout sites; 6,522 in the Gulf of Alaska and 14,723 in the Bering Sea/Aleutian Islands (Fritz et al. 2008b). A composite pup count for 2009-2010 includes counts from 172 sites in 2009, and 6 sites in 2010. There were 5,457 pups counted in the Gulf of Alaska and 5,584



**Figure 1.** Approximate distribution of Steller sea lions in the North Pacific. Major U.S. haulouts and rookeries (50 CFR 226.202, 27 August 1993) and active Asian haulouts and rookeries (Burkanov and Loughlin, 2005) are depicted (points). Black dashed line (144° W) indicates stock boundary (Loughlin 1997). Note: Haulouts and rookeries in British Columbia are not shown.

pups counted in the Bering Sea/Aleutian Islands for a total of 11,041 for the stock in Alaska. Combining the pup count data from 2009-2010 (11,041) and non-pup count data from 2008 (31,245) results in a minimum abundance estimate of 42,286 Steller sea lions in the western U.S. stock in 2008-2010.

An estimate of the total population size of western Steller sea lion in Alaska may be obtained by multiplying the best estimate of total pup production (11,041) by 4.5 (Calkins and Pitcher 1982), which equals 49,685. This would not be a minimum abundance estimate since it is based on extrapolating total population size from pup counts based on survival and fecundity estimates in a life table. The 4.5 multiplier may not be appropriate for use in estimating the abundance of the western stock, as it is based on a life history table using age-specific fecundity and survival for the stable, mid-1970s population. The demographics of central Gulf of Alaska populations suggest that these rates have changed considerably since the mid-1970s (Holmes and York 2003; Holmes et al. 2007).

Holmes and York (2003) and Holmes et al. (2007) estimated changes in adult and juvenile survival and natality in the female segment of the population that were consistent with time series of pup and non-pup counts, and changes in the juvenile proportion of the population in the central Gulf of Alaska (Kodiak archipelago). They found that the rapid decline of the central Gulf sea lion population in the 1980s was associated with a large drop in juvenile survival and smaller declines in adult survival and natality. As the rate of population decline lessened in the 1990s, rates of juvenile and adult survival increased to pre-decline levels in the 1998-2004 period. Rates of natality, however, continued to decline throughout the 1990s and into the 2000s. Thus, the authors concluded that factors that caused the population decline (those contributing to lower rates of juvenile survival) were likely quite different from those that are now affecting recovery (those contributing to lower reproductive rates of adult females).


In 2006-2009, over 18,000 Steller sea lions were counted in Russia. Methods used to survey Steller sea lions in Russia differ from those used in Alaska, with less use of aerial photography and more use of skiff surveys and ground counts. Burkanov and Loughlin (2005) estimated that the size of the Steller sea lion population (pups and non-pups) in Russia was 16,000 in 2005. Data collected since then indicate that Steller sea lion numbers in the Kuril Islands and the Sea of Okhotsk increased while those in the western Bering Sea, eastern Kamchatka and Commander Islands have remained stable or declined.

### Minimum Population Estimate

The 2008 count of non-pups (31,245) plus the number of pups in 2009-2010 (11,041) is 42,286, which will be used as the minimum population estimate ( $N_{MIN}$ ) for the U.S. portion of the western stock of Steller sea lion (Wade and Angliss 1997). This is considered a minimum estimate because it has not been corrected to account for animals that were at sea during the surveys.

### Current Population Trend

The first reported trend counts (an index to examine population trends) of Steller sea lions in Alaska were made in 1956-60. Those counts indicated that there were at least 140,000 (no correction factors applied) sea lions in the Gulf of Alaska and Aleutian Islands (Merrick et al. 1987). Subsequent surveys indicated a major population decrease, first detected in the eastern Aleutian Islands in the mid-1970s (Braham et al. 1980). Counts from 1976 to 1979 indicated about 110,000 sea lions (no correction factors applied, Table 1). The decline appears to have spread eastward to the Kodiak Island area during the late 1970s and early 1980s, and then westward to the central and western Aleutian Islands during the early and mid-1980s (Merrick et al. 1987, Byrd 1989). The greatest declines since the 1970s occurred in the eastern Aleutian Islands and western Gulf of Alaska, but declines also occurred in the central Gulf of Alaska and central Aleutian Islands. Counts of Steller sea lions at trend sites for the western U. S. stock decreased 40% from 1991 to 2000 (Table 1), an average annual decline of 5.4% (Loughlin and York 2000).



**Figure 2.** Counts of adult and juvenile Steller sea lions at rookery and haulout trend sites throughout the range of the western U.S. stock in Alaska, 1990-2008. Correction factor applied to 2004 and 2008 counts for film format differences (Fritz and Stinchcomb 2005).

Counts of non-pup Steller sea lions at trend sites for the western U.S. stock increased 5.5% from 2000 to 2002, and at a similar rate between 2002 and 2004 (Table 1, Fig. 2). These were the first region-wide increases for the western stock since standardized surveys began in the 1970s. Aerial surveys for non-pup Steller sea lions were conducted in 2006 and 2007, but were incomplete due to a court-ordered cessation of research that caused a delay to the start of the survey in 2006, and loss of survey days due to bad weather and aircraft maintenance requirements in both years. Although some trend sites were not surveyed in both 2006 and 2007, available data indicated that the size of the adult and juvenile portion of the western Steller sea lion population throughout much of its range (Cape St. Elias to Tanaga Island, 145°-178° W) in Alaska remained largely unchanged between 2004 (N=23,107) and 2007 (N=23,118) (Fritz et al. 2008a). Results of the aerial survey conducted in 2008 (Fritz et al. 2008b) confirmed that the recent (2004-2008) overall trend in the western population of adult and juvenile Steller sea lions in Alaska is stable. There continues to be considerable regional variability in recent (2004-2008) trends (percentages listed below are % change between years):

- the population in the eastern Aleutian Islands is the only one that has consistently increased from 2004-2008 (+7%);
- the populations in the central and western Aleutian Islands declined (-30% and -16%, respectively);
- the populations in the central and western Gulf of Alaska increased between 2004 and 2007, but declined slightly between 2007 and 2008; and
- non-pup counts in the eastern Gulf of Alaska increased by 35%, with some of this increase likely related to timing of the 2008 survey (earlier than usual) and seasonal movement of animals into this area from the central Gulf and Southeast Alaska (eastern stock).

Counts in the area from the central Gulf of Alaska through the western Aleutian Islands (85% of the 2008 population) declined slightly (-1%) between 2004 and 2008, indicating that the overall increase observed between 2004 and 2008 (3%) was entirely in the eastern Gulf of Alaska. The increase in the eastern Gulf of Alaska may be partially explained by movement of animals from the eastern stock, since counts at index sites in Southeast Alaska were approximately 1,200 lower in 2008 than in 2002, despite the overall 3% per year increase in the Steller sea lion population observed in Southeast Alaska through 2005 (NMFS 2008).

In 2009 (DeMaster 2009), NMML conducted a non-pup survey in late June ('late' compared with the 'early' 2008 survey) to further investigate seasonal movement of sea lions in the northern Gulf of Alaska and how it could affect trend counts in both the eastern and western stocks. In the 'late' 2009 compared to the 'early' 2008 survey, NMFS counted:

- 2,636 more non-pups on all sites in SE Alaska;
- 812 fewer non-pups on all sites in the eastern Gulf of Alaska; and
- 404 more non-pups on 28 of the 33 trend sites surveyed consistently since 1991 in the central Gulf of Alaska.

These results are consistent with the hypothesis proposed in 2008 (Fritz et al. 2008a) that seasonal movement into the eastern Gulf of Alaska may have affected non-pup trend analyses in this area as well as for the western DPS as a whole. DeMaster (2009) estimated the number of sea lions from SE Alaska (eastern stock) and from the central Gulf of Alaska (western stock) that were counted in the eastern Gulf of Alaska (western stock) in 2008 by comparing actual to predicted 2008 sub-area totals based on the 2000-2009 overall trends. These analyses indicated that approximately 570 animals from the eastern stock may have been counted within the range of the western stock in 2008. If 570 non-pups are subtracted from the 2008 total, the overall western stock increase between 2004 and 2008 is reduced from 3% to 1%. The 2009 non-pup survey results in the northern Gulf of Alaska supports the earlier conclusion that the increase observed between 2000 and 2004 in the size of the western stock of Steller sea lion did not continue, and that the population was generally stable between 2004 and 2008.

**Table 1.** Counts of adult and juvenile Steller sea lions observed at rookery and haulout trend sites surveyed consistently since the late 1970s by year and geographical area for the western U. S. stock (NMFS 1995, Sease et al. 2001, Fritz et al. 2008b, NMFS 2008). Counts from 1976 to 1979 (NMFS 1995) were combined to produce complete regional counts that are comparable to the 1990-2008 data. Data from 2004 and 2008 reflect a 3.64% reduction from actual counts to account for improvements in survey protocol in 2004 relative to previous years (Fritz and Stinchcomb 2005).

Area	late 1970s	1990	1991	1992	1994	1996	1998	2000	2002	2004	2008
Gulf of Alaska	65,296	16,409	14,598	13,193	11,862	9,784	8,937 <sup>1</sup>	7,995	9,087	8,993	10,931
Bering Sea/Aleutians	44,584	14,116	14,807	14,106	12,274	12,426	11,501	10,330	10,253	11,507	10,559
Total	109,880	30,525	29,405	27,299	24,136	22,210	20,438 <sup>1</sup>	18,325	19,340	20,500	21,489

<sup>1</sup>Identifies 637 non-pups counted at six trend sites in 1999 in the eastern Gulf of Alaska which were not surveyed in 1998.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

There are no estimates of maximum net productivity rate for Steller sea lions. Hence, until additional data become available, it is recommended that the theoretical maximum net productivity rate ( $R_{MAX}$ ) for pinnipeds of 12% be employed for this stock (Wade and Angliss 1997).

### POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor:  $PBR = N_{MIN} \times 0.5R_{MAX} \times F_R$ . The recovery factor ( $F_R$ ) for this stock is 0.1, the default value for stocks listed as “endangered” under the Endangered Species Act (Wade and Angliss 1997). Thus, for the U.S. portion of the western stock of Steller sea lions,  $PBR = 253$  animals ( $42,286 \times 0.06 \times 0.1$ ).

The PBR levels for some stocks of marine mammals in the U.S. with an obvious declining trend have been called “undetermined” (e.g., PBR levels for Cook Inlet beluga whales, Hawaiian monk seals); this has not been proposed for the western stock of Steller sea lions. The PBR management approach was developed with the assumption that direct human-related mortalities would be the primary reason for observed declines in abundance for marine mammal stocks in U. S. waters. For at least this stock, this assumption seems unwarranted. Because direct human-related mortalities are at a low level and are unlikely to either be responsible for the decline or to contribute substantially towards extinction risk, calling the PBR level “undetermined” is unnecessarily conservative for this population of over 40,000 animals.

### ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

#### Fisheries Information

Until 2003, there were six different federally regulated commercial fisheries in Alaska that could have interacted with Steller sea lions. These fisheries were monitored for incidental mortality by fishery observers. As of 2003, changes in fishery definitions in the List of Fisheries have resulted in separating these 6 fisheries into 22 fisheries (69 FR 70094, 2 December 2004). This change does not represent a change in fishing effort, but provides managers with better information on the component of each fishery that is responsible for the incidental serious injury or mortality of marine mammal stocks in Alaska. Between 2007-2009, there were incidental serious injuries and mortalities of western Steller sea lions in the following fisheries: Bering Sea/Aleutian Islands Atka mackerel trawl, Bering Sea/Aleutian Islands flatfish trawl, Bering Sea/Aleutian Islands Pacific cod trawl, Bering Sea/Aleutian Islands pollock trawl, Gulf of Alaska Pacific cod trawl, Gulf of Alaska pollock trawl, Bering Sea/Aleutian Islands Pacific cod longline, and Gulf of Alaska Pacific cod longline (Table 2).

Observers also monitored the Prince William Sound salmon drift gillnet fishery in 1990 and 1991, recording 2 mortalities in 1991, extrapolated to 29 (95% CI: 1-108) kills for the entire fishery (Wynne et al. 1992). No mortalities were observed during 1990 for this fishery (Wynne et al. 1991), resulting in a mean kill rate of 14.5 (CV = 1.0) animals per year for 1990 and 1991. In 1990, observers boarded 300 (57.3%) of the 524 vessels that fished in the Prince William Sound salmon drift gillnet fishery, monitoring a total of 3,166 sets, or roughly 4% of the estimated number of sets made by the fleet. In 1991, observers boarded 531 (86.9%) of the 611 registered vessels and monitored a total of 5,875 sets, or roughly 5% of the estimated sets made by the fleet (Wynne et al. 1992). The Alaska Peninsula and Aleutian Islands salmon drift gillnet fishery was also monitored during 1990 (roughly 4% observer coverage) and no Steller sea lion mortalities were observed. It is not known whether these incidental mortality levels are representative of the current incidental mortality levels in these fisheries.

An observer program for the Cook Inlet salmon set and drift gillnet fisheries was implemented in 1999 and 2000 in response to the concern that there may be significant numbers of marine mammal injuries and mortalities that occur incidental to these fisheries. Observer coverage in the Cook Inlet drift gillnet fishery was 1.75% and 3.73% in 1999 and 2000, respectively. The observer coverage in the Cook Inlet set gillnet fishery was 7.3% and 8.3% in 1999 and 2000, respectively (Manly 2006). There were no mortalities of Steller sea lions observed in the set or drift gillnet fisheries in either 1999 or 2000 (Manly 2006). An observer program conducted for a portion of the Kodiak drift gillnet fishery in 2002 did not observe any serious injuries or mortalities of Steller sea lions, although Steller sea lions were frequently observed in the vicinity of the gear (Manly et al. 2003).

Combining the mortality estimates from the Bering Sea and Gulf of Alaska groundfish trawl and Gulf of Alaska longline fisheries presented above (14.6) with the mortality estimate from the Prince William Sound salmon drift gillnet fishery (14.5) results in an estimated mean annual mortality rate in the observed fisheries of 29.1 (CV = 0.50) sea lions per year from this stock (Table 2).

**Table 2.** Summary of incidental mortality of Steller sea lions (western U. S. stock) due to fisheries from 2007 through 2009 (or most recent data available) and calculation of the mean annual mortality rate. Mean annual mortality in brackets represents a minimum estimate from stranding data. The most recent 3 years of available data are used in the mortality for a particular fishery. N/A indicates that data are not available. Details of how percent observer coverage is measured is included in Appendix 6.

Fishery name	Years	Data type	Observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Bering Sea/Aleutian Is. Atka mackerel trawl	2007	obs	94	0	0	0
	2008	data	100	0	0	
	2009		99	0	0	
Bering Sea/Aleutian Is. flatfish trawl	2007	obs	72	3	3.91	5.98 (CV = 0.11)
	2008	data	100	11	11.02	
	2009		100	3	3.01	
Bering Sea/Aleutian Is. Pacific cod trawl	2007	obs	52	1	1.51	0.50 (CV = 0.58)
	2008	data	56	0	0	
	2009		64	0	0	
Bering Sea/Aleutian Is. pollock trawl	2007	obs	85	2	2.19	6.18 (CV = 0.15)
	2008	data	85	8	9.21	
	2009		86	6	7.14	
Gulf of Alaska pollock trawl	2007	obs	27	0	0	0
	2008	data	34	0	0	
	2009		43	0	0	
Bering Sea/Aleutian Is. Pacific cod longline	2007	obs	63	0	0	0
	2008	data	63	0	0	
	2009		61	0	0	
Gulf of Alaska Pacific cod longline	2007	obs	45	0	0	0.97 (CV = 0.81)
	2008	data	32	1	2.92	
	2009		43	0	0	
Prince William Sound salmon drift gillnet	1990-1991	obs data	4-5%	0 2	0 29	14.5 (CV = 1.0)
	1990	obs data	3%	0	0	
Alaska Peninsula/Aleutian Islands salmon drift gillnet	1990	obs data	4%	0	0	0
Cook Inlet salmon set gillnet <sup>1</sup>	1999-2000	obs data	2-5%	0 0	0, 0	0
	1999-2000	obs data	2-5%	0 0	0, 0	
Kodiak Island salmon set gillnet	2002	obs data	6.0%	0	0	0

Fishery name	Years	Data type	Observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Observer program total						28.13 (CV = 0.50)
				<b>Reported mortalities</b>		
Alaska sport salmon troll (non-commercial)	2005-2009	strand	N/A	0, 0, 0, 0, 1	N/A	[0.2]
Miscellaneous fishing gear	2005-2009	strand	N/A	0, 0, 0, 0, 1	N/A	[0.2]
Minimum total annual mortality						28.53 (CV = 0.50)

<sup>1</sup>Data from the 1999 Cook Inlet observer program are preliminary.

Reports from the NMFS stranding database of Steller sea lions entangled in fishing gear or with injuries caused by interactions with gear are another source of mortality data. During the 5-year period from 2005 to 2009, there were three confirmed fishery-related Steller sea lion stranding in the range of the western stock. One sighting involved a Steller sea lion that was reported to be in bad body condition and observed with a flasher lure hanging from its mouth; it was believed to have the hooks inside the mouth (Table 2). The other two events involved one animal found on a Bering Sea/ Aleutian Islands pollock trawl vessel while offloading the catch, which is accounted for in the estimated mortality for this fishery, and one animal entangled in unidentified gear on the Pribilof Islands. Fishery-related strandings during 2005-2009 result in an estimated annual mortality of 0.4 animals from this stock. This estimate is considered a minimum because not all entangled animals strand and not all stranded animals are found or reported. Steller sea lions reported in the stranding database as shot are not included in this estimate, as they may result from animals struck and lost in the Alaska Native subsistence harvest.

NMFS studies using satellite tracking devices attached to Steller sea lions suggest that they rarely go beyond the U.S. Exclusive Economic Zone into international waters. Given that the high-seas gillnet fisheries have been prohibited and other net fisheries in international waters are minimal, the probability that Steller sea lions are taken incidentally in commercial fisheries in international waters is very low. NMFS concludes that the number of Steller sea lions taken incidental to commercial fisheries in international waters is insignificant.

The minimum estimated mortality rate incidental to U. S. commercial fisheries is 28.5 sea lions per year, based on observer data (28.1) and stranding data (0.4) where observer data were not available. Observer data on state fisheries dates as far back as 1990; however, these are the best data available to estimate takes in these fisheries. No observers have been assigned to several fisheries that are known to interact with this stock making the estimated mortality a minimum estimate.

### Subsistence/Native Harvest Information

Information on the subsistence harvest of Steller sea lions comes via two sources: the Alaska Department of Fish and Game (ADFG) and the Ecosystem Conservation Office (ECO) of the Aleut Community of St. Paul. The ADFG conducted systematic interviews with hunters and users of marine mammals in approximately 2,100 households in about 60 coastal communities within the range of the Steller sea lion in Alaska (Wolfe et al. 2005). The interviews were conducted once per year in the winter (January to March), and covered hunter activities for the previous calendar year. As of 2009, data on community subsistence harvests are no longer being collected. Therefore, the most recent 5-years of data (2004-2008) will be retained and used for estimating an annual mortality estimate for all areas except St. Paul. Data from St. Paul are still being collected and will be updated with the most recent 5-year period available. The ECO collects data on the harvest in near real-time on St. Paul Island, and records hunter activities within 36 hours of the harvest (Zavadil et al. 2010). Information on subsistence harvest levels is provided in Table 3a; data from ECO (e.g., Zavadil et al. 2010) are relied upon as the source of data for St. Paul Island and all other data are from the ADFG (e.g., Wolfe et al. 2005).

The mean annual subsistence take from this stock over the 5-year period from 2004 through 2008, combined with the mean take over the 2005-2009 period from St. Paul, was 198 Steller sea lions/year (Table 3a).

**Table 3a.** Summary of the subsistence harvest data for the western U. S. stock of Steller sea lions. As of 2009, data on community subsistence harvests are no longer being collected. Therefore, the most recent 5-years of data (2004-2008) will be retained and used for estimating an annual mortality estimate for all areas except St. Paul. Data from St. Paul are still being collected and will be updated with the most recent 5-year period available (2005-2009).

Year	All areas except St. Paul Island			St. Paul Island	
	Number harvested	Number struck and lost	Total	Number harvested + struck and lost	Total take
2004	136.8	49.1	185.9 <sup>1</sup>		
2005	153.2	27.6	180.8 <sup>2</sup>	22 <sup>6</sup>	203
2006	114.3	33.1	147.4 <sup>3</sup>	26 <sup>7</sup>	173
2007	165.7	45.2	210.9 <sup>4</sup>	34 <sup>8</sup>	245
2008	114.7	21.6	136.3 <sup>5</sup>	22 <sup>9</sup>	158
2009	N/A	N/A	N/A	26 <sup>10</sup>	N/A
Mean annual take	136.9	35.3	172.3	26	198

<sup>1</sup>Wolfe et al. 2005; <sup>2</sup>Wolfe et al. 2006; <sup>3</sup>Wolfe et al. 2008; <sup>4</sup>Wolfe et al. 2009a; <sup>5</sup>Wolfe et al. 2009b; <sup>6</sup>Lestenkof and Zavidil 2006; <sup>7</sup>Lestenkof et al. 2007; <sup>8</sup>Lestenkof et al. 2008, <sup>9</sup>Jones 2009, <sup>10</sup>Zavidil 2010.

### Other Mortality

Illegal shooting of sea lions was thought to be a potentially significant source of mortality prior to the listing of sea lions as “threatened” under the U.S. Endangered Species Act (ESA) in 1990. Such shooting has been illegal since the species was listed as threatened. (Note: the 1994 Amendments to the MMPA made intentional lethal take of any marine mammal illegal except for subsistence take by Alaska Natives or where imminently necessary to protect human life). Records from NMFS enforcement indicate that there were two cases of illegal shootings of Steller sea lions in the Kodiak area in 1998, both of which were successfully prosecuted (NMFS, Alaska Enforcement Division). There have been no cases of successfully prosecuted illegal shootings between 1999 and 2003 (NMFS, Alaska Enforcement Division).

Mortalities may occasionally occur incidental to marine mammal research activities authorized under MMPA permits issued to a variety of government, academic, and other research organizations. Between 2003-2007, there was a total of 3 mortalities resulting from research on the western stock of Steller sea lions, which results in an average of 0.6 mortalities per year from this stock (Tammy Adams, Permits, Conservation, and Education Division, Office of Protected Resources, NMFS, 1315 East-West Highway, Silver Spring, MD 20910).

### STATUS OF STOCK

The current annual level of incidental U. S. commercial fishery-related mortality (28.5) exceeds 10% of the PBR (25) and, therefore, cannot be considered insignificant and approaching a zero mortality and serious injury rate. Based on available data, the estimated annual level of total human-caused mortality and serious injury (28.5 + 198 + 0.6 = 227.1) is below the PBR level (253) for this stock. The western U. S. stock of Steller sea lion is currently listed as “endangered” under the ESA, and therefore designated as “depleted” under the MMPA. As a result, the stock is classified as a strategic stock. However, given that the population has declined for unknown reasons that are not explained by the level of direct human-caused mortality, there is no reason to believe that limiting those mortalities to the level of the PBR will reverse the decline, if in fact the population is still declining.

The slight increase in the population estimate and PBR level should be interpreted and applied with caution. The increase in number of nonpups in the summer 2008 aerial survey may be attributable to an increase in numbers of eastern Steller sea lions hauled out in the eastern Gulf of Alaska at the time the aerial survey was conducted. A concurrent decrease in numbers in the eastern Steller sea lion stock counts occurred and NMFS is currently investigating the possibility that the increase in counts in the eastern Gulf of Alaska was due to seasonal movements of eastern Steller sea lion stock animals rather than recruitment into the stock.

### Habitat Concerns

The decline in the western U. S. stock of Steller sea lion caused a change in the listing status of the stock in 1997 from “threatened” to “endangered” under the U. S. Endangered Species Act of 1973. Survey data collected since 2000 suggest that the decline has slowed or stopped in some portions of the range of the western U. S. stock, but continues in others. Many factors have been suggested as causes of the steep decline observed in the 1980s,

(e.g., competitive effects of fishing, environmental change, disease, killer whale predation, incidental take, illegal and legal shooting). Decreases in rates of survival, particularly for juveniles, were associated with the steep 1980s declines (Holmes et al. 2007). Factors causing direct mortality were likely the most important. The slowing of the decline in the 1990s, and the periods of increase and stability observed between 2000 and 2008 were associated with increases in survival of both adults and juveniles, but also with continuation of a chronic decline in reproductive rate that may have been initiated in the early 1980s (Pitcher et al. 1998, Holmes et al. 2007). Nutritional stress related to competition with commercial fisheries or environmental change, along with predation by killer whales, have been identified as potentially important threats to recovery (NMFS 2008). Additional potential threats to Steller sea lion recovery are shown in Table 3b.

**Table 3b.** Potential threats and impacts to Steller sea lion recovery and associated references. Threats and impact to recovery as described by the Draft Steller Sea Lion Recovery Plan (NMFS 2008). Reference examples identify research related to corresponding threats and may or may not support the underlying hypotheses.

Threat	Impact on Recovery	Reference Examples
Environmental variability	Potentially high	Fritz and Hinckley 2005, Trites and Donnelly 2003
Competition with fisheries	Potentially high	Dillingham et al. 2006, Fritz and Brown 2005, Hennen 2004, Fritz and Ferrero 1998
Predation by killer whales	Potentially high	DeMaster et al. 2006, Trites et al. 2007, Williams et al. 2004, Springer et al. 2003
Toxic substances	Medium	Albers and Loughlin 2003, Lee et al. 1996, Calkins et al. 1994
Incidental take by fisheries	Low	Perez 2006, Nikulin and Burkanov 2000, Wynne et al. 1992
Subsistence harvest	Low	Wolfe et al. 2005, Loughlin and York 2000, Haynes and Mishler 1991
Illegal shooting	Low	NMFS 2001, Loughlin and York 2000
Entanglement in marine debris	Low	Calkins 1985
Disease and parasitism	Low	Burek et al. 2005
Disturbance from vessel traffic and tourism	Low	Kucey and Trites 2006
Disturbance or mortality due to research activities	Low	Atkinson et al. 2008, Kucey and Trites 2006, Kucey 2005, Loughlin and York 2000, Calkins and Pitcher 1982

A number of management actions were implemented between 1990 and 1998 to promote the recovery of the western U. S. stock of Steller sea lions, including 3 nautical mile (nmi) no-entry zones around rookeries, prohibition of groundfish trawling within 10-20 nmi of certain rookeries, and spatial and temporal allocation of Gulf of Alaska pollock and Aleutian Island Atka mackerel total allowable catch. In 2000, NMFS issued a Biological Opinion (BO) on effects of the groundfish fisheries in the Bering Sea/Aleutian Islands and Gulf of Alaska regions on listed species. In this BO, NMFS determined that the continued prosecution of the groundfish fisheries as described in the Fishery Management Plan for Bering Sea/Aleutian Islands Groundfish and in the Fishery Management Plan for Gulf of Alaska Groundfish was likely to jeopardize the continued existence of the western population of Steller sea lion and to adversely modify critical habitat. NMFS also identified several other factors that could contribute to the decline of the population, including a shift in the large-scale weather regime and predation. To avoid jeopardy, NMFS identified a Reasonable and Prudent Alternative that included components such as 1) adoption of a more precautionary rule for setting “global” harvest limits, 2) extension of 3 nmi protective zones around rookeries and haulouts not currently protected, 3) closures of many areas around rookeries and haulouts to 20 nmi, 4) establishment of four seasonal and area catch limits, and 5) establishment of a procedure (“fishing in proportion to biomass”) for setting seasonal catch limits on removal levels in critical habitat based on the biomass of the target species residing in critical habitat.

In 2001, NMFS developed a programmatic SEIS to consider the impacts on Steller sea lions of different management regimes for the Alaska groundfish fisheries. A committee composed of 21 members from fishing groups, processor groups, Alaska communities, environmental advocacy groups, and NMFS representatives met to recommend conservation measures for Steller sea lions and to develop a "preferred alternative" for the SEIS. Although consensus was not reached, a "preferred alternative" was identified and included in the SEIS. The

preferred alternative included complicated, area-specific management measures (e.g., area restrictions and closures) designed to reduce direct and indirect interactions between the Atka mackerel, pollock, and Pacific cod fisheries and Steller sea lions, particularly in waters within 10 nmi of haulouts and rookeries. The suite of conservation measures, which were implemented in 2002, were developed after working with the: 1) State of Alaska to explore whether there are potential adverse effects of state fisheries on Steller sea lions, and 2) the North Pacific Fishery Management Council (Council) to further minimize overcapitalization of fisheries and concentration of fisheries in time and space. The 2002 suite of conservation measures also removed the broad prohibition of fishing with trawl gear within 10 (or 20) nmi of rookeries in the western stock in U.S. waters, and did not apply the “fishing in proportion to biomass” procedure for regulating seasonal catch for the three Steller sea lion prey species in the same manner as was initially applied in the 2000 BO. All Steller sea lion-fishery management measures were reviewed in a draft programmatic, status quo ESA Biological Opinion on the effects of groundfish fisheries on listed species released for public review in August 2010 (NMFS 2010). NMFS concluded that the groundfish fisheries in the Bering Sea/Aleutian Islands area and in the Gulf of Alaska, as currently managed (as of 2010) were likely to jeopardize the continued existence (recovery) and adversely modify the critical habitat of the western stock of Steller sea lion. NMFS has proposed the following new measures (reasonable and prudent alternatives to the status quo suite of fishery management regulations) that mitigate jeopardy and adverse modification: closure of the western Aleutian Islands region (170°-177°E) region to directed fishing for Atka mackerel and Pacific cod, and additional measures in the central Aleutian Islands (170°W-177°E) to reduce catches of Atka mackerel and Pacific cod in critical habitat and disperse the fisheries temporally and spatially.

NMFS reconstituted the Steller Sea Lion Recovery Team in 2002 to write a revised recovery plan for the eastern and western U.S. stocks. The Team’s draft plan was reviewed by five independent reviewers in February 2006, prior to its delivery to NMFS, who then released the Plan for public review in May 2006. NMFS addressed the peer and public review comments and released the second draft Plan for another round of public and independent peer (one by the Council of Independent Experts and another commissioned by the Council) review in May 2007. NMFS released the final recovery plan in March 2008 (NMFS 2008). The de-listing criteria approved by NMFS for the western stock of Steller sea lion are:

1. The population for the U.S. region of this [stock] has increased (statistically significant) for 30 years (at an average annual growth rate of 3%), based on counts of non-pups (i.e., juveniles and adults). Based on an estimated population size of about 42,500 animals in 2000, this would represent approximately 103,000 animals in 2030.
2. The trends in non-pups in at least 5 of the 7 sub-regions are stable or increasing, consistent with the trend observed under criterion #1. The population trend in any two adjacent sub-regions can not be declining significantly. The population trend in any subregion cannot have declined by more than 50%. The 7 sub-regions are:
  - a. Eastern Gulf of Alaska (US)
  - b. Central Gulf of Alaska (US)
  - c. Western Gulf of Alaska (US)
  - d. Eastern Aleutian Islands (including the eastern Bering Sea) (US)
  - e. Central Aleutian Islands (US)
  - f. Western Aleutian Islands (US)
  - g. Russia/Asia
3. The ESA listing factor criteria are met.

## CITATIONS

- Albers, P. H., and T. R. Loughlin. 2003. Effects of PAHs on marine birds, mammals, and reptiles. Pp. 243-261 *In*: P. E. T. Douben (ed.) PAHs: An ecotoxicological perspective. John Wiley and Sons, London.
- Atkinson, S., D. P. DeMaster, and D. G. Calkins. 2008. Anthropogenic causes of the western Steller sea lion *Eumetopius jubatus* population decline and their threat to recovery. *Mammal Rev.* 38(1):1-18.
- Baker, A. R., T. R. Loughlin, V. Burkanov, C. W. Matson, T. G. Trujillo, D. G. Calkins, J. K. Wickliffe, and J. W. Bickham. 2005. Variation of mitochondrial control region sequences of Steller sea lions: the three-stock hypothesis. *J. Mammal.* 86:1075-1084.
- Bickham, J. W., J. C. Patton, and T. R. Loughlin. 1996. High variability for control-region sequences in a marine mammal: Implications for conservation and biogeography of Steller sea lions (*Eumetopias jubatus*). *J. Mammal.* 77:95-108.

- Braham, H. W., R. D. Everitt, and D. J. Rugh. 1980. Northern sea lion decline in the eastern Aleutian Islands. *J. Wildl. Manage.* 44:25-33.
- Burek, K. A., F. M. D. Gulland, G. Sheffield, K. B. Beckmen, E. Keyes, T. R. Spraker, A. W. Smith, D. E. Skilling, J. F. Evermann, J. L. Stott, J. T. Saliki, and A. W. Trites. 2005. Infectious disease and the decline of the Steller sea lions (*Eumetopias jubatus*) in Alaska, USA: insights from serologic data. *J. Wildl. Dis.* 41(3):512-524.
- Burkanov, V., and T. R. Loughlin. 2005. Distribution and abundance of Steller sea lions on the Asian coast, 1720's – 2005. *Mar. Fish. Rev.* 67(2):1-62.
- Byrd, G. V. 1989. Observations of northern sea lions at Ugamak, Buldir, and Agattu Islands, Alaska in 1989. Unpubl. rep., U.S. Fish and Wildlife Service. Alaska Maritime National Wildlife Refuge, P.O. Box 5251, NSA Adak, FPO Seattle, WA 98791.
- Calkins, D. G., and K. W. Pitcher. 1982. Population assessment, ecology and trophic relationships of Steller sea lions in the Gulf of Alaska. Environmental Assessment of the Alaskan Continental Shelf. Final reports 19:455-546.
- Calkins, D. G. 1985. Steller sea lion entanglement in marine debris. Pp. 308-314 *In* R. S. Shomura and H. O. Yoshida (editors), Proceedings of the workshop on the fate and impact of marine debris, 27-29 November 1984, Honolulu, Hawaii. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFC-54.
- Calkins, D. G., E. Becker, T. R. Spraker, and T. R. Loughlin. 1994. Impacts on Steller sea lions. Pp. 119-139 *In* T. R. Loughlin (ed.), *Marine Mammals and the Exxon Valdez*. Academic Press, N.Y.
- DeMaster, D. P. 2009. Aerial Survey of Steller Sea Lions in Alaska, June-July 2009 and Update on the Status of the Western Stock in Alaska. Memorandum to D. Mecum, K. Brix and L. Rotterman, December 2, 2009. Available AFSC, National Marine Mammal Laboratory, NOAA, NMFS 7600 Sand Point Way NE, Seattle WA 98115.
- DeMaster, D. P., A. W. Trites, P. Clapham, S. Mizroch, P. Wade, R. J. Small, and J. V. Hoef. 2006. The sequential megafaunal collapse hypothesis: Testing with existing data. *Prog. Oceanogr.* 68(2-4): 329-342.
- Dillingham, P. W., J. R. Skalski, and K. E. Ryding. 2006. Fine-scale geographic interactions between Steller sea lion (*Eumetopias jubatus*) trends and local fisheries. *Can. J. Fish. Aquat. Sci.* 63:107-119.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Fritz, L. W., and Ferrero, R. C. 1998. Options in Steller sea lion recovery and groundfish fishery management. *Biosphere Conserv.* 1(1): 7-19.
- Fritz, L. W., and E. S. Brown. 2005. Survey-and fishery-derived estimates of Pacific cod (*Gadus macrocephalus*) biomass: implications for strategies to reduce interactions between groundfish fisheries and Steller sea lions (*Eumetopias jubatus*). *Fish. Bull.* 103:501-515.
- Fritz, L. W. and T. Gelatt. 2011. Surveys of Steller Sea Lions in Alaska, June-July 2010. Memorandum to the Record, NMFS Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle WA 98115. <http://www.afsc.noaa.gov/nmml/PDF/SSL-Survey-memo-2010.pdf>.
- Fritz, L. W., and S. Hinckley. 2005. A critical review of the regime shift “junk food”- nutritional stress hypothesis for the decline of the western stock of Steller sea lion. *Mar. Mamm. Sci.* 21(3):476-518.
- Fritz, L. W., and C. Stinchcomb. 2005. Aerial, ship and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in the western stock in Alaska, June and July 2003 and 2004. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-153, 56 p.
- Fritz, L., M. Lynn, E. Kunisch, and K. Sweeney. 2008a. Aerial, ship, and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in the western stock in Alaska, June and July 2005-2007. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-183, 70 p.
- Fritz, L. W., K. Sweeney, C. Gudmundson, T. Gelatt, M. Lynn and W. Perryman. 2008b. Survey of Adult and Juvenile Steller Sea Lions, June-July 2008. Memorandum to the Record, NMFS Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle WA 98115. <http://www.afsc.noaa.gov/nmml/pdf/SSLNon-Pups2008memo.pdf>.
- Haynes, T. L., and C. Mishler. 1991. The subsistence harvest and use of Steller sea lions in Alaska. Alaska Dep. Fish and Game Technical Paper No. 198, 44 pp.
- Hennen, D. R. 2004. The Steller sea lion (*Eumetopias jubatus*) decline and the Gulf of Alaska/Bering Sea commercial fishery. Unpubl. Ph.D. dissertation, Montana State University, Bozeman, MT. 224 pp.

- Hoffman, J. I., C. W. Matson, W. Amos, T. R. Loughlin, and J. W. Bickham. 2006. Deep genetic subdivision within a continuously distributed and highly vagile marine mammal, the Steller's sea lion (*Eumetopias jubatus*). *Mol. Ecol.* 15:2821-2832.
- Holmes, E. E., L. W. Fritz, A. E. York, K. Sweeney. 2007. Age-structured modeling provides evidence for a 28-year decline in the birth rate of western Steller sea lions. *Ecolog. Applic.* 17(8):2214-2232.
- Holmes, E. E., and A. E. York. 2003. Using age structure to detect impacts on threatened populations case study using Steller sea lions. *Conserv. Biol.* 17:1794-1806.
- Jones, D. J. 2009. 2008 subsistence harvest of Steller sea lion on St. Paul Island. Memorandum for the Record, April 27, 2009, Aleut Community of St. Paul, Tribal Government, Ecosystem Conservation Office. St. Paul Island, Pribilof Islands, Alaska.
- Kucey, L. 2005. Human disturbance and the hauling out behaviour of Steller sea lions (*Eumetopias jubatus*). M.Sc. thesis, University of British Columbia, Vancouver. 67 pp.
- Kucey, L., and A.W. Trites. 2006. A review of the potential effects of disturbance on sea lions: assessing response and recovery. In A.W. Trites, S. Atkinson, D.P. DeMaster, L.W. Fritz, T.S. Gelatt, L.D. Rea, and K. Wynne (eds.) *Sea Lions of the World*, Alaska Sea Grant Program AK-SG-06-01.
- Lee, J. S., S. Tanabe, H. Umino, R. Tatsukawa, T. R. Loughlin and D. C. Calkins. 1996. Persistent organochlorines in Steller sea lion (*Eumetopias jubatus*) from the bulk of Alaska and the Bering Sea, 1976-1981. *Mar. Pollut. Bull.* 32(7):535-544.
- Lestenkof, A. D., and P. A. Zavadil. 2006. 2005 subsistence harvest of Steller sea lion on St. Paul Island. Memorandum for the Record, August 31, 2006, Aleut Community of St. Paul, Tribal Government, Ecosystem Conservation Office. St. Paul Island, Pribilof Islands, Alaska.
- Lestenkof, A. D., P. A. Zavadil, and D. J. Jones. 2007. 2006 subsistence harvest of Steller sea lion on St. Paul Island. Memorandum for the Record, April 11, 2007, Aleut Community of St. Paul, Tribal Government, Ecosystem Conservation Office. St. Paul Island, Pribilof Islands, Alaska.
- Lestenkof, A. D., P. A. Zavadil, and D. J. Jones. 2008. 2007 subsistence harvest of Steller sea lion on St. Paul Island. Memorandum for the Record, March 4, 2008, Aleut Community of St. Paul, Tribal Government, Ecosystem Conservation Office. St. Paul Island, Pribilof Islands, Alaska.
- Loughlin, T. R. 1997. Using the phylogeographic method to identify Steller sea lion stocks. Pp. 329-341 In A. Dizon, S. J. Chivers, and W. Perrin (eds.), *Molecular genetics of marine mammals, incorporating the proceedings of a workshop on the analysis of genetic data to address problems of stock identity as related to management of marine mammals*. Soc. Mar. Mammal., Spec. Rep. No. 3.
- Loughlin, T. R., D. J. Rugh, and C. H. Fiscus. 1984. Northern sea lion distribution and abundance: 1956-1980. *J. Wildl. Manage.* 48:729-740.
- Loughlin, T.R., and A.E. York. 2000. An accounting of the sources of Steller sea lion mortality. *Mar. Fish. Rev.* 62(4):40-45.
- Manly, B. F. J. 2006. Incidental catch and interactions of marine mammals and birds in the Cook Inlet salmon driftnet and setnet fisheries, 1999-2000. Draft report to NMFS Alaska Region. 83 pp.
- Manly, B. F. J., A. S. Van Atten, K. J. Kuletz, and C. Nations. 2003. Incidental catch of marine mammals and birds in the Kodiak Island set gillnet fishery in 2002. Final report to NMFS Alaska Region. 91 pp.
- Merrick, R. L., T. R. Loughlin, and D. G. Calkins. 1987. Decline in abundance of the northern sea lion, *Eumetopias jubatus*, in 1956-86. *Fish. Bull., U.S.* 85:351-365.
- National Marine Fisheries Service. 1995. Status review of the United States Steller sea lion (*Eumetopias jubatus*) population. Prepared by the National Marine Mammal Laboratory, AFSC, NMFS, NOAA, 7600 Sand Point Way NE, Seattle, WA 98115. 61 pp.
- National Marine Fisheries Service. 2001. Endangered Species Act, Section 7 Consultation Biological Opinion and Incidental Take Statement on the authorization of the Bering Sea/Aleutian Islands and Gulf of Alaska Groundfish Fishery Management Plan Amendments 61 and 70. NMFS Alaska Region, Protected Resources Division, Juneau, AK.
- National Marine Fisheries Service. 2008. Recovery Plan for the Steller sea lion (*Eumetopias jubatus*). Revision. National Marine Fisheries Service, Silver Spring, MD. 325 pp.
- National Marine Fisheries Service. 2010. Endangered Species Act, Section 7 Consultation Biological Opinion and Incidental Take Statement on the authorization of the Bering Sea/Aleutian Islands and Gulf of Alaska Groundfish Fishery Management Plan Amendments 61 and 70. NMFS Alaska Region, Protected Resources Division, Juneau, AK.  
[http://www.fakr.noaa.gov/protectedresources/stellers/esa/biop/final/biop1210\\_chapters.pdf](http://www.fakr.noaa.gov/protectedresources/stellers/esa/biop/final/biop1210_chapters.pdf)

- Nikulin, V. S., and V. N. Burkanov. 2000. Species composition of marine mammal by-catch during Japanese driftnet salmon fishery in southwestern Bering Sea. Unpubl. manuscript, 2 pp. Available, National Marine Mammal Laboratory, AFSC, 7600 Sand Point Way, NE, Seattle, WA 98115.
- Perez, M. A. 2006. Analysis of marine mammal bycatch data from the trawl, longline, and pot groundfish fisheries of Alaska, 1998-2004, defined by geographic area, gear type, and target groundfish catch species. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-167.
- Phillips, C.D., T.S. Gelatt, J.C. Patton, and J.W. Bickham. 2011. Phylogeography of Steller sea lions: relationships among climate change, effective population size, and genetic diversity. *J. Mammal.* 92(5):1091–1104.
- Pitcher, K. W., D. G. Calkins, and G. W. Pendleton. 1998. Reproductive performance of female Steller sea lions: an energetics-based reproductive strategy? *Can. J. Zool.* 76:2075-2083.
- Sease, J. L., W. P. Taylor, T. R. Loughlin, and K. W. Pitcher. 2001. Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1999 and 2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-122, 52 pp.
- Sease, J. L., and A. E. York. 2003. Seasonal distribution of Steller's sea lions at rookeries and haul-out sites in Alaska. *Mar. Mamm. Sci.* 19(4): 745-763.
- Springer, A. M., J. A. Estes, G. B. van Vliet, T. M. Williams, D. F. Doak, E. M. Danner, K.A. Forney and B. Pfister. 2003. Sequential megafaunal collapse in the North Pacific Ocean: an ongoing legacy of industrial whaling? *Proc. Natl. Acad. Sci.* 100: 12223-12228.
- Trites, A. W., and C. P. Donnelly. 2003. The decline of Steller sea lions in Alaska: a review of the nutritional stress hypothesis. *Mamm. Rev.* 33: 3-28.
- Trites, A. W., V. B. Deecke, E. J. Gregr, J. K. B. Ford, and P. F. Olesiuk. 2007. Killer whales, whaling and sequential megafaunal collapse in the North Pacific: a comparative analysis of the dynamics of marine mammals in Alaska and British Columbia following commercial whaling. *Mar. Mamm. Sci.* 23(4):751-765.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Williams, T. M., J. A. Estes, D. F. Doak, and A. M. Springer. 2004. Killer appetites: assessing the role of predators in ecological communities. *Ecology* 85(12):3373-3384.
- Wolfe, R. J., J. A. Fall, and M. Riedel. 2008. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2006. Alaska Dep. Fish and Game, Juneau, AK, Subsistence Div. Tech. Paper No. 339. Juneau, AK.
- Wolfe, R. J., J. A. Fall, and M. Riedel. 2009a. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2007. Alaska Dep. Fish and Game, Juneau, AK, Subsistence Div. Tech. Paper No. 345. Juneau, AK.
- Wolfe, R. J., J. A. Fall, and M. Riedel. 2009b. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2008. Alaska Dep. Fish and Game, Juneau, AK, Subsistence Div. Tech. Paper No. 347. Juneau, AK.
- Wolfe, R. J., J. A. Fall, and R. T. Stanek. 2005. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2004. Alaska Dep. Fish and Game, Juneau, AK, Subsistence Div. Tech. Paper No. 303. Juneau, AK.
- Wolfe, R. J., J. A. Fall, and R. T. Stanek. 2006. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2005. Alaska Dep. Fish and Game, Juneau, AK, Subsistence Div. Tech. Paper No. 319. Juneau, AK.
- Wynne, K. M., D. Hicks, and N. Munro. 1991. 1990 salmon gillnet fisheries observer programs in Prince William Sound and South Unimak Alaska. Annual Rept. NMFS/NOAA Contract 50ABNF000036. 65 pp. NMFS, Alaska Region, Office of Marine Mammals, P.O. Box 21668, Juneau, AK 99802.
- Wynne, K. M., D. Hicks, and N. Munro. 1992. 1991 Marine mammal observer program for the salmon driftnet fishery of Prince William Sound Alaska. Annual Rept. NMFS/NOAA Contract 50ABNF000036. 53 pp. NMFS, Alaska Region, Office of Marine Mammals, P.O. Box 21668, Juneau, AK 99802.
- York, A. E., R. L. Merrick, and T. R. Loughlin. 1996. An analysis of the Steller sea lion metapopulation in Alaska. Chapter 12, Pp. 259-292 *In* D. R. McCullough (ed.), *Metapopulations and wildlife conservation*. Island Press, Covelo, California.

Zavadil, P. A. 2010. 2009 subsistence harvest of Steller sea lion on St. Paul Island. Memorandum for the Record, April 2010, Aleut Community of St. Paul, Tribal Government, Ecosystem Conservation Office. St. Paul Island, Pribilof Islands, Alaska.