

BOTTLENOSE DOLPHIN (*Tursiops truncatus*): Gulf of Mexico Bay, Sound, and Estuarine Stocks

STOCK DEFINITION AND GEOGRAPHIC RANGE

Bottlenose dolphins are distributed throughout the bays, sounds, and estuaries of the Gulf of Mexico (Mullin 1988). The identification of biologically-meaningful “stocks” of bottlenose dolphins in these waters is complicated by the high degree of behavioral variability exhibited by this species (Shane *et al.* 1986; Wells and Scott 1999), and by the lack of requisite information for much of the region.

Previous stock assessment reports have provisionally identified distinct stocks in each of 33 areas of contiguous, enclosed, or semi-enclosed bodies of water adjacent to the Gulf of Mexico (Table 1, Waring *et al.* 1997), based on descriptions of relatively discrete dolphin “communities” in some of these areas. A “community” includes resident dolphins that regularly share large portions of their ranges, exhibit similar distinct genetic profiles, and interact with each other to a much greater extent than with dolphins in adjacent waters. The term, as adapted from Wells *et al.* (1987), emphasizes geographic, genetic, and social relationships of dolphins. Bottlenose dolphin communities do not constitute closed demographic populations, as individuals from adjacent communities are known to interbreed. Nevertheless, the geographic nature of these areas and long-term stability of residency patterns suggest that many of these communities exist as functioning units of their ecosystems and, under the Marine Mammal Protection Act, must be maintained as such. Also, the stable patterns of residency observed within communities suggest that long periods would be required to repopulate the home range of a community were it eradicated or severely depleted. Thus, in the absence of information supporting management on a larger scale, it is appropriate to adopt a risk-averse approach and focus management efforts at the level of the community rather than at some larger demographic scale. Support for this risk-averse approach derives from several sources.

Long-term (year-round, multi-year) residency by at least some individuals has been reported from nearly every site where photographic identification or tagging studies have been conducted in the Gulf of Mexico. In Texas, some of the dolphins in the Matagorda-Espiritu Santo Bay area (Gruber 1981; Lynn 1995; Würsig and Lynn 1996), Aransas Pass (Shane 1977; Weller 1998), San Luis Pass (Maze 1997), and Galveston Bay (Bräger 1993; Bräger *et al.* 1994; Fertl 1994) have been reported as long-term residents. Hubard (1998) reported sightings of dolphins tagged 12-15 years previously in Mississippi Sound. In Florida, long-term residency has been reported from Choctawhatchee Bay (1989-1993, F. Townsend, unpublished data), Tampa Bay (Wells 1986; Wells *et al.* 1996a), Sarasota Bay (Irvine and Wells 1972; Irvine *et al.* 1981; Wells 1986, 1991; Scott *et al.* 1990; Wells *et al.* 1987), Lemon Bay (Wells *et al.* 1996b), and Charlotte Harbor/Pine Island Sound (Shane 1990; Wells *et al.* 1996b, 1997). In many cases, residents emphasize use of the bay, sound, or estuary waters, with limited movements through passes to the Gulf of Mexico (Shane 1977, 1990; Gruber 1981; Irvine *et al.* 1981; Lynn 1995, Maze 1997). These habitat use patterns are reflected in the ecology of the dolphins in some areas; for example, residents of Sarasota Bay, Florida lacked squid in their diet, unlike non-resident dolphins stranded on nearby Gulf beaches (Barros and Wells 1998).

Genetic data also support the concept of relatively discrete bay, sound, and estuary stocks. Analyses of mitochondrial DNA haplotype distributions indicate the existence of clinal variations along the Gulf of Mexico coastline (Duffield and Wells, in press).

Differences in reproductive seasonality from site to site also suggest genetic-based distinctions between communities (Urian *et al.* 1996). Mitochondrial DNA analyses suggest finer-scale structural levels as well. For example, Matagorda Bay, Texas dolphins appear to be a localized population (NMFS unpublished data), and differences in haplotype frequencies distinguish between adjacent communities in Tampa Bay, Sarasota Bay, and Charlotte Harbor/Pine Island Sound, along the central west coast of Florida (Duffield and Wells 1991, in press). Examination of protein electrophoretic data resulted in similar conclusions for the Florida dolphins (Duffield and Wells 1986).

The long-term structure and stability of at least some of these communities is exemplified by the residents of Sarasota Bay, Florida. This community has been observed since 1970 (Irvine and Wells 1972; Scott *et al.* 1990; Wells 1991). The number of dolphins regularly occupying the Sarasota Bay area has remained consistently at about 100. At least four generations of identifiable residents currently inhabit the region, including half of those first identified in 1970. Maximum immigration and emigration rates of about 2-3% have been estimated (Wells and Scott 1990).

Genetic exchange occurs between resident communities; hence the application of the demographically and behaviorally-based term “community” rather than “population” (Wells 1986). Up to about 30% of calves in Sarasota Bay apparently have been sired by non-residents (Duffield and Wells in press). A variety of potential exchange mechanisms occur in the Gulf. Small numbers of inshore dolphins traveling between regions have been reported, with patterns ranging from traveling through adjacent communities (Wells

1986; Wells *et al.* 1996a,b) to movements over distances of several hundred km in Texas waters (Gruber 1981; Würsig and Lynn 1996; Würsig unpublished data). In many areas year-round residents co-occur with non-resident dolphins, providing potential opportunities for genetic exchange. About 17% of group sightings involving resident Sarasota Bay dolphins include at least one non-resident as well (Wells *et al.* 1987). Similar mixing of inshore residents and non-residents is seen off San Luis Pass, Texas (Maze 1997). Non-residents exhibit a variety of patterns, ranging from apparent nomadism recorded as transience in a given area, to apparent seasonal or non-seasonal migrations. Passes, especially the mouths of the larger estuaries, serve as mixing areas. For example, several communities mix at the mouth of Tampa Bay, Florida (Wells 1986), and most of the dolphins identified in the mouths of Galveston Bay and Aransas Pass, Texas were considered transients (Henningesen 1991; Bräger 1993; Weller 1998).

Seasonal movements of dolphins into and out of some of the bays, sounds, and estuaries provide additional opportunities for genetic exchange with residents, and complicate the identification of stocks in coastal and inshore waters. In small bay systems such as Sarasota Bay, Florida and San Luis Pass, Texas residents move into Gulf coastal waters in fall/winter, and return inshore in spring/summer (Irvine *et al.* 1981; Maze 1997). In larger bay systems, seasonal changes in abundance suggest possible migrations, with increases in more northerly bay systems in summer, and in more southerly systems in winter. Fall/winter increases in abundance have been noted for Matagorda Bay (Gruber 1981; Lynn 1995; Würsig and Lynn 1996), Aransas Pass (Shane 1977; Weller 1998), Tampa Bay (Scott *et al.* 1989), and Charlotte Harbor/Pine Island Sound (Thompson 1981; Scott *et al.* 1989). Spring/summer increases in abundance have been reported for Galveston Bay (Henningesen 1991; Bräger 1993; Fertl 1994) and Mississippi Sound (Hubard 1998).

Much uncertainty remains regarding the structure of bottlenose dolphin stocks in many of the Gulf of Mexico bays, sounds, and estuaries. Given the apparent co-occurrence of resident and non-resident dolphins in these areas, and the demonstrated variations in abundance, it appears that consideration should be given to the existence of a complex of stocks, and to the roles of bays, sounds, and estuaries for stocks emphasizing Gulf of Mexico coastal waters. A starting point for management strategy should be the protection of the long-term resident communities, with their multi-generational geographic, genetic, demographic, and social stability. These localized units would be at greatest risk from geographically-localized impacts. Complete characterization of many of these basic units would benefit from additional photo-identification, telemetry, and genetic research (Wells 1994).

The current provisional stocks follow the designations in Table 1, with a few revisions. Available information suggests that Block B35, Little Sarasota Bay, can be subsumed under Sarasota Bay, and B36, Caloosahatchee River, can be considered a part of Pine Island Sound. As more information becomes available, additional combination or division may be warranted. For example, a number of geographically and socially distinct subgroupings of dolphins in regions such as Tampa Bay, Charlotte Harbor, Pine Island Sound, Aransas Pass, and Matagorda Bay have been identified, but the importance of these distinctions to stock designations remain undetermined (Shane 1977; Gruber 1981; Wells *et al.* 1996a,b, 1997; Würsig and Lynn 1996).

Understanding the full complement of the stock complex using the bay, sound, and estuarine waters of the Gulf of Mexico will require much additional information. The development of biologically-based criteria to better define and manage stocks in this region should integrate multiple approaches, including studies of ranging patterns, genetics, morphology, social patterns, distribution, life history, stomach contents, isozyme analyses, and contaminant concentrations. Spatially-explicit population modeling could aid in evaluating the implications of community-based stock definition. As these studies provide new information on what constitutes a bottlenose dolphin "biological stock,"

current provisional definitions will likely need to be revised. As stocks are more clearly identified, it will be possible to conduct abundance estimates using standardized methodology across sites (thereby avoiding some of the previous problems of mixing results of aerial and boat-based surveys), identify fisheries and other human impacts relative to specific stocks, and perform individual stock assessments. As recommended by the Atlantic Scientific Review Group (November 1998, Portland, Maine), a workshop will be held during 1999 to review current information pertaining to bottlenose dolphin stock structure in Gulf of Mexico bays, sounds, and estuaries, conduct simulations of alternative stock structure and, if warranted, propose a new stock structure.

Table 1. Bottlenose dolphin abundance (N_{BEST}), coefficient of variation (CV), minimum population estimate (N_{MIN}), and Potential Biological Removal (PBR) in U.S. Gulf of Mexico bays, sounds, and other estuaries. Blocks refer to aerial survey blocks illustrated in Fig. 1. Blocks with an abundance of zero were surveyed but not considered stocks at this time (but see Note 1 below).

Blocks	Gulf of Mexico Estuary	N_{BEST}	CV	N_{MIN}	PBR	Year	Reference
B51	Laguna Madre	80	1.57	31	0.3	1992	A
B52	Nueces Bay, Corpus Christi Bay	58	0.61	36	0.4	1992	A
B50	Compano Bay, Aransas Bay, San Antonio Bay, Redfish Bay, Espiritu Santo Bay	55	0.82	30	0.3	1992	A
B54	Matagorda Bay, Tres Palacios Bay, Lavaca Bay	61	0.45	42	0.4	1992	A
B55	West Bay	29	1.10	14	0.1	1992	A
B56	Galveston Bay, East Bay, Trinity Bay	152	0.43	107	1.1	1992	A
B57	Sabine Lake	0 ¹	-			1992	A
B58	Calcasieu Lake	0 ¹	-			1992	A
B59	Vermillion Bay, West Cote Blanche Bay, Atchafalaya Bay	0 ¹	-			1992	A
B60	TerreBonne Bay, Timbalier Bay	100	0.53	66	0.7	1993	A
B61	Barataria Bay	219	0.55	142	1.4	1993	A
B30	Mississippi River Delta	0 ¹	-			1993	A
B02-05, 29,31	Bay Boudreau, Mississippi Sound	1,401	0.13	1,256	13	1993	A
B06	Mobile Bay, Bonsecour Bay	122	0.34	92	0.9	1993	A
B07	Perdido Bay	0 ¹	-			1993	A
B08	Pensacola Bay, East Bay	33	0.80	18	0.2	1993	A
B09	Choctawhatchee Bay	242	0.31	188	1.9	1993	A
B10	St. Andrew Bay	124	0.57	79	0.8	1993	A
B11	St. Joseph Bay	0 ¹	-			1993	A
B12-13	St. Vincent Sound, Apalachicola Bay, St. Georges Sound	387	0.34	293	2.9	1993	A
B14-15	Apalachee Bay	491	0.39	358	3.6	1993	A
B16	Waccasassa Bay, Withlacoochee Bay, Crystal Bay	100	0.85	54	0.5	1994	A
B17	St. John's Sound, Clearwater Harbor	37	1.06	18	0.2	1994	A
B32-34	Tampa Bay	559	0.24	458	4.6	1994	A
B20	Sarasota Bay	97	na ³	97	1.0	1992	B
B35	Little Sarasota Bay	2 ²	0.24	2	0.0	1985	C
B21	Lemon Bay	0 ¹	-			1994	A
B22-23	Pine Sound, Charlotte Harbor, Gasparilla Sound	209	0.38	153	1.5	1994	A
B36	Caloosahatchee River	0 ^{1,2}	-			1985	C
B24	Estero Bay	104	0.67	62	0.6	1994	A
B25	Chokoloskee Bay, Ten Thousand Islands, Gullivan Bay	208	0.46	144	1.4	1994	A
B27	Whitewater Bay	242	0.37	179	1.8	1994	A
B28	Florida Keys (Bahia Honda to Key West)	29	1.00	14	0.1	1994	A

References: A- Blaylock and Hoggard 1994; B- Wells 1992; C- Scott et al. 1989

Notes:

¹ During earlier surveys (Scott *et al.* 1989), the range of seasonal abundances was as follows: B57, 0-2 (CV=0.38); B58, 0-6 (0.34); B59, 0-0; B30, 0-182(0.14); B07, 0-0; B21, 0-15(0.43); and B36, 0-0.

² Block not surveyed during surveys reported in Blaylock and Hoggard 1994.

³ No CV because N_{BEST} was a direct count of known individuals.




Figure 1. U.S. Gulf of Mexico bays and sounds. Each of the alpha-numerically designated blocks corresponds to one of the NMFS Southeast Fisheries Science Center logistical aerial survey areas listed in Table 1. The bottlenose dolphins inhabiting each bay and sound are considered to comprise a unique stock for purposes of this assessment.

POPULATION SIZE

Population size (Table 1) for all of the stocks except Sarasota Bay, Florida, was estimated from preliminary analyses of line-transect data collected during aerial surveys conducted in September–October 1992 in Texas and Louisiana; in September–October 1993 in Louisiana, Mississippi, Alabama, and the Florida panhandle (Blaylock and Hoggard 1994); and in September–November 1994 along the west coast of Florida (NMFS unpublished data). Standard line-transect perpendicular sighting distance analytical methods (Buckland *et al.* 1993) and the computer program DISTANCE (Laake *et al.* 1993) were used. Stock size in Sarasota Bay, Florida, was obtained through direct count of known individuals (Wells 1992).

Minimum Population Estimate

The minimum population estimate (Table 1) is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The minimum population estimate was calculated for each block from the estimated population size and its associated coefficient of variation. Where the population size resulted from a direct count of known individuals, the minimum population size was identical to the estimated population size.

Current Population Trend

The data are insufficient to determine population trends for all of the Gulf of Mexico bay, sound, and estuary bottlenose dolphin communities. The Sarasota Bay community, however, has been monitored since 1970 and has remained relatively constant over the last 20+ years at approximately 105 animals (Wells 1998). Three anomalous mortality events have occurred among portions of these dolphin communities between 1990 and 1994; however, it is not possible to accurately partition the mortalities between bay and coastal stocks, thus the impact of these mortality events on communities is not known.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are not known for the dolphin communities that comprise these stocks. While productivity rates may be estimated for individual females within communities, such estimates are confounded at the stock level due to the influx of dolphins from adjacent areas which balance losses, and the unexplained loss of some individuals which offset births and recruitment (Wells 1998). Continued monitoring and expanded survey coverage will be required to address and develop estimates of productivity for these dolphin communities. The maximum net productivity rate was assumed to be 0.04. This value is based

on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (Wade and Angliss 1997). The “recovery” factor, which accounts for endangered, depleted, and threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.5 because these stocks are of unknown status. PBR for each stock is given in Table 1.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

There are a number of difficulties associated with the interpretation of stranding data. It is possible that some or all of the stranded dolphins may have been from a nearby coastal stock; however, the proportion of stranded dolphins belonging to another stock cannot be determined because of the difficulty of determining from where the stranded carcass originated. Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the dolphins which die or are seriously injured in fishery interactions wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction, and the condition of the carcass if badly decomposed can inhibit the interpretation of cause of death.

A total of 1,881 bottlenose dolphins were found stranded in the USA Southeast Gulf of Mexico from 1993 to 1997 (Table 2) (NMFS unpublished data). Of these, 57 or 3% showed evidence of human interactions as the cause of death (*e.g.*, gear entanglement, mutilation, gunshot wounds). Bottlenose dolphins are known to become entangled in recreational and commercial fishing gear (Wells *et al.* 1998; Gorzelany 1998; Wells and Scott 1994) and some are struck by recreational and commercial vessels (Wells and Scott 1997). In 1998 alone, two resident bottlenose dolphins and an associated calf were killed by vessel strikes and a resident young-of-the-year died from entanglement in a crab-pot float line (R.S. Wells personal communication).

The Gulf of Mexico menhaden fishery was observed to take 9 bottlenose dolphins (three fatally) between 1992 and 1995 (NMFS unpublished data). There were 1,366 sets observed out of 26,097 total sets, which if extrapolated for all years suggests that as many as 172 bottlenose dolphins could have been taken in this fishery with up to 57 animals killed. An observer program is urgently needed to obtain statistically reliable information for this fishery on the number of sets annually, the incidental take and mortality rates, and the communities from which bottlenose dolphins are being taken.

Some of the bay, sound and estuarine communities were the focus of a live-capture fishery for bottlenose dolphins which supplied dolphins to the U.S. Navy and to oceanaria for research and public display for almost two decades (NMFS unpublished data). During the period between 1972-89, 490 bottlenose dolphins, an average of 29 dolphins annually, were removed from a few locations in the Gulf of Mexico, including the Florida Keys. Mississippi Sound sustained the highest level of removals with 202 dolphins taken from this stock during this period, representing 41% of the total and an annual average of 12 dolphins (compared to a current PBR of 13). The annual average number of removals never exceeded current PBR levels, but it may be biologically significant that 73% of the dolphins removed during 1982-88 were females. The impact of those removals on the stocks is unknown.

Fishery Information

Annual fishing effort for the shrimp trawl fishery in the USA Gulf of Mexico bays, sounds, and estuaries during 1988-1993 averaged approximately 2.20 million hours of tows (CV = 0.11) (NMFS unpublished data). There have been no reports of incidental mortality or injury in any of these stocks associated with the shrimp trawl fishery.

A fishery for blue crabs operates in estuarine areas throughout the Gulf of Mexico coast employing traps attached to a buoy with rope. Bottlenose dolphins have been reported stranded with polypropylene rope around their flukes (NMFS 1991; McFee and Brooks, Jr. 1998; NMFS unpublished data), indicating the possibility of entanglement with crab pot lines. This fishery has not been monitored by observers and there are no estimates of bottlenose dolphin mortality or serious injury for this fishery.

Gillnets are not used in Texas, and gillnets over 46 m² in area were not allowed in Florida past July 1995, but fixed and runaround gillnets are currently in use in Louisiana, Mississippi, and Alabama. These fisheries, for the most part, operate year around. They are state-controlled and licensed, and vary widely in intensity and target species. No marine mammal mortalities associated with gillnet fisheries have been reported in these states, but stranding data suggest that gillnet and marine mammal interaction does occur, causing mortality and serious injury.

Table 2. Bottlenose dolphin strandings in the USA Gulf of Mexico (West Florida to Texas) from 1993 to 1997. Data are from the Southeast Marine Mammal Stranding Database (SESUS).

State		1993	1994	1995	1996	1997	Total
Florida	No. Stranded	134	51	101	133	63	482
	No. Human Interactions	4	2	3	2	0	11
	% With Human Interactions	3%	4%	3%	2%	0%	2%
Alabama	No. Stranded	48	16	15	17	14	110
	No. Human Interactions	1	0	1	0	1	3
	% With Human Interactions	2%	0%	7%	0%	7%	3%
Mississippi	No. Stranded	64	25	32	59	42	222
	No. Human Interactions	4	0	4	2	2	12
	% With Human Interactions	6%	0%	12%	3%	5%	5%
Louisiana	No. Stranded	14	74	31	92	42	253
	No. Human Interactions	0	0	1	3	1	5
	% With Human Interactions	0%	0%	3%	3%	2%	2%
Texas	No. Stranded	133	227	110	208	136	814
	No. Human Interactions	4	6	7	7	2	26
	% With Human Interactions	0%	3%	6%	3%	0%	3%
Totals	No. Stranded	393	393	289	509	297	1881
	No. Human Interactions	13	8	16	14	6	57
	% With Human Interactions	3%	2%	6%	3%	2%	3%

Other Mortality

The near shore habitat occupied by many of these stocks is adjacent to areas of high human population, and in some bays, such as Mobile Bay in Alabama and Galveston Bay in Texas, is highly industrialized. The area surrounding Galveston Bay, for example, has a coastal population of over 3 million people. More than 50% of all chemical products manufactured in the USA are produced there and 17% of the oil produced in the Gulf of Mexico is refined there (Henningesen and Würsig 1991). Many of the enclosed bays in Texas are surrounded by agricultural lands which receive periodic pesticide applications.

Concentrations of chlorinated hydrocarbons and metals were examined in conjunction with an anomalous mortality event of bottlenose dolphins in Texas bays in 1990 and found to be relatively low in most; however, some had concentrations at levels of possible toxicological concern (Varanasi *et al.* 1992). No studies to date have determined the amount, if any, of indirect human-induced mortality resulting from pollution or habitat degradation. However, a recent health assessment of 35 bottlenose dolphins from Matagorda Bay, Texas associated high levels of chlorinated hydrocarbons with low health assessment scores (Reif *et al.*, in review). Morbillivirus has also been implicated in the deaths of bottlenose dolphins in some of these communities (Duignan *et al.* 1996).

STATUS OF STOCK

The status of these stocks relative to OSP is unknown and this species is not listed as threatened or endangered under the Endangered Species Act. The occurrence of three anomalous mortality events among bottlenose dolphins along the USA Gulf of Mexico coast since 1990 (NMFS unpublished data) is cause for concern; however, the effects of the mortality events on stock abundance have not yet been determined. The available evidence suggests that bottlenose dolphin stocks in the northern and western coastal portion of the USA Gulf of Mexico may have experienced a morbillivirus epidemic in 1993 (Lipscomb 1993; Lipscomb *et al.* 1994). Seven of 35 live-captured bottlenose dolphins (20%) from Matagorda Bay, Texas, in 1992, tested positive for previous exposure to cetacean morbillivirus (Reif *et al.*, in review), and it is possible that other estuarine resident stocks have been exposed to the morbillivirus (Duignan *et al.* 1996).

The relatively high number of bottlenose dolphin deaths which occurred during the mortality events in the last decade suggests that some of these stocks may be stressed. Fishery-related mortality and serious injury for each of these stocks is not known, but considering the evidence from stranding data, the total fishery-related mortality and serious injury exceeds 10% of the total PBR, and, therefore, it is not insignificant and approaching the zero mortality and serious injury rate. For these reasons, and because the PBR for most of these stocks would be exceeded with the incidental capture of a single dolphin, each of these stocks is a strategic stock.

REFERENCES

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background and a Summary of the 1995 Assessments. NOAA Technical Memorandum. NMFS-OPR-6, 73 pp.
- Barros, N.B. and R.S. Wells. 1998. Prey and feeding patterns of resident bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *J. Mamm.* 79(3):1045-1059.
- Blaylock, R.A. and W. Hoggard. 1994. Preliminary estimates of bottlenose dolphin abundance in southern U.S. Atlantic and Gulf of Mexico continental shelf waters. NOAA Technical Memorandum. NMFS-SEFSC-356, 10 pp.
- Bräger, S. 1993. Diurnal and seasonal behavior patterns of bottlenose dolphins (*Tursiops truncatus*). *Mar. Mamm. Sci.* 9: 434-440.
- Bräger, S., B. Würsig, A. Acevedo, and T. Henningsen. 1994. Association patterns of bottlenose dolphins (*Tursiops truncatus*) in Galveston Bay, Texas. *J. Mamm.* 75(2): 431-437.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, and J.L. Laake. 1993. Distance sampling: Estimating abundance of biological populations. *Chapman & Hall*, London. 446 pp.
- Duffield, D.A. and R.S. Wells. 1986. Population structure of bottlenose dolphins: Genetic studies of bottlenose dolphins along the central west coast of Florida. Contract Report to National Marine Fisheries Service, Southeast Fisheries Center, Contract No. 45-WCNF-5-00366, 16 pp.
- Duffield, D.A. and R.S. Wells. 1991. The combined application of chromosome, protein and molecular data for the investigation of social unit structure and dynamics in *Tursiops truncatus*. Pages 155-169 in A.R. Hoelzel (ed), Genetic Ecology of Whales and Dolphins. *Rep. int. Whal. Commn., Special Issue 13, Cambridge, U.K.*
- Duffield, D.A. and R.S. Wells. In press. The molecular profile of a resident community of bottlenose dolphins, *Tursiops truncatus*. In C.J. Pfeiffer (ed), *Cell and Molecular Biology of Marine Mammals*.
- Duignan, P.J., C. House, D.K. Odell, R.S. Wells, L. Hansen, M.T. Walsh, D.J. St. Aubin, B.K. Rima and J.R. Geraci. 1996. Morbillivirus infection in bottlenose dolphin: evidence for recurrent epizootics in the Western Atlantic and Gulf of Mexico. *Mar. Mamm. Sci.* 12(4):499-515.
- Fertl, D.C. 1994. Occurrence, movements, and behavior of bottlenose dolphins (*Tursiops truncatus*) in association with the shrimp fishery in Galveston Bay, Texas. M. Sc. thesis, Texas A&M University, College Station. 117 pp.
- Gruber, J.A. 1981. Ecology of the Atlantic bottlenosed dolphin (*Tursiops truncatus*) in the Pass Cavallo area of Matagorda Bay, Texas. M. Sc. thesis, Texas A&M University, College Station. 182 pp.
- Henningsen, T. 1991. Zur Verbreitung und Ökologie des Großen Tümmlers (*Tursiops truncatus*) in Galveston, Texas. Diploma thesis, Christian-Albrechts-Universität, Kiel, Germany. 80 pp.
- Henningsen, T. and B. Würsig. 1991. Bottle-nosed dolphins in Galveston Bay, Texas: Numbers and activities. Pages 36-38 in P. G. H. Evans (ed), European research on cetaceans - 5. Proceedings of the Fifth Annual Conference of the European Cetacean Society, Sandefjord, Norway, 21-23 February, 1991. Cambridge, UK.
- Hubard, C.W. 1998. Abundance, distribution, and site fidelity of bottlenose dolphins in Mississippi Sound, Mississippi. M. Sc. thesis, University of Alabama, Tuscaloosa. 101 pp.
- Irvine, B. and R.S. Wells. 1972. Results of attempts to tag Atlantic bottlenose dolphins (*Tursiops truncatus*). *Cetology* 13:1-5.

- Irvine, A.B., M.D. Scott, R.S. Wells and J.H. Kaufmann. 1981. Movements and activities of the Atlantic bottlenose dolphin, *Tursiops truncatus*, near Sarasota, Florida. *Fish. Bull. U.S.* 79:671-688.
- Laake, J. L., S.T. Buckland, D.R. Anderson, and K.P. Burnham. 1993. DISTANCE user's guide, V2.0. Colorado Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins. 72 pp.
- Lipscomb, T.P. 1993. Some answers to questions about morbillivirus. Pages 4-5 in R.A. Blaylock, B. Mase, and D. K. Odell (eds), *Strandings*, Vol. 2, No. 3, SEFSC Miami Laboratory, Miami, Florida, 7 pp.
- Lipscomb, T.P. 1994. Morbilliviral disease in an Atlantic bottlenose dolphin (*Tursiops truncatus*) from the Gulf of Mexico. *Journal of Wildlife Diseases* 30(4): 572-576.
- Lynn, S.K. 1995. Movements, site fidelity, and surfacing patterns of bottlenose dolphins on the central Texas coast. M. Sc. thesis, Texas A&M University, College Station. 92 pp.
- Maze, K.S. 1997. Bottlenose dolphins of San Luis Pass, Texas: Occurrence patterns, site fidelity, and habitat use. M. Sc. thesis, Texas A&M University, College Station. 79 pp.
- McFee, W.E. and W. Brooks, Jr. 1998. Fact finding meeting of marine mammal entanglement in the crab pot fishery: A summary. U.S. Fish and Wildlife Service unpublished report.
- Mullin, K.D. 1988. Comparative seasonal abundance and ecology of bottlenose dolphins (*Tursiops truncatus*) in three habitats of the north-central Gulf of Mexico. Ph. D. dissertation, Mississippi State University, Starkville. 135 pp.
- NMFS. 1991. Proposed regime to govern the interactions between marine mammals and commercial fishing operations after October 1, 1993. Draft Environmental Impact Statement, June 1991.
- Reif, J.S., L.J. Hansen, S. Galloway, G. Mitchum and T. L. Schmitt. In review. The relationship between chlorinated hydrocarbon contaminants and selected health parameters in bottlenose dolphins (*Tursiops truncatus*) from Matagorda Bay, Texas, 1992. Colorado State University, Fort Collins, and NMFS, Southeast Fisheries Science Center, Miami, Florida.
- Scott, G.P., D.M. Burn, L.J. Hansen and R.E. Owen. 1989. Estimates of bottlenose dolphin abundance in the Gulf of Mexico from regional aerial surveys. *CRD88/89-07*.
- Scott, G.P. 1990. Management-oriented research on bottlenose dolphins by the Southeast Fisheries Center. Pages 623-639 in S. Leatherwood and R. R. Reeves (eds), *The bottlenose dolphin*. Academic Press, San Diego. 653 pp.
- Scott, M.D., R.S. Wells, and A.B. Irvine. 1990. A long-term study of bottlenose dolphins on the west coast of Florida. Pages 235-244 in S. Leatherwood and R. R. Reeves (eds), *The bottlenose dolphin*. Academic Press, San Diego. 653 pp.
- Shane, S.H. 1977. The population biology of the Atlantic bottlenose dolphin, *Tursiops truncatus*, in the Aransas Pass area of Texas. M. Sc. thesis, Texas A&M University, College Station. 238 pp.
- Shane, S.H. 1990. Behavior and ecology of the bottlenose dolphin at Sanibel Island, Florida. Pages 245-265 in S. Leatherwood and R.R. Reeves (eds), *The bottlenose dolphin*. Academic Press, San Diego. 653 pp.
- Shane, S.H., R.S. Wells and B. Würsig. 1986. Ecology, behavior, and social organization of the bottlenose dolphin: A review. *Mar. Mamm. Sci.* 2(1):34-63.
- Thompson, N.B. 1981. Estimates of abundance of *Tursiops truncatus* in Charlotte Harbor, Florida. NOAA/NMFS/SEFSC/Miami Laboratory, Fishery Data Analysis Technical Report.
- Urian, K.W., D.A. Duffield, A.J. Read, R.S. Wells and D.D. Shell. 1996. Seasonality of reproduction in bottlenose dolphins, *Tursiops truncatus*. *J. Mamm.* 77:394-403.
- Varanasi, U., K.L. Tilbury, D.W. Brown, M.M. Krahn, C.A. Wigren, R.C. Clark, and S.L. Chan. 1992. Pages 56-86 in L.J. Hansen (ed), *Report on Investigation of 1990 Gulf of Mexico Bottlenose Dolphin Strandings*, Southeast Fisheries Science Center Contribution MIA-92/93-21, 219 pp.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, Seattle, Washington. NOAA Technical Memorandum. NMFS-OPR-12, 93 pp.
- Waring, G.T., D.L. Palka, K.D. Mullin, J.H.W. Hain, L.J. Hansen and K.D. Bisack. 1997. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments -- 1996. NOAA Technical Memorandum. NMFS-NE-114.
- Weller, D.W. 1998. Global and regional variation in the biology and behavior of bottlenose dolphins. Ph. D. dissertation, Texas A&M University, College Station. 142 pp.
- Wells, R.S. 1986a. Population structure of bottlenose dolphins: behavioral studies along the central west coast of Florida. Contract report to NMFS, SEFSC. Contract No. 45-WCNF-5-00366, 58 pp.
- Wells, R.S. 1986b. Structural aspects of dolphin societies. Ph. D. dissertation, University of California, Santa Cruz. 234 pp.

- Wells, R.S. 1991. The role of long-term study in understanding the social structure of a bottlenose dolphin community. Pages 199-225 in K. Pryor and K.S. Norris (eds), *Dolphin Societies: Discoveries and Puzzles*. University of California Press, Berkeley. 397 pp.
- Wells, R.S. 1992. The marine mammals of Sarasota Bay. Pages 9.1-9.23 in *Sarasota Bay: Framework for action*. Sarasota Bay National Estuary Program, Sarasota, Florida.
- Wells, R.S. 1994. Determination of bottlenose dolphin stock discreteness: Application of a combined behavioral and genetic approach. Pages 16-20 in K.R. Wang, P.M. Payne, and V.G. Thayer (compilers), *Coastal Stock(s) of Atlantic Bottlenose Dolphin: Status Review and Management*. Proceedings and Recommendations from a Workshop held in Beaufort, NC, 13-14 September 1993. NOAA Technical Memorandum. NMFS-OPR-4. 120 pp.
- Wells, R.S. 1998. Progress report: Sarasota long-term bottlenose dolphin research. Unpublished contract report to the U.S. Department of Commerce, NOAA Fisheries, Southeast Fisheries Science Center, Miami. 5 pp.
- Wells, R.S. and M.D. Scott. 1990. Estimating bottlenose dolphin population parameters from individual identification and capture-release techniques. Pages 407-415 in P.S. Hammond, S.A. Mizroch and G.P. Donovan (eds), *Individual Recognition of Cetaceans: Use of Photo-Identification and Other Techniques to Estimate Population Parameters*. *Rep. int. Whal. Commn., Special Issue 12*, Cambridge, U.K. 440 pp.
- Wells, R.S. and M.D. Scott. 1999. Bottlenose dolphins. Pages 137-182 in S.H. Ridgway and R. Harrison (eds), *Handbook of Marine Mammals, Vol. 6, the Second Book of Dolphins and Porpoises*. Academic Press, San Diego.
- Wells, R.S., M.D. Scott and A.B. Irvine. 1987. The social structure of free-ranging bottlenose dolphins. Pages 247-305 in Genoways, H. (ed), *Current Mammalogy, Vol. 1*. New York, Plenum Press.
- Wells, R.S., K.W. Urian, A.J. Read, M.K. Bassos, W.J. Carr and M.D. Scott. 1996a. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Tampa Bay, Florida: 1988-1993. NOAA Technical Memorandum. NMFS-SEFSC-385, 25 pp. + 6 Tables, 8 Figures, and 4 Appendices.
- Wells, R.S., M.K. Bassos, K.W. Urian, W.J. Carr and M.D. Scott. 1996b. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Charlotte Harbor, Florida: 1990-1994. NOAA Technical Memorandum. NMFS-SEFSC-384, 36 pp. + 8 Tables, 10 Figures, and 5 Appendices.
- Wells, R.S., M.K. Bassos, K.W. Urian, S.H. Shane, E.C.G. Owen, C.F. Weiss, W.J. Carr and M.D. Scott. 1997. Low-level monitoring of bottlenose dolphins, *Tursiops truncatus*, in Pine Island Sound, Florida: 1996. Contract report to National Marine Fisheries Service, Southeast Fisheries Center. Contribution No. 40-WCNF601958
- Würsig, B. and S.K. Lynn. 1996. Movements, site fidelity, and respiration patterns of bottlenose dolphins on the central Texas coast. NOAA Technical Memorandum. NMFS-SEFSC-383, 43 pp. + 10 Tables, 15 Figures, and 6 Appendices.