

NATIONAL MARINE FISHERIES SERVICE
APPLICATION INSTRUCTIONS FOR PERMITS FOR THE INCIDENTAL TAKE
OF ENDANGERED OR THREATENED SPECIES UNDER THE ENDANGERED SPECIES ACT

In coordination with, but not substituting for 50 CFR 222.307
OMB control number (0648-0230) Expiration date for clearance: 03/31/2012


Information Required in the Application

The Assistant Administrator may issue permits to take endangered or threatened marine species incidentally to an otherwise lawful activity under section 10(a)(1)(B) of the Endangered Species Act of 1973 (ESA). The information collection associated with the following application instructions is required for the purpose of obtaining such a permit. The information provided will be used to process the incidental take permit in accordance with the ESA, including the solicitation of public comments on the justification of the take of ESA-listed species incidental to proposed activities. The information provided by an applicant in accordance with these instructions is not confidential and is subject to public exposure for comments. Notwithstanding any other provision of the law, no person is required to respond to, nor shall any person be subject to a penalty for failure to comply with, a collection of information subject to the requirements of the Paperwork Reduction Act, unless that collection of information displays a currently valid OMB Control Number. Public reporting burden for this collection of information is estimated to average 80 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to the address below.


An application for a permit should provide all of the following information. The information needed in the application should be presented in the same structure and format shown below to increase processing efficiency. When a question does not apply, do not overlook the category, but indicate Not Applicable (N.A.). In some cases, a brief explanation as to why the category is not applicable may expedite processing. Please note that for the title and closing statement of the application, specific wording is required.

If the applicant represents an individual or a single entity, such as a corporation, the application should be for an individual incidental take permit. If the applicant represents a group or organization whose members conduct the same or a similar activity in the same geographical area with similar impacts on endangered or threatened marine species, the application should be for a general incidental take permit. To be covered by a general incidental take permit, each individual conducting the activity must have a certificate of inclusion issued under paragraph (f) of 50 CFR 222.307. NMFS estimates a public reporting burden of .5 hour for each certificate of inclusion. The sufficiency of applications will be determined by the Assistant Administrator in accordance with the requirements of 50 CFR 222.307.

I. One of the titles below as appropriate:

A.  Application for an Individual Incidental Take Permit under the Endangered Species Act of 1973.

Dan Forster or
Director, Wildlife Resources Division
Georgia Department of Natural Resources
2070 U.S. Hwy 278 SE
Social Circle, GA 30025


Spud Woodward
Director, Coastal Resources Division
Georgia Department of Natural Resources
One Conservation Way
Brunswick, GA 31520

B. Application for a General Incidental Take Permit under the Endangered Species Act of 1973.

II. Date of the application.

February 27, 2012

- III. The name, address, telephone, and fax number of the applicant. If the applicant is a partnership, corporate entity or is representing a group or organization, include applicable details.

Dan Forster (770-918-6400) and/or Spud Woodward (912-264-7218)

See above for respective addresses

- IV. A description of the endangered or threatened species, by common and scientific name, and a description of the status, distribution, seasonal distribution, habitat needs, feeding habits and other biological requirements of the affected species.

Refer to previously provided report "Altamaha Sturgeon-Section 6 Final Report (Bahn and Peterson, 2010)".

- V. A detailed description of the proposed activity, including, but not limited to:

- A. The anticipated dates and duration of the activity.

GA commercial shad season dates can be found on pages 17-18 of "Georgia's Commercial Saltwater Fishing Regulations" that was previously provided. GADNR request that this permit be valid for a term of 10 years beginning January 1, 2012.

- B. The specific location of the activity. Please include latitude/longitude coordinates if possible.

Waters open to commercial shad fishing can be found on pages 17-18 of "Georgia's Commercial Saltwater Fishing Regulations" that was previously provided.

- C. For a general incidental take application, include an estimate of the total level of activity expected to be conducted.

According to mandatory individual records (trip tickets) reported to GADNR Coastal Resources Division (CRD), from 2007 through 2011 total statewide annual commercial shad fishing trips in GA have declined from 388 trips to 241 trips/yr and averaged 316 trips/yr during this time. GADNR anticipates that commercial fishing activity will remain stable or slightly decline over the duration of the requested permit.

- VI. The application must include a conservation plan based on the best scientific and commercial data, which specifies:

- A. The anticipated impact of the proposed activity on the listed species, including:

1. The estimated number of animals of the listed species and, if applicable, the subspecies or population group, and range.

Estimated total number of shortnose sturgeon incidentally captured by shad set-net fishermen in the Altamaha River ranged from 53-498 fish during 2007-2009 (Bahn and Petereson, 2010). This same study also estimated the Altamaha River population at approximately 6,300 fish. New commercial shad regulations that were instituted January 1, 2011 should substantially reduce incidental bycatch of sturgeon since these rules closed the section of the Altamaha River with the highest bycatch rates. Bahn and Peterson (2010) stated "In fact, we estimate that more shortnose sturgeon were incidentally captured in the upper river during January 2009 (333 fish) than in all months of all three years combined in the lower river (216 fish; Table 2)". For the section of the Atlamaha that is currently open to commercial shad fishing, this study reported that during 2007-2009 the

highest total annual bycatch of sturgeon by fishermen was estimated at 111 fish. GADNR also records incidental sturgeon captures while conducting an American shad fishery independent gill net survey on the Altamaha River and from 2001-2010 a total of 73 shortnose sturgeon were captured and released alive. The catch rate of shortnose sturgeon from the American shad gill net survey averaged 0.41 fish/day over this 10-yr period. During this same 10-yr period, the highest catch rate from any consecutive 3-year period (2001-2002) was 0.94 fish/day. These catch rates were significantly impacted by one year in which 41 of the 73 shortnose sturgeon were captured. Other than 2002, the highest number of shortnose sturgeon captured during the GADNR gill net survey in one year was 8 fish. From 2001-2010, reported commercial shad fishing trips on the Altamaha River averaged 265 trips. Utilizing catch rates from the GADNR gill net survey resulted in an estimated range of 109-250 shortnose sturgeon being incidentally captured per year in the commercial shad fishery. Due to the high variability in shortnose sturgeon bycatch rates, GADNR proposes utilizing 3-year running averages to monitor shortnose sturgeon bycatch. GADNR estimates that 3-year averages of incidental bycatch will not likely exceed 175 fish/yr in the Altamaha River.

Bahn and Peterson observed extremely low catch rates of Atlantic sturgeon in the commercial shad fishery during their 2007-2009 study, with only 6 Atlantic sturgeon being captured over the entire 3-year study. Due to the low catch rates an accurate estimate of total Atlantic sturgeon incidental capture could not be produced from the 2007-2009 study (personal comm). GADNR does record incidental Atlantic sturgeon captures while conducting an American shad fishery independent gill net survey on the Altamaha River and from 2001-2010 a total of 33 Atlantic sturgeon were captured and released alive. All of these were sub-adult fish with an average total length of 526 mm. The catch rate of Atlantic sturgeon from the American shad gill net survey averaged 0.19 fish/day over this 10-yr period. During this same 10-yr period, the highest catch rate from any consecutive 3-year period (2006-2008) was 0.41 fish/day. From 2001-2010, reported commercial shad fishing trips on the Altamaha River averaged 265 trips. Utilizing the catch rate of 0.41 fish/day results in an estimate of 109 Atlantic sturgeon being incidentally captured per year. Based on this data, GADNR estimates that 3-year averages of incidental bycatch will not likely exceed 140 fish/yr in the Altamaha River.

A similar study was completed on the Savannah River in the 1990's. Collins et al. (1996) reported that during the 1990-92 shad seasons a total of 240-shortnose sturgeon were captured by Savannah River shad fishermen. The Savannah River is open to commercial shad fishing from U.S. Hwy 301 (rkm 192), downstream to the Atlantic Ocean, an area approximately 103 rkm or 35% smaller than previously open to commercial shad fishing. Closing the upper portion of the river should decrease incidental bycatch and protect suspected spawning sites. It is estimated that 3-year averages of shortnose sturgeon incidental bycatch by GA shad fishermen will not exceed 75 fish/yr in the Savannah River.

GADNR does not conduct a fishery independent gill net survey on the Savannah River and does not have any recent data regarding the incidental bycatch of Atlantic sturgeon by the commercial shad fishery for the Savannah River. Therefore, GADNR proposes utilizing bycatch rate developed from the Altamaha fishery independent gill net survey to estimate the anticipated number of Atlantic sturgeon that may be intercepted in the Savannah River. From 2001-2010, Savannah River commercial shad fishing effort reported to GADNR has averaged an estimated 85 trips/yr. Utilizing the catch rate of 0.41 fish/day derived from the Altamaha River results in an estimate of 35 Atlantic sturgeon being incidentally captured per year. Based on this data, GADNR estimates that 3-year averages of incidental bycatch will not likely exceed 50 fish/yr in the Savannah River.

Incidental bycatch of sturgeon by the commercial shad fishery has not been evaluated in the Ogeechee River. This is a very small commercial fishery and based on the total number of commercial shad fishing trips from 2007-2011,

approximately 2% of the total statewide effort is exerted on the Ogeechee River. New regulations closed approximately 137 rkm or 66% of the river previously open to commercial fishing and also limited legal gear to drift nets only. GADNR believes that 3-year averages of incidental bycatch will likely not exceed 10 shortnose and 10 Atlantic sturgeon/yr in the Ogeechee River.

2. The type of anticipated taking, such as harassment, predation, competition for space and food, etc.

GA commercial regulations require that all sturgeon incidentally captured must be immediately released unharmed (pg 18 “Georgia’s Commercial Saltwater Fishing Regulations”)

3. The effects of the take on the listed species, such as descaling, altered spawning activities, potential for mortality, etc.

Bahn and Peterson (2010) reported a very low mortality rate of 2.3% for shortnose sturgeon that were captured in set nets targeting American shad in the Altamaha River. Sub-lethal effects are unclear.

- B. The anticipated impact of the proposed activity on the habitat of the species and the likelihood of restoration of the affected habitat.

The American shad gill net fishery is a low impact fishery and should have extremely minor physical affects on aquatic habitat utilized by shortnose sturgeon. In addition, the newly established commercial fishery boundaries will provide protection to confirmed and suspected spawning sites in Georgia’s rivers.

- C. The steps that will be taken to monitor, minimize, and mitigate such impacts, including:

1. Specialized equipment, methods of conducting activities, or other means.

Refer to page 18 of “Georgia’s Commercial Saltwater Fishing Regulations” for information on legal shad fishing gear.

2. Detailed monitoring plans.

See monitoring plan document that was previously submitted.

3. Funding available to implement measures taken to monitor, minimize and mitigate impacts.

In 2011, Georgia Department of Natural Resources management and monitoring of commercial fisheries operated under state appropriations and federal awards totaling approximately \$180,000. GADNR is mandated by ASMFC to annually monitor commercial shad fisheries and sturgeon populations. GADNR will utilize state appropriated funds, federal awards and existing staff to monitor the commercial shad fishery and incorporate sturgeon bycatch monitoring.

- D. The alternative actions to such taking that were considered and the reasons why those alternatives are not being used.

See alternative regulation document that was previously submitted.

- E. A list of all sources of data used in preparation of the plan, including reference reports, environmental assessments and impact statements, and personal communications with recognized experts on the species or activity who may have access to data not published in current literature.

Bahn and Peterson (2010)
Collins et al (1996)
GA Commercial Saltwater Fishing Regulations
GADNR (personal comm.)

An application for a certificate of inclusion under a General incidental take permit must include the following:

1. General incidental take permit under which the applicant wants coverage;
2. Applicant's name, address and telephone number (if the applicant is a partnership or corporate entity, then the applicable details);
3. Description of the activity the applicant wants covered under the general permit, including anticipated geographic range and season; and
4. Signed statement that the applicant has read and understood the general incidental take permit and the conservation plan, will apply with the applicable terms and conditions, and will fund the applicable measures of the conservation plan.

Modifications to Permits

Requests for modifications to incidental take permits should address all applicable sections of these instructions, including a detailed description of the proposed changes. Appropriate changes should also be made to the Conservation Plan. Modification requests involving an increased number of animals, additional species, an increased risk to the animals, or a significant change in the location of incidental take are subject to the 30-day public review and are granted or denied at the discretion of the Assistant Administrator for Fisheries.

Where to Send the Application

The application may be submitted electronically, if possible (either by email or by mailing a disk), but one signed original of the complete application must be sent to one of the following addresses.

Send applications for incidental take of all species except sea turtles and Pacific salmon to:

Chief, Endangered Species Division
National Marine Fisheries Service, F/PR3
1315 East-West Highway
Silver Spring, Maryland 20910
Telephone 301-713-1401
Fax 301-713-0376

Send applications for incidental take of sea turtles to:

Chief, Marine Mammal and Turtle Division
National Marine Fisheries Service, F/PR2
1315 East-West Highway
Silver Spring, Maryland 20910
Telephone 301-713-2322
Fax 301-713-4060
Web Site <http://www.nmfs.noaa.gov/pr/>

Please see separate application instructions for incidental take permits for sea turtles, available on-line at http://www.nmfs.noaa.gov/pr/permits/esa_permits.htm

Send applications for incidental take of anadromous fish in the Pacific to one of these offices:

Pacific Salmon
Northwest Regional Office
National Marine Fisheries Service
7600 Sand Point Way NE
Building 1
Seattle, WA 98115
Phone: (206) 526-6150
Fax: (206) 526-6426

NMFS Northern California Coast Salmon
National Marine Fisheries Service
1655 Heindon Road
Arcata, CA 95521
Phone: (707) 825-5163
Fax: (707) 825-4840

NMFS Central California Coast Salmon
National Marine Fisheries Service
777 Sonoma Ave., Room 325
Santa Rosa, CA 95404
Phone: (707) 575-6050
Fax: (707) 578-3435

NMFS California Central Valley Salmon
National Marine Fisheries Service
650 Capitol Mall, Suite 8-300
Sacramento, CA 95819
Phone: (916) 930-3600 Fax: (916) 930-3629

NMFS Southern California Salmon
National Marine Fisheries Service
501 West Ocean Blvd
Long Beach, CA 90802-4250
Phone: (562) 980-4020 Fax: (562) 980-4027

GA American Shad Fishery Sturgeon Bycatch Monitoring Plan

The Georgia Department of Natural Resources (GADNR) proposes to utilize a combination of a trip ticket system and direct observations to monitor the bycatch of shortnose sturgeon in the commercial shad fishery. Georgia regulations currently require commercial fishermen to complete trip tickets to document species, sex and pounds of shad harvested each day. In addition to the information on shad harvest, these tickets capture the fisherman's name and license number, name of dealer that purchases fish, river fished, gear type (set or drift net), length of net, total soak time, and number of net sets. Fishermen and/or dealers are required to return completed trip tickets to the Georgia Department of Natural Resources by the 10th of each following month (i.e. January tickets would be due by February 10). The current trip ticket will be modified to require fisherman to record information on sturgeon bycatch (total numbers of sturgeon intercepted and released) and data will be utilized to monitor sturgeon interactions with the shad fishery. Modified trip tickets will have rows and/or columns for fishermen to separately record incidental catches of shortnose and Atlantic sturgeon.

GADNR will make a concerted effort to educate commercial shad fishermen on the importance of both accurately recording sturgeon incidental catches and returning the trip tickets in a timely manner, at least by the 10th of each following month. GADNR will develop an informational packet on sturgeon identification, proper handling (emphasizing the importance of fishermen frequently checking their nets and immediately releasing any sturgeon that are incidentally caught), and the importance of reporting incidental sturgeon catches. Prior to each shad season, this informational packet will be provided to all known commercial shad fishermen.

A list of names and addresses of commercial shad fishermen will be compiled from prior trip tickets, the commercial fishing license database, and a list of cooperators in shad tagging studies. A set of trip tickets, self-addressed return envelopes, and information on how to obtain additional trip tickets will also be provided to each fisherman on this list. In addition to these direct handouts and mailings, GADNR Law Enforcement staff will be supplied additional trip tickets to be provided to shad fishermen encountered during routine patrol.

According to results reported by Bahn and Peterson (2010), estimated shortnose sturgeon bycatch determined from direct observations of commercial shad fishing activities did not differ significantly from those estimated from commercial shad fishermen log book data for the same time period. However, GADNR believes that it is still important to periodically observe commercial shad fishing activities. Thus, GADNR staff will utilize the same list of names obtained from trip tickets, the commercial fishing license database, and the list of cooperators in shad tagging studies to establish contact information (i.e. phone numbers) for a subset of individuals that commercially fish for shad on the Altamaha, Ogeechee, and Savannah rivers.

Once contact information has been established for a set of fishermen for each river, GADNR staff will contact fishermen to determine when they will be fishing and to establish a time and location to observe fishermen pulling their nets. The goal will be to make observations within 24-48 hours of contact with the fisherman. Numbers of direct observations for each river will be based on current shad fishing pressure and spawning migrations of shad and sturgeon.

GADNR will attempt to observe a minimum of 10% of the commercial shad fishing trips on each river. Based on averaging the last 3 years of commercial fishing effort, GA DNR would need to observe approximately 25, 5, and 1 trip each year, respectively, for the Altamaha, Savannah, and Ogeechee rivers. Since commercial shad fishing effort is extremely low on the Ogeechee River, GADNR will attempt to observe at least 2 trips per year on the Ogeechee River.

Monthly observations for a river system may also vary. Shad fishing effort is typically lower on all three rivers in January than in February and March due to the fact that shad abundance is less early in the season. Therefore, the number of direct observations will likely be lower for January than for the following months.

GADNR monitors the shad spawning migration every week during the commercial shad season, which allows staff to know when the spawning run and resulting fishing pressure are peaking. This information will allow GADNR to make necessary adjustments in monitoring efforts to ensure that at least 10% of all commercial shad fishing trips are observed annually. Monitoring efforts will also be adaptive to the timing of the sturgeon spawning migration and the number of sturgeon intercepts. GADNR will increase direct observations if high numbers of sturgeon intercepts are detected. GADNR is confident that this approach will ensure that an adequate number of observations are made during the peak of both the shad and sturgeon spawning migrations so that sturgeon bycatch is accurately estimated.

If unusually high catch rates are being observed, GADNR will immediately increase law enforcement presence and educational efforts. Staff will also begin evaluating additional modifications to the commercial shad fishing regulations for the next year. Data collected from the trip tickets and direct observations will be summarized and provided to the National Marine Fisheries Service no later than the end of February, March, and April each year.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

CHAPTER 1
INTRODUCTION AND LITERATURE REVIEW¹

¹ Bahn, R. A., D. J. Farrae, and D. L. Peterson *in part to be submitted to*
Reviews in Fish Biology and Fisheries summer 2010

22 The shortnose sturgeon, *Acipenser brevirostrum* LeSueur 1818, is the
23 smallest member of Acipenseridae, and inhabits coastal rivers and estuaries
24 along the Atlantic Coast of North America from the St. John River, Canada, to the
25 St. John's River in northeast Florida (Vladykov and Greeley 1963; Moser and
26 Ross 1995; Bain et al. 2007). Like other members of the genus, shortnose
27 sturgeon are long-lived, late maturing, diadromous fishes with a protracted
28 spawning periodicity (Vladykov and Greeley 1963; Bemis and Kynard 1997).
29 Historical abundance estimates are scarce, however, shortnose sturgeon were
30 exploited for decades along with the sympatric Atlantic sturgeon, *Acipenser*
31 *oxyrinchus* (Smith et al. 1984). During the last century, shortnose sturgeon had
32 become sufficiently rare that they were listed as an endangered species in the
33 United States in 1967 (National Marine Fisheries Service (NMFS) 1998). Today,
34 few healthy populations exist and many anthropogenic factors impede restoration
35 efforts (Kynard 1997). Many populations, particularly in southern rivers, continue
36 to be threatened with extinction. With federal protection in place, the two primary
37 factors currently affecting population recovery in the Southeastern U.S. are
38 habitat degradation and fishing mortality as a result of unintended capture or
39 "bycatch" in commercial fisheries targeting other species (Collins et al. 2000).

40

41 **Life History**

42 Sturgeon are long-lived, late maturing, diadromous fishes with a
43 protracted spawning periodicity (Bemis and Kynard 1997). Populations of
44 shortnose sturgeon have life history differences in their northern and southern

45 ranges, but southern populations have not been well studied. In southern rivers,
46 shortnose sturgeon mature sooner, spawn earlier in the year, grow faster, and
47 have shorter life spans compared to those in the northern part of the range
48 (Vladykov and Greeley 1963; Heidt and Gilbert 1978; Dadswell 1979).

49 As an amphidromous species, shortnose sturgeon require riverine habitats
50 to complete their life cycle, but they will migrate to estuarine and marine habitats
51 for purposes other than spawning (Bemis and Kynard 1997). Shortnose
52 sturgeon typically mature at 500-600 mm total length (TL), which is reached by 2-
53 3 years for males and 3-5 years for females in southern populations (Dadswell
54 1979; Kynard 1997). After maturity, males spawn every 1-2 years; females
55 spawn every 3-5 years (Dadswell 1979). Southern shortnose sturgeon are
56 estimated to live less than 20 years, compared to 30-67 years for their northern
57 counterparts (Rogers and Weber 1994; Kynard 1997). Spawning occurs from
58 late January (D. Peterson, unpublished data) to March in southern rivers, where
59 shortnose sturgeon migrate to the upstream portion of their population range
60 (Heidt and Gilbert 1978; Bain 1997; Kynard 1997). In the Altamaha River,
61 spawning is thought to occur between river kilometer (rkm) 167 and 215 (DeVries
62 2006; D. Peterson, unpublished data).

63

64 **Bycatch**

65 Fishing mortality from bycatch is a problem for many species that have life
66 histories dependent on late maturation and protracted spawning periodicity
67 (Boreman 1997; Stein et al. 2004). Although they are long-lived, sturgeons only

68 spawn once every 3-5 years (Dadswell 1979). Hence, sturgeon populations are
69 especially sensitive to loss of reproductive potential from bycatch mortality
70 (Boreman 1997).

71 Bycatch of sturgeon in riverine, estuarine, and marine fisheries is a threat
72 to the recovery of many sturgeon populations (Stein et al. 2004; Munro et al.
73 2007). Although shortnose sturgeon are federally protected, they are frequently
74 captured across their range in commercial fisheries targeting other riverine
75 species (Kynard 1997). Most of this bycatch occurs in anchored and drifted gill
76 net fisheries for American shad (*Alosa sapidissima*; Collins et al. 1996; Kynard
77 1997).

78 Bycatch of shortnose sturgeon by commercial shad fisheries is well
79 documented (Heidt and Gilbert 1978; Dadswell 1979; Collins et al. 1996; Weber
80 1996; Kynard 1997; Collins et. al 2000). Collins et. al (2000) states that the use
81 of anchored gill nets in essential habitats by commercial fishermen is a threat to
82 the recovery of sturgeon populations. In Georgia, commercial shad fisheries are
83 open from January 1 to March 31. Based on total fishing effort, the shad fishery
84 is one of the largest commercial fisheries operated in Georgia (Collins et al.
85 1996). Adult shortnose sturgeon are vulnerable to incidental capture by
86 commercial shad fisheries because their upstream spawning migration coincides
87 with the peak commercial fishing effort (Collins et al. 2000). Soak time directly
88 affects sturgeon mortality rates in anchored gill net fisheries (Atlantic Sturgeon
89 Status Review Team (ASSRT) 2007). In the Altamaha River, commercial
90 fishermen use both drifted and anchored gill nets in different portions of the river.

91 Anchored gill nets must have a minimum of 11.43 cm stretched mesh with a
92 maximum length of 30.48 m. Nets must be spaced at least 182.88 m apart with
93 one end attached to the shore, allowing open fish passage through at least ½ of
94 the river channel. Most gill nets deployed upstream of the estuary in the
95 Altamaha River from 2004-08 were anchored gill nets (D. Peterson, unpublished
96 data). Drifted gill nets can be used throughout the river, but are mostly used in
97 the estuary. Only drifted gill nets are permitted in the Altamaha Sound. Collins
98 et al. (1996) and Stein et al. (2004) state that the time non-target species spend
99 tangled in drifted gill nets is likely less than that of anchored gill nets because
100 drifted gill nets must be tended constantly to prevent these nets from becoming
101 entrained on benthic debris. Collins et al. (1996) also states that catch per unit
102 effort (CPUE) of sturgeon may be lower in drifted gill nets because they often do
103 not fish the lower portion of the water column.

104 Previous studies of shad fisheries have shown that shortnose sturgeon
105 bycatch can be significant. Collins et al. (1996) reported that shad fishermen
106 captured 240 shortnose sturgeon from 1990-92 in the Savannah River. In this
107 study, 97% of captured shortnose sturgeons were mature adults (TL 560 -1060
108 mm). In 1994, the shortnose sturgeon population in the Savannah River was
109 calculated to be 1676, but this estimate was deemed incorrect because not all
110 assumptions of the Schnabel model were met (NMFS 1998).

111 Both shortnose sturgeon and American shad migrate to upstream
112 spawning sites in southern rivers during February and March (Hall et al. 1991;
113 Collins and Smith 1995). Spawning shortnose sturgeon leave the estuary in mid-

December, migrating upstream for several hundred kilometers throughout the winter (DeVries 2006). Although Georgia's commercial shad fishery does not open until January, DeVries (2006) documented adult shortnose sturgeon continuing upstream migrations throughout February and early March. Hence, the temporal and spatial overlap of shortnose sturgeon migrations and the commercial fishery creates a potential for incidental capture of spawning shortnose sturgeon. Although commercial fishermen must immediately release any sturgeon caught, soak time of commercial gear is not regulated. Consequently, most commercial fishermen check their nets once daily, thereby increasing the potential for injury or death of entangled shortnose sturgeon. Aside from direct mortality caused by long soak times of anchored gill nets, prolonged entanglement of sturgeon can have sublethal effects, but they have not been well studied (Moser and Ross 1995; Boreman 1997; Kynard 1997). Previous studies have reported instances where radio-tagged shortnose sturgeon aborted their spawning migrations after being captured in commercial anchored gill nets (Moser and Ross 1995; Weber 1996).

Mortality and injury of sturgeons because of bycatch in shad fisheries has been identified as a serious threat to southern sturgeon populations (Kynard 1997; Collins et al. 2000). Because the Altamaha River contains the largest population of adult shortnose sturgeon (~1800 individuals) south of the Delaware River, bycatch of shortnose sturgeon in the shad fishery is a concern to both state and federal agencies (NMFS 1998; DeVries 2006). The observed mortality rate of over 30% in the Altamaha River shortnose sturgeon population (DeVries

2006) is high compared to 22% in the Hudson River (Secor and Woodland 2005). The effect of bycatch on the mortality rate of shortnose sturgeon in the Altamaha River is unknown; however, Collins et al. (1996) documented a 16% mortality rate and a 20% injury rate among shortnose sturgeon captured in the commercial shad fishery of Winyah Bay, SC.

Research Objectives and Justification

The objective of my study was to estimate the bycatch of shortnose sturgeon in the commercial shad fishery of the Altamaha River, GA. The National Marine Fisheries Service has identified studies of shortnose sturgeon bycatch in commercial fisheries as a research priority throughout the Atlantic Coast (NMFS 1998). In a previous study of shortnose sturgeon bycatch in the Savannah River, Collins et al. (1996) recommended the use of a standardized creel survey methodology for future assessments in other southern rivers. Because the effects of sturgeon bycatch have not been well studied, little is known about how Georgia's commercial shad fisheries may be affecting recovery of shortnose sturgeon throughout the state. Although surveys conducted during the 1980s and 1990s documented mortality of shortnose sturgeon in Georgia's shad fisheries, the population level effects were difficult to quantify because shortnose sturgeon abundance estimates were not available (Collins et al. 1996). A recent study by DeVries (2006) however, reported new abundance estimates for Altamaha River shortnose sturgeon, providing a context for quantifying the effects of bycatch. The results of this study provide the first quantified estimates

of bycatch and mortality rates of shortnose sturgeon in the Altamaha River
commercial shad fishery. The application of these results will provide a
framework for evaluating current commercial shad fishing regulations in Georgia
and on other rivers where shortnose sturgeon populations exist.

References

- Atlantic Sturgeon Status Review Team (ASSRT). 2007. Status Review of Atlantic sturgeon (*Acipenser oxyrinchus oxyrinchus*). Report to National Marine Fisheries Service, Northeast Regional Office. February 23, 2007. 174 pp.
- Bain, M. B. 1997. Atlantic and shortnose sturgeons of the Hudson River: common and divergent life history attributes. *Environmental Biology of Fishes* 48:347-358.
- Bain, M. B., N. Haley, D. L. Peterson, K. K. Arend, K. E. Mills, and P. J. Sullivan. 2007. Recovery of a US Endangered Fish. *PLoS ONE* 2(1): e168.
- Bemis, W. E. and B. Kynard. 1997. Sturgeon rivers: An introduction to acipensiform biogeography and life history. *Environmental Biology of Fishes* 48:167-183.
- Boreman, J. 1997. Sensitivity of North American sturgeons and paddlefish to fishing mortality. *Environmental Biology of Fishes* 48:399-405.
- Collins, M. R. and T. I. J. Smith. 1995. Characteristics of the adult segment of the Savannah River population of shortnose sturgeon. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 47(1993):485-491.
- Collins, M. R., S. G. Rogers, and T. I. J. Smith. 1996. Bycatch of sturgeons along the southern Atlantic Coast of the USA. *North American Journal of Fisheries Management* 16:24-29.

205 Collins, M. R., S. G. Rogers, T. I. J. Smith, and M. L. Moser. 2000. Primary
 206 factors affecting sturgeon populations in the southeastern United States:
 207 Fishing mortality and degradation of essential habitats. *Bulletin of Marine*
 208 *Science* 66(3):917-928.

209 Dadswell, M. J. 1979. Biology and population characteristics of the shortnose
 210 sturgeon, *Acipenser brevirostrum* LeSueur 1818 (Osteichthes:
 211 *Acipenseridae*), in the Saint John River Estuary, New Brunswick, Canada.
 212 *Canadian Journal of Zoology* 57:2186-2210.

213 DeVries, R. J. 2006. Population dynamics, movements, and spawning habitat of
 214 the shortnose sturgeon, *Acipenser brevirostrum*, in the Altamaha River
 215 system, Georgia. M.S. Thesis, University of Georgia, Athens, Georgia.

216 Hall, J. W., T. I. J. Smith, and S. D. Lamprecht. 1991. Movements and habitats
 217 of shortnose sturgeon, *Acipenser brevirostrum*, in the Savannah River.
 218 *Copeia* 1991:695-702.

219 Heidt, A. R. and R. J. Gilbert. 1978. The shortnose sturgeon in the Altamaha
 220 River drainage, Georgia. Pages 54-60 *in* R. R. Odum and L. Landers,
 221 editors. *Proceedings of the rare and endangered wildlife symposium*.
 222 Georgia Department of Natural Resources, Game and Fish Division,
 223 Technical Bulletin WL 4, Athens, Georgia.

224 Kynard, B. 1997. Life history, latitudinal patterns, and status of the shortnose
 225 sturgeon, *Acipenser brevirostrum*. *Environmental Biology of Fishes*
 226 48:319-334.

227 Moser, M. L. and S. W. Ross. 1995. Habitat use and behavior of shortnose and
 228 Atlantic sturgeons in the lower Cape Fear River, North Carolina.
 229 Transactions of the American Fisheries Society 124:225-234.

230 Munro, J., R. E. Edwards, and A. W. Kahnle. 2007. Anadromous sturgeons:
 231 Habitats, threats, and management. Synthesis and summary. Pages 1-
 232 15 in J. Munro, D. Hatin, J. E. Hightower, K. McKown, K. J. Sulak, A. W.
 233 Kahnle, and F. Caron, editors. Anadromous sturgeons: habitats, threats,
 234 and management. American Fisheries Society, Symposium 56, Bethesda,
 235 Maryland.

236 National Marine Fisheries Service (NMFS). 1998. Final recovery plan for the
 237 shortnose sturgeon, *Acipenser brevirostrum*. Prepared by the Shortnose
 238 Sturgeon Recovery Team for the National Marine Fisheries Service, Silver
 239 Spring, Maryland. 104pp.

240 Rogers, S. G. and W. Weber. 1994. Movements of shortnose sturgeon in the
 241 Altamaha River System, Georgia. Contributions Series No. 57. Coastal
 242 Resources Division, Georgia Department of Natural Resources.
 243 Brunswick, Georgia.

244 Secor, D. H., and R. J. Woodland. 2005. Recovery and status of shortnose
 245 sturgeon in the Hudson River. Hudson River Foundation for Science and
 246 Environmental Research, Inc. 105pp.

247 Smith, T. I. J., D. E. Marchette, and G. F. Ulrich. 1984. The Atlantic sturgeon
 248 fishery in South Carolina. North American Journal of Fisheries
 249 Management 4:164-176.

- Stein, A. B., K. D. Friedland, and M. Sutherland. 2004. Atlantic sturgeon marine
bycatch and mortality on the continental shelf of the northeast United
States. North American Journal of Fisheries Management 24:171-183.
- Vladykov, V. D. and J. R. Greeley. 1963. Order Acipenseroidei. Pages 24-60 in
V. H. Olsen, editor. Fishes of the western North Atlantic, part III. Memoirs
of the Sears Foundation for Marine Research, New Haven Connecticut.
630 pp.
- Weber, W. 1996. Population size and habitat use of shortnose sturgeon,
Acipenser brevirostrum, in the Ogeechee River System, Georgia. M.S.
Thesis, University of Georgia, Athens, Georgia.

273

274

275

CHAPTER 2

276

BYCATCH OF SHORTNOSE STURGEON IN THE COMMERCIAL SHAD

277

FISHERY OF THE ALTAMAHA RIVER, GEORGIA²

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

² Bahn, R. A. and D. L. Peterson *to be submitted to*

Transactions of the American Fisheries Society summer 2010

295 **Abstract**

296 Although the shortnose sturgeon (*Acipenser brevirostrum*) has been
297 federally protected as an endangered species since 1967, incidental capture of
298 shortnose sturgeon in commercial shad fisheries has been documented as a
299 source of mortality that may limit recovery of some populations. As such,
300 shortnose sturgeon bycatch assessments were recently identified as a priority by
301 the National Marine Fisheries Service, as part of the iterative process of
302 identifying and reducing threats to East Coast sturgeon. The objective of our
303 study was to estimate total bycatch and mortality of shortnose sturgeon in the
304 anchored gill net portion of the Altamaha River commercial shad fishery from
305 2007 - 09. Using a roving creel survey design, we conducted on-the-water
306 counts of commercial shad nets to estimate fishing effort. Catch-per-unit effort
307 was estimated from log books and direct observations of net retrievals by
308 randomly selected commercial fishermen. During the 3 years of the study, total
309 estimated bycatch of shortnose sturgeon was 71, 53, and 498 fish, respectively.
310 Catch rates were highest during January and February of 2009 in upriver
311 commercial nets near previously confirmed spawning locations in the river.
312 Mortality of captured shortnose sturgeon was low in all three years (< 8%),
313 although we did not assess post-release survival. Future studies are needed to
314 better assess population level effects and sub-lethal effects of incidental capture
315 on shortnose sturgeon. Because bycatch is highly variable annually, future
316 studies need to be conducted over several seasons and throughout the extent of
317 the population range in a particular river.

318 **Introduction**

319 Shortnose sturgeon (*Acipenser brevirostrum*) are an amphidromous
320 species that ranges from the St. John River, Canada, to the St. John's River in
321 northeast Florida (Vladykov and Greeley 1963). Although shortnose sturgeon
322 were once common in most major East Coast river systems, commercial
323 exploitation and habitat degradation have reduced populations significantly
324 (Kynard 1997; Collins et al. 2000). The shortnose sturgeon has been federally
325 listed as an endangered species since 1967 (National Marine Fisheries Service
326 (NMFS) 1998).

327 Northern and southern populations of shortnose sturgeon are known to
328 exhibit several important differences in life history; however, southern
329 populations have not been well studied. In southern rivers, shortnose sturgeon
330 mature sooner, spawn earlier in the year, grow faster, and have shorter life spans
331 compared to those in the northern part of the range (Vladykov and Greeley 1963;
332 Heidt and Gilbert 1978; Dadswell 1979). As an amphidromous species,
333 shortnose sturgeon require riverine habitats to complete their life cycle, but they
334 will feed in estuarine and marine habitats during the winter months (Bemis and
335 Kynard 1997). Shortnose sturgeon typically mature at 500-600 mm total length
336 (TL), which is reached by 2-3 years for males and 3-5 years for females in
337 southern populations (Dadswell 1979; Kynard 1997). After maturity, males
338 spawn every 1-2 years; females every 3-5 years (Dadswell 1979). Southern
339 shortnose sturgeon are estimated to live less than 20 years, compared to 30-67
340 years for their northern counterparts (Rogers and Weber 1994; Kynard 1997).

Spawning occurs from late January (D. Peterson, unpublished data) to March in southern rivers, where shortnose sturgeon migrate to the upstream portion of their population range (Heidt and Gilbert 1978; Bain 1997; Kynard 1997).

Although shortnose sturgeon have been federally protected for more than 40 years, they are frequently captured across their range in commercial fisheries targeting other riverine species (Kynard 1997). Most of this "bycatch" occurs in anchored and drifted gill net fisheries for American shad (*Alosa sapidissima*; Collins et al. 1996; Kynard 1997). Several authors have shown that fishing mortality from bycatch poses an especially serious threat to species with reproductive strategies that depend on late maturation and protracted spawning periodicity (Boreman 1997; Stein et al. 2004; Munro et al. 2007). Despite their long life spans, shortnose sturgeon spawn only once every 2-5 years after reaching maturity (Dadswell 1979), making them particularly sensitive to the cumulative losses of reproductive potential resulting from chronic bycatch mortality (Boreman 1997).

Bycatch of shortnose sturgeon in commercial shad fisheries has been well documented (Heidt and Gilbert 1978; Dadswell 1979; Collins et al. 1996; Weber 1996; Kynard 1997; Collins et al. 2000), but population level effects are poorly understood. Previous studies of commercial shad fisheries have shown that shortnose sturgeon bycatch can be significant and Collins et al. (2000) suggest that this bycatch may be among the most serious impediments to the recovery of southern shortnose sturgeon populations. In South Carolina, previous studies have shown that shad fishermen captured 240 shortnose sturgeon from 1990-92

in the Savannah River and that 97% of those captured were mature adults (TL 560 -1060 mm; Collins et al. 1996). In 1994, the shortnose sturgeon population in the Savannah River was estimated at 1,676 individuals, suggesting that annual bycatch in this commercial fishery may have resulted in the incidental capture of up to 15% of the entire adult population.

Although shortnose sturgeon accidentally captured in commercial shad fisheries must be immediately released, delayed mortality and injury resulting from incidental capture has been identified as a serious threat to populations in several southern rivers (Kynard 1997; Collins et al. 2000). Collins et al. (1996), for example, documented a 16% mortality rate and a 20% injury rate for shortnose sturgeon captured in commercial shad nets in Winyah Bay, SC.

In many Atlantic Coast rivers, spawning runs of American shad largely overlap with those of shortnose sturgeon (Hall et al. 1991; Collins et al. 1996; NMFS 1998). Consequently, adult shortnose sturgeon are particularly vulnerable to incidental capture in commercial shad fisheries because their annual upstream migrations coincide with the peak commercial fishing effort (Collins et al. 2000). Because bycatch is a known problem for recovering shortnose sturgeon populations, NMFS has identified studies of bycatch in commercial fisheries as a research priority as part of the iterative process of identifying and reducing threats to the recovery of sturgeons (NMFS 1998).

In Georgia, the Altamaha River contains the largest population of shortnose sturgeon (~1,800 adults) within the southern portion of the range (Peterson and DeVries 2006). Hence, bycatch of shortnose sturgeon in the

Altamaha commercial shad fishery is of particular concern to both state and federal management agencies (NMFS 1998). In the Altamaha River, the commercial shad fishery is open from 1 January to 31 March and fishermen may use both drifted and anchored gill nets, depending on where they operate. Drifted gill nets can be used throughout the river, but their use is largely restricted to estuarine waters because of an abundance course woody debris above the head of tide. Anchored gill nets can be used upstream of the estuary. Because drifted nets must be tended constantly, the average duration of fish entanglement is typically much lower in drifted nets compared to anchored nets (Collins et al. 1996; Stein et al. 2004). Collins et al. (1996) also noted that catch-per-unit-effort (CPUE) of shortnose sturgeon may be lower in drifted gill nets because they usually do not extend down to the benthos where shortnose sturgeon are typically found. Anchored nets must have a minimum of 11.43-cm stretched mesh with a maximum length of 30.48 m. Nets must be spaced at least 182.88 m apart with one end attached to the shore, allowing unhindered fish passage through at least ½ of the river channel. Most gill nets deployed upstream of the estuary in the Altamaha River from 2004-06 were anchored gill nets (D. Peterson, unpublished data).

In southern rivers, both shortnose sturgeon and American shad migrate to upstream spawning sites in southern rivers from December to March (Hall et al. 1991; Collins and Smith 1993; Bahn et al. 2010). Although Georgia's commercial shad fishery does not open until January, DeVries (2006) documented adult shortnose sturgeon moving upstream in December, and continuing their

410 migration through February and early March. Hence, the temporal and spatial
411 overlap of shortnose sturgeon spawning migrations and the commercial shad
412 fishery creates a potential for incidental capture of spawning shortnose sturgeon.
413 Soak time directly affects sturgeon mortality rates in anchored gill net fisheries
414 (Atlantic Sturgeon Status Review Team (ASSRT) 2007). Although commercial
415 fishermen must immediately release any shortnose sturgeon caught, soak time of
416 commercial gear is not regulated. Consequently, most commercial fishermen
417 check their nets only once daily, thereby increasing the potential for injury or
418 death of entangled shortnose sturgeon. Aside from direct mortality caused by
419 long soak times of anchored gill nets, sublethal effects of prolonged
420 entanglement have been documented for shortnose sturgeon (Moser and Ross
421 1995; Kynard 1997). Previous studies have reported several instances where
422 radio-tagged shortnose sturgeon aborted spawning migrations after capture in
423 anchored gill nets (Moser and Ross 1995; Weber 1996).

424 Because the effects of sturgeon bycatch have not been well studied, little
425 is known about how Georgia's commercial shad fisheries may be affecting
426 recovery of shortnose sturgeon throughout the state. The objective of our study
427 was to quantify bycatch of shortnose sturgeon in the anchored gill net
428 commercial shad fishery in the Altamaha River from 2007-2009. Although
429 surveys conducted during the 1980s and 1990s documented mortality of
430 shortnose sturgeon in Georgia's shad fisheries, the population level effects were
431 difficult to quantify because shortnose sturgeon abundance estimates were not
432 available (Collins et al. 1996). A recent study by Peterson and DeVries (2006)

however, provided new abundance estimates for Altamaha River shortnose sturgeon, providing the key context necessary for quantifying the effects of bycatch in this population. In this study, we report the first quantified estimates of total bycatch and mortality rates of shortnose sturgeon in the Altamaha River commercial shad fishery. The application of these results may provide an important new framework for evaluating current commercial shad fishing regulations in Georgia and on other rivers where shortnose sturgeon populations coexist with commercial shad fisheries.

Study Site

The Altamaha River is formed on the coastal plain of Georgia by the confluence of the Ocmulgee and Oconee rivers near Hazlehurst, GA (Figure 1). The river flows southeast 215 km to the Atlantic Ocean near Darien, GA. The watershed contains approximately 800 km of unimpounded channel habitat accessible to diadromous fishes including shortnose sturgeon. Because the stream drains over one-quarter of the state, channel depths are highly variable depending on seasonal rainfall patterns and hydropower operation on reservoirs in the Ocmulgee and Oconee rivers. The head of tide is typically located between rkm 45-55, again depending on discharge. Mean channel depth is typically 50-70 m in width and 2-3 m in depth (Heidt and Gilbert 1978). Depths greater than 10 m are common in the tidally influenced section of the river. Deep cutbanks (10 m and greater) and channel scours below bridges are found above the head of tide.

Methods

Experimental Design

To estimate the number of shortnose sturgeon incidentally captured in the commercial shad fishery, we conducted a standardized fishery assessment of the Altamaha River mainstem from 1 January to 31 March, 2007-2009. Based on a priori knowledge of known and suspected shortnose sturgeon spawning locations (Peterson and DeVries 2006), we divided the river into two strata (Figure 1). The upper river stratum began at rkm 215 and extended downstream to rkm 184. The lower river stratum began at rkm 184 and extended downstream to rkm 21.

Using a roving creel survey design (Malvestuto 1996), we conducted weekly counts of anchored gill nets by traversing the entire 215 rkm of the study area by boat. In 2007 and 2008, these weekly counts were completed in two consecutive days, beginning with a random starting location and direction of travel. In 2009, counts were conducted continuously from upstream to downstream, so that they could be completed in one day. In each year, a running count of shad nets was made by checking each floating net buoy encountered during these counts to confirm that an actively fishing net was present. Nets that did not comply with published fishing regulations were included in all net count totals, but were not reported to law enforcement until the end of the season to prevent any potential bias in fisherman behavior.

For each month of each season, CPUE was obtained using a combination of direct observations of net retrievals and log books from five to seven commercial fishermen. The individual fishermen selected to provide this

information were chosen based on the river section where they fished and their willingness to participate in the study. Specific locations of their nets were independent of each other and interspersed throughout the study area. Each fisherman was compensated US\$500 annually in return for their cooperation in allowing us to observe randomly selected net pulls and for keeping accurate log books of both effort and catch. Direct observations of fishermen were randomized with some allowance for the individual schedules of each. Fishermen were not compensated, however, until accuracy of log books had been verified at the conclusion of each fishing season. Accuracy of log books was verified using two methods: 1) using a matched-pair t-test to compare days when observers were and were not present, and 2) using a matched-pair t-test to identify any significant differences of effort and catch data in log books versus those obtained through direct observations.

Direct observations of catch were conducted at least three times for each participating fishermen during each shad season. During each observation, we followed the fishermen to his nets in a separate boat so that we could record the number of each species captured as the net was retrieved. After all nets had been pulled, we recorded soak times, net dimensions, and mesh sizes. During 2008 and 2009, we also recorded total length (TL) and weight (g) of each shortnose sturgeon that was captured.

Data Analysis

To estimate total annual effort, we first calculated the mean number of nets fished in each stratum for each month of the season. Total net-hours was

then calculated for each month based on the number of nets counted each week and the total number of fishing hours that the season was open. This included 12 hours for opening and closing days and 24 hours for all other days. Total monthly fishing effort for each stratum was then calculated using the formula:

Total fishing effort (net hrs) = $\Sigma ((\text{Mean nets observed} / \text{mo}) \times (\text{Total fishing hrs} / \text{mo}))$

Accuracy of log book data from each fisherman was evaluated using a one sample matched-pair t-test ($\alpha = 0.05$) to compare the mean of the differences between days when observers were and were not present. We then used a one sample matched-pair t-test ($\alpha = 0.05$) to compare the mean of the differences between logged and observational data. To perform this test, the total annual number of shortnose sturgeon observed in the catch of each individual fishermen was standardized to the total number of net-hours recorded in his log book to calculate a monthly CPUE for each fisherman. Estimates of total monthly effort and catch were then calculated for each fisherman by supplementing the direct observational data with those from the log books recorded on days when observers were not present. A total monthly CPUE for shortnose sturgeon (SNS) was then estimated for each stratum using the formula:

CPUE = $(\text{Number SNS observed} + \text{number SNS logged}) / \text{Total net hrs}$

The variance of each of these estimates was used to calculate 0.95 confidence intervals. Assuming a linear relationship between effort and catch, we then estimated total monthly bycatch in each stratum using the formula:

525 Total monthly catch = (Total fishing hrs / mo) x (Mean monthly CPUE)
526 To identify any potential bias of mean CPUE calculations and to evaluate
527 the accuracy of CPUE variance estimates, we resampled our original data using
528 bootstrap analysis with replacement as described by Efron and Tibshirani (1994)
529 using SAS (SAS Institute, Cary, NC). We constructed resample sets of both 100
530 and 1,000 bootstrap samples to compare resampled means and variances to
531 those of the original data. For each month in each year in each stratum, we
532 randomly constructed 100 and 1,000 bootstrap samples containing the same
533 number of observations as the year-month-stratum data from which we were
534 resampling (e.g. from 70 field observations we generated 100 and 1,000
535 bootstrap resample sets with 70 observations each). For example, because the
536 original data from the lower stratum in January 2007 contained $i = 70$
537 observations, each bootstrap sample in the resample sets for the lower stratum
538 in January 2007 also contained $i = 70$ observations. We then calculated the
539 mean of each bootstrap sample and used these means to calculate grand means
540 and variances for the resample sets (by year-month-stratum, both 100 and 1,000
541 bootstrap samples) for comparison with original field data.

542

543 **Results**

544 During each of the three commercial fishing seasons sampled, we
545 conducted a total of 7-12 net counts totaling 1,358-2,328 rkm sampled annually.
546 We also collected catch data from 192-336 direct observations, and 10,382 –
547 15,410 net hours of log book entry data (Table 1). From these data, we

548 estimated that the total anchored gill fishery was comprised of 13-20 fishermen
549 annually. Of these participants, 2-4 operated in the upper stratum compared to
550 11-16 in the lower stratum. Over the three fishing seasons, data collected from
551 log books and direct observations annually accounted for 48% – 66% of all
552 fishing effort in the anchored gill net fishery (Table 1).

553 Total estimated effort for the entire anchored gill net fishery varied from
554 22,689 – 27,405 hours annually (Table 2). Weekly effort varied from 6 – 35 nets
555 per week during all three years of the study (Figure 2). In the upper river, fishing
556 effort peaked in February of each year; however, effort was not consistent among
557 months or years in the lower river (Figure 2). In the upper river, mean weekly
558 effort ranged from 0.8 – 4.0 nets per week. Mean weekly effort in the lower river
559 varied from 14.0 – 28.7 nets per week (Figure 2). Monthly effort varied from 495
560 – 1536 hours in the upper river compared to 5,712 – 11,700 hours in the lower
561 river (Table 2). Despite this variability, several spatial and temporal trends in
562 bycatch were evident. Most fishing effort (56.3%) occurred between rkm 35 -
563 100; however, most bycatch occurred in the upper river. In fact, we estimate that
564 more shortnose sturgeon were incidentally captured in the upper river during
565 January 2009 (333 fish) than in all months of all three years combined in the
566 lower river (216 fish; Table 2).

567 Analysis of log book data from all three years showed that catch data
568 recorded on days when observers were present was not significantly different
569 than on days when observers were absent ($p > 0.61$ for all three years).
570 Furthermore, total catch of shortnose sturgeon recorded during direct

observations was not significantly different than that provided in fishermen log books ($p > 0.42$ for all three years).

Total estimated bycatch varied from a low of 53 shortnose sturgeon in 2008 to 498 shortnose sturgeon in 2009 (Table 2). We estimated that 387 shortnose sturgeon were incidentally captured in the upper river during the 2009 shad season. No bycatch was recorded in the upper river in March during all three years of the study. In 2008 and 2009, bycatch peaked in February in the lower river (36 and 74 fish, respectively), and then declined in March (Table 2). This trend was not observed in 2007, however.

During months when shortnose sturgeon were incidentally captured in the upper river, CPUE was always higher than that of the lower river (Figure 3). For example, in January 2009, CPUE in the upper river was 0.5007 SNS/hr, compared to only 0.0015 SNS/hr in the lower river (Figure 3). During February 2007 and 2009, CPUE in the upper river was also higher (0.0126 and 0.0512 SNS/hr, respectively) than during the same period in the lower river (0.0019 and 0.0110 SNS/hr, respectively; Figure 3). During 2008 and 2009, CPUE in the lower river was lowest in January, followed by an increase of over 100% in February, and then a decline in March (Figure 3).

Bootstrap results of both the 100 and 1,000 resample sets showed that the observed mean CPUE values for our study were unbiased (Table 3). The associated standard errors for the randomized bootstrap sample sets were smaller than those of the estimated mean CPUE for both strata, indicating that

the variance estimates of mean CPUE in both strata were also accurate (Table 3).

Except for one juvenile fish captured in the upper river during January 2009, all shortnose sturgeon we observed during 2008 – 09 measured ≥ 590 mm TL. Most fish appeared to be in healthy condition and swam away after release, however, we were unable to assess any sublethal or post-release effects of incidental capture. Only 4 of the 172 shortnose sturgeon captured in commercial gill nets were dead upon net retrieval, yielding a mortality rate of 2.3% (Table 2).

Discussion

The results of this study provide the first quantified estimate of annual bycatch and mortality of shortnose sturgeon in the anchored gill net commercial shad fishery of the Altamaha River. Although shortnose sturgeon were captured during all three years of the study, a key finding of this study was that bycatch varied by as much as 900% across years. During the 2007 and 2008 seasons, fewer than 40 shortnose sturgeon were observed in the commercial catch, but in 2009, we recorded 105 captures yielding an expanded estimate of 498 captures over the entire three month fishery. Because of stochastic variables in habitat conditions and the protracted spawning periodicity of shortnose sturgeon, we caution against future researchers forming conclusions about sturgeon from short-term data.

The Altamaha River is thought to have the largest shortnose sturgeon population among southern rivers; however, the adult abundance is low

616 compared to that of northern river systems. Throughout the study, all but one
617 fish observed in commercial nets were adults (≥ 590 mm TL). A recent study by
618 Peterson and DeVries (2006) showed that the Altamaha population contains
619 1,500-2,000 adults, so we can estimate that in 2009 between 19 and 49 percent
620 of the adult population was "caught" in a net. In southern rivers, females spawn
621 every 3-5 years, and males every 1-2 years. We estimated that 470 (95% CI
622 278-686) adult shortnose sturgeon were captured in January and February,
623 suggesting that 25 to 80 percent of the spawning run was captured. The
624 observed mortality rate of 2.3% is lower than the 16% previously observed by
625 Collins et al. (1996) in southern shad fisheries. However, studies on sub-lethal
626 and post-release effects of bycatch are lacking. Because incidental capture of
627 spawning adults has been shown to negatively affect spawning behavior, bycatch
628 has indirect population level effects (Moser and Ross 1995; Weber 1996).

629 The highest bycatch rates occurred in the upper river strata, during the
630 month of February. In this stratum, there were never more than five fishermen
631 operating at any one time; however, many of their nets were fished in known
632 spawning areas of shortnose sturgeon. During January 2009, we observed
633 several net retrievals in this reach of the river in which 4-16 shortnose sturgeon
634 were captured in one net. In total, 36 adult shortnose sturgeon were recorded in
635 the upper river during January and February 2009, and many of the males were
636 running ripe. In contrast, no sturgeon were captured in the upper river during
637 March in any year, suggesting that the spawning period was probably limited to a
638 four to six week interval lasting from mid-January to late-February.

In all three years of the study, few shortnose sturgeon were captured in the lower river in January. Previous telemetry studies by Peterson and DeVries (2006) suggest that spawning shortnose sturgeon have already reached their spawning grounds by the start of the commercial fishing season while non-spawners remain in the estuary. Although many shortnose sturgeon were captured in the lower river during 2009, CPUE of shortnose sturgeon in the lower 184 km of the river was only 0.0015 compared to 0.5007 in the upper river during the same period. These findings suggest that spawning adult shortnose sturgeon are highly vulnerable to incidental capture in the upper 30 km of the Altamaha River.

Reducing bycatch of shortnose sturgeon in commercial fisheries is a critical component of recovering populations throughout the Atlantic coast. Further studies are needed in southern rivers, including the Altamaha, to quantify both direct (mortality) and indirect (sub-lethal and post-release) population level effects of bycatch on shortnose sturgeon populations. Although several potential management strategies already exist to minimize bycatch, the results of this study suggest that river-specific research and monitoring programs are needed to provide quantified data on the spatial and temporal variation in shortnose sturgeon movements for implementation of an effective adaptive fisheries management plan. For example, Collins et al. (2000) suggested the establishment of riverine and estuarine reserves that are completely closed to commercial gill net fisheries. Although closure of critical habitats may or may not be an important component, our results suggest that on the Altamaha River,

662 delaying the opening of commercial shad fishing in the upper river stratum until 1
663 March, would almost completely eliminate bycatch of migrating shortnose
664 sturgeon with only a minimal (5-15%) impact of total shad landings (Bahn et al.
665 2010). Regardless of which specific management actions are used, an adaptive
666 approach that incorporates real-time monitoring of commercial bycatch is the
667 only reasonable means of adequately protecting shortnose populations exposed
668 to commercial gill netting operations. Although complete closure of shad
669 fisheries is probably unnecessary, the annual variability of shortnose sturgeon
670 spawning runs and commercial fishing behavior will preclude any type of “one
671 size fits all” management approach. Consequently, future efforts to minimize
672 shortnose sturgeon bycatch while maintaining the economic and social benefits
673 provided by commercial fisheries will require close cooperation among federal
674 and state management agencies as well as commercial fishermen.

References

- Atlantic Sturgeon Status Review Team (ASSRT). 2007. Status Review of Atlantic sturgeon (*Acipenser oxyrinchus oxyrinchus*). Report to National Marine Fisheries Service, Northeast Regional Office. Washington, D. C.
- Bain, M. B. 1997. Atlantic and shortnose sturgeons of the Hudson River: common and divergent life history attributes. *Environmental Biology of Fishes* 48:347-358.
- Bahn, R. A., D. Harrison, J. Fleming, and D. L. Peterson. 2010. The anchored gill net shad fishery of the Altamaha River, Georgia. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 63:183-187.
- Bemis, W. E. and B. Kynard. 1997. Sturgeon rivers: An introduction to acipensiform biogeography and life history. *Environmental Biology of Fishes* 48:167-183.
- Boreman, J. 1997. Sensitivity of North American sturgeons and paddlefish to fishing mortality. *Environmental Biology of Fishes* 48:399-405.
- Collins, M. R. and T. I. J. Smith. 1993. Characteristics of the adult segment of the Savannah River population of shortnose sturgeon. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 47:485-491.
- Collins, M. R., S. G. Rogers, and T. I. J. Smith. 1996. Bycatch of sturgeons along the southern Atlantic Coast of the USA. *North American Journal of Fisheries Management* 16:24-29.

708 Collins, M. R., S. G. Rogers, T. I. J. Smith, and M. L. Moser. 2000. Primary
 709 factors affecting sturgeon populations in the southeastern United States:
 710 fishing mortality and degradation of essential habitats. *Bulletin of Marine*
 711 *Science* 66(3):917-928.

712 Dadswell, M. J. 1979. Biology and population characteristics of the shortnose
 713 sturgeon, *Acipenser brevirostrum* LeSueur 1818 (Osteichthes:
 714 Acipenseridae), in the Saint John River Estuary, New Brunswick, Canada.
 715 *Canadian Journal of Zoology* 57:2186-2210.

716 DeVries, R. J. 2006. Population dynamics, movements, and spawning habitat of
 717 the shortnose sturgeon, *Acipenser brevirostrum*, in the Altamaha River
 718 system, Georgia. Master's thesis. University of Georgia, Athens,
 719 Georgia.

720 Efron, B. and R. J. Tibshirani. 1994. *An Introduction to the Bootstrap*. Chapman
 721 & Hall. Boca Raton, Florida.

722 Hall, J. W., T. I. J. Smith, and S. D. Lamprecht. 1991. Movements and habitats
 723 of shortnose sturgeon, *Acipenser brevirostrum*, in the Savannah River.
 724 *Copeia* 1991:695-702.

725 Heidt, A. R. and R. J. Gilbert. 1978. The shortnose sturgeon in the Altamaha
 726 River drainage, Georgia. Pages 54-60 *in* R. R. Odum and L. Landers,
 727 editors. *Proceedings of the rare and endangered wildlife symposium*.
 728 Georgia Department of Natural Resources, Game and Fish Division,
 729 Technical Bulletin of Wildlife 4, Athens, Georgia.

730 Kynard, B. 1997. Life history, latitudinal patterns, and status of the shortnose
 731 sturgeon, *Acipenser brevirostrum*. Environmental Biology of Fishes
 732 48:319-334.

733 Malevestuto, S. P. 1996. Sampling the Recreational Creel. Pages 591-623 in
 734 B. R. Murphy and D. W. Willis, editors. Fisheries techniques, 2nd edition.
 735 American Fisheries Society, Bethesda, Maryland.

736 Moser, M. L. and S. W. Ross. 1995. Habitat use and behavior of shortnose and
 737 Atlantic sturgeons in the lower Cape Fear River, North Carolina.
 738 Transactions of the American Fisheries Society 124:225-234.

739 Munro, J., R. E. Edwards, and A. W. Kahnle. 2007. Anadromous sturgeons:
 740 habitats, threats, and management. Synthesis and summary. Pages 1-15
 741 in J. Munro, D. Hatin, J. E. Hightower, K. McKown, K. J. Sulak, A. W.
 742 Kahnle, and F. Caron, editors. Anadromous sturgeons: habitats, threats,
 743 and management. American Fisheries Society, Symposium 56, Bethesda,
 744 Maryland.

745 National Marine Fisheries Service (NMFS). 1998. Final recovery plan for the
 746 shortnose sturgeon, *Acipenser brevirostrum*. Prepared by the Shortnose
 747 Sturgeon Recovery Team for the National Marine Fisheries Service, Silver
 748 Spring, Maryland.

749 Peterson, D. L. and R. J. DeVries. 2006. Population dynamics and critical
 750 habitats of the shortnose sturgeon, *Acipenser brevirostrum*, in the
 751 Altamaha River System, Georgia. Final report to NMFS.

752 Rogers, S. G. and W. Weber. 1994. Movements of shortnose sturgeon in the
 753 Altamaha River System, Georgia. Contributions Series Number 57.
 754 Coastal Resources Division, Georgia Department of Natural Resources.
 755 Brunswick, Georgia.

756 Stein, A. B., K. D. Friedland, and M. Sutherland. 2004. Atlantic sturgeon marine
 757 bycatch and mortality on the continental shelf of the northeast United
 758 States. North American Journal of Fisheries Management 24:171-183.

759 Vladykov, V. D. and J. R. Greeley. 1963. Order Acipenseroidei. Pages 24-60 in
 760 V. H. Olsen, editor. Fishes of the western North Atlantic, part III. Memoirs
 761 of the Sears Foundation for Marine Research, Yale University, New
 762 Haven, Connecticut.

763 Weber, W. 1996. Population size and habitat use of shortnose sturgeon,
 764 *Acipenser brevirostrum*, in the Ogeechee River System, Georgia.
 765 Master's thesis. University of Georgia, Athens, Georgia.

766
 767
 768
 769
 770
 771
 772
 773
 774

Table 1. Summary data from Altamaha River shortnose sturgeon bycatch study, 2007-09.

Year	Number of net counts	Number of direct observations	Logged net hours	Percent of fishery Observed
2007	7	336	14,271	66.4
2008	11	252	15,410	59.4
2009	12	192	10,382	48.2

797 Table 2. Raw number of shortnose sturgeon captured (number dead in parentheses), CPUE, 95% CI, estimated total
 798 fishing effort (h), and estimated shortnose sturgeon bycatch (95% CI in parentheses) by river strata of the anchored gill
 799 net commercial shad fishery in the Altamaha River, Georgia, 2007 – 09. * = No data available. ** = Estimate was lower
 800 than observed value.
 801

Upper River

Year	Month	Number of SNS captured	CPUE	95% CI	Estimated total fishing effort (h)	Mean estimated bycatch (95% CI)
2007	Jan	*	*	*	1,050	*
	Feb	4	0.0126	± 0.0115	1,536	19 (4 - 37)
	Mar	0	0.0000	± 0.0000	1,185	0
2008	Jan	0	0.0000	± 0.0000	333	0
	Feb	0	0.0000	± 0.0000	612	0
	Mar	0	0.0000	± 0.0000	594	0
2009	Jan	33 (1)	0.5007	± 0.1695	666	333 (220 - 446)
	Feb	3	0.0512	± 0.0645	1,056	54 (3 - 122)
	Mar	0	0.0000	± 0.0000	495	0

Lower River

2007	Jan	13 (1)	0.0023	± 0.0013	9,744	22 (9 - 35)
	Feb	17	0.0019	± 0.0010	5,712	** **
	Mar	5 (2)	0.0021	± 0.0023	6,489	13 (5 - 28)
2008	Jan	9	0.0013	± 0.0009	7,236	9 (9 - 16)
	Feb	14	0.0031	± 0.0028	11,700	36 (14 - 69)
	Mar	5	0.0012	± 0.0012	6,930	8 (5 - 16)
2009	Jan	8	0.0015	± 0.0012	6,180	9 (8 - 16)
	Feb	47	0.0110	± 0.0042	6,720	74 (47 - 102)
	Mar	14	0.0037	± 0.0021	7,572	28 (14 - 44)


802 Table 3. Comparison of mean and associated standard errors (SE) of observed CPUE and CPUE of bootstrap resample
803 sets, 100 and 1000 bootstrap samples. * = No data available.
804

Upper River

Year	Month	Observed CPUE	SE	100 bootstrap resamples	SE	1,000 bootstrap resamples	SE
2007	Jan	*	*	*	*	*	*
	Feb	0.0126	0.00585	0.0129	0.00155	0.0131	0.00182
	Mar	0.0000	—	—	—	—	—
2008	Jan	0.0000	—	—	—	—	—
	Feb	0.0000	—	—	—	—	—
	Mar	0.0000	—	—	—	—	—
2009	Jan	0.5007	0.08650	0.5121	0.04673	0.5169	0.04778
	Feb	0.0512	0.03292	0.0550	0.01491	0.0616	0.01552
	Mar	0.0000	—	—	—	—	—

Lower River


2007	Jan	0.0023	0.00065	0.0023	0.00006	0.0023	0.00002
	Feb	0.0019	0.00053	0.0019	0.00005	0.0019	0.00005
	Mar	0.0021	0.00118	0.0022	0.00013	0.0021	0.00013
2008	Jan	0.0013	0.00045	0.0012	0.00005	0.0013	0.00005
	Feb	0.0031	0.00145	0.0032	0.00013	0.0031	0.00013
	Mar	0.0012	0.00064	0.0013	0.00007	0.0012	0.00007
2009	Jan	0.0015	0.00060	0.0017	0.00007	0.0015	0.00008
	Feb	0.0110	0.00215	0.0113	0.00021	0.0113	0.00023
	Mar	0.0037	0.00107	0.0037	0.00012	0.0037	0.00014



806

807 Figure 1. The Altamaha River with locations of commercial fishermen observed during
 808 the study. ● = Six locations and river kilometer of fishermen surveyed in each year of
 809 the study. The Seaboard Coastline Railroad Bridge (rkm 42) divides the river into two
 810 strata under current GDNr regulations. The line downstream of rkm 203 is the U.S. 1
 811 Bridge (rkm 184) which demarcates the lower and upper river strata used during this
 812 study.

813



814

815 Figure 2. Mean number of anchored gill nets with associated 95% confidence intervals
 816 observed in the Altamaha River by strata by month and year from 2007 – 09. J =
 817 January, F = February, M = March

818

819

820

821

822

823

824




Figure 3. CPUE of shortnose sturgeon with associated 95% confidence intervals in the Altamaha River by strata by month and year from 2007 – 09. J = January, F = February, M = March, * = No data

835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857

CHAPTER 3
ABUNDANCE AND RECRUITMENT OF
JUVENILE ATLANTIC STURGEON IN THE ALTAMAHA RIVER, GEORGIA³

Paul Schueller and Douglas L. Peterson*

Warnell School of Forestry and Natural Resources

University of Georgia

Athens, Georgia 30602

*Corresponding author: dpeterson@warnell.uga.edu

³ Manuscript in press, Transactions of the American Fisheries Society

858 Running Title: Juvenile Atlantic Sturgeon

859

860

861 **Abstract**

862 Juvenile Atlantic sturgeon remain in natal rivers for several years prior to out-migrating
863 to marine environments during later portions of their life history. Data regarding river-resident
864 juvenile population dynamics are unknown. During the summers of 2004 – 2007, we performed
865 mark-recapture of juvenile Atlantic sturgeon in the Altamaha River to assess age-specific
866 abundance, apparent survival, per capita recruitment, and factors influencing recruitment. The
867 objectives of this study were to estimate age-specific abundance, overall juvenile recruitment and
868 apparent survival, and to determine factors influencing recruitment. Estimates indicated that
869 juvenile abundance ranged from 1072 – 2033 individuals, and age-1 and age-2 individuals
870 comprised greater than 87% of the juvenile population, while abundance of age-3 or older
871 individuals was less than 13% of the population. Estimates of apparent survival and per capita
872 recruitment from Pradel models indicated that the juvenile population experienced high annual
873 turnover, as apparent survival rates were low ($< 33\%$) and per capita recruitment was high (from
874 0.82 to 1.38). Fall discharge, which had a positive relationship with recruitment, was the only
875 factor assessed that significantly explained time variation in per capita recruitment. The findings
876 of this study suggest that juvenile populations at the southern extreme of the Atlantic sturgeon's
877 range may remain in natal rivers for less time than northern counterparts. This is further
878 evidence of difference in life history between northern and southern populations of Atlantic
879 sturgeon. Potential findings of density dependence could have major implications for both
880 population recovery and management of this species.

881

882

883

884 **Introduction**

885 Atlantic sturgeon (*Acipenser oxyrinchus*) are a long-lived, anadromous species that spend
886 the early portion of their juvenile stage in freshwater (Scott and Crossman 1973). Adults inhabit
887 marine environments in most years, but females enter coastal rivers for spawning every 3 – 5
888 years while males spawn every 1 – 5 years (Smith 1985). In southern rivers females typically
889 spawn by age-10 and males by age-8 (Smith 1985), but age at maturity in northern populations
890 may require 20 years or more (Scott and Crossman 1973). Spawning occurs well upriver from
891 the saltwater interface of most rivers (Van Eenennaam et al. 1996, Caron et al. 2002, Hatin et al.
892 2002), as embryos and larvae are intolerant of salinity (Van Eenennaam et al. 1996). At
893 hatching, embryonic Atlantic sturgeon seek cover within interstitial spaces of rocky substrates,
894 but after 8 – 10 d they emerge as true larvae and disperse downstream (Kynard and Horgan
895 2002). Larval migration continues for approximately 12 d, and although most movements occur
896 at night during the first 6 d, little diel preference has been observed thereafter (Kynard and
897 Horgan 2002). In early juvenile development, individuals primarily use deep water habitats near
898 the fresh/saltwater interface (Moser and Ross 1995, Bain 1997). After 2 – 6 years in these
899 habitats, juveniles leave their natal rivers for marine environments (Dovel and Berggren 1983).

900 Throughout their range, Atlantic sturgeon populations have suffered declines resulting
901 from decades of anthropogenic activities. Throughout much of the 20th Century, adults were
902 harvested during spring spawning migrations for both meat and caviar (Smith 1985). As northern
903 stocks declined, commercial fishing shifted to southern rivers, particularly during the 1970s and

904 1980s (Colligan et al 1998). While overexploitation was likely a primary cause of most
905 population declines, habitat degradation may be impeding or limiting recovery of many
906 populations (Smith 1985). Degraded water quality from industrial effluents and poor land use
907 practices has adversely affected spawning and nursery habitats throughout the species' range
908 (Smith 1985, Colligan et al. 1998). Especially in southern rivers, thermal effluents and excessive
909 ground water pumping often degrades juvenile habitats by increasing water temperatures and
910 lowering dissolved oxygen (Rochard et al. 1990, Collins et al. 2000, Niklitscheck and Secor
911 2005).

912 Although Atlantic sturgeon have been federally protected since 1996 (ASMFC),
913 recovery has been difficult to assess because (1) historical abundance data are largely lacking,
914 (2) the cryptic and complex life cycle of the species makes quantitative assessments difficult, and
915 (3) latitudinal variation in ecology and population dynamics confounds direct comparisons of
916 data from northern and southern river systems. Despite uncertainties regarding recruitment
917 mechanisms and other basic aspects of juvenile ecology, long-term monitoring of juvenile
918 abundance (i.e. recruitment) is currently one of the most critical research needs for assessing
919 species recovery (Atlantic Sturgeon Status Review Team. 2007). In the Hudson River for
920 example, Peterson et al. (2000) estimated abundance of age-1 juveniles to demonstrate the
921 severity of recruitment declines resulting from decades of overfishing. Unfortunately, those
922 authors relied on the presence of hatchery-reared juveniles to estimate the abundance of wild
923 juveniles, an experimental approach which may not be appropriate or even possible on other
924 rivers systems. Furthermore, studies of recruitment mechanisms in Atlantic sturgeon have not
925 been attempted in any Atlantic coast river system.

While both scientists and managers agree that quantified methods of assessing sturgeon recruitment are essential for evaluating population trends and identifying key environmental factors that affect year class formation, early life stages of most sturgeon species are notoriously difficult to sample. In both freshwater and estuarine environments, juvenile sturgeons are widely dispersed and/or invulnerable to most types of sampling gear. Consequently, quantified estimates of abundance and mortality of juvenile sturgeons have persisted as critical information gaps in our understanding of recruitment mechanisms of sturgeon stocks worldwide (Pine et al. 2001, Secor et al. 2002; Peterson et al. 2006). Recently, however, some notable successes have been obtained using both empirical data and modeling methods. For example, Pine et al. (2001) used age-structured models to estimate first year survival in Gulf sturgeon. In a field study of lake sturgeon on the Peshtigo River, Wisconsin, Caroffino et al. (2010) sampled eggs, larvae, and age-0 juveniles to estimate first-year survival. Similar studies have been completed for a few other species, but quantified estimates of post-recruit juveniles are lacking. The Altamaha River, Georgia is currently thought to contain the 2nd largest population of Atlantic sturgeon in US waters (Peterson et al. 2008, Atlantic Sturgeon Status Review Team. 2007), but unlike the Hudson River, recruitment studies of Atlantic sturgeon have not been attempted there. The objectives of this study were to: 1) estimate annual age-specific abundance, 2) estimate annual apparent survival and per capita recruitment and 3) identify key factors that influence recruitment processes of juvenile Atlantic sturgeon in the Altamaha River.

Methods

Study Site/Fish Sampling

The study was conducted entirely within the tidally influenced portion of the Altamaha River system, near Darien, Georgia (Figure 1). To ensure spatial distribution of sampling

949 locations, specific sampling sites were randomly distributed within three contiguous 10-km strata
950 compromising the lower 30 rkm of the Altamaha Estuary. Within each stratum, channel habitats
951 deeper than 3 m were sampled weekly from June to August, 2004 – 2007. Juvenile Atlantic
952 sturgeon (Ages 1 – 3+) were captured using both trammel nets and experimental gill nets
953 measuring 91 m by 3 m. Experimental gill nets consisted of three 30.5-m panels of 7.6, 10.2,
954 and 15.2-cm monofilament mesh (stretch measure). Trammel nets were made from 7.6-cm mesh
955 inner panel and two 30.5-cm mesh outer panels. Nets were deployed perpendicular to the
956 current, anchored to the bottom, and fished for 25 – 90 min during slack tides only.

957 As nets were retrieved, juvenile Atlantic sturgeon were removed and placed in a floating
958 net pen, where they were allowed to recover for 10-15 minutes prior to data collection. Each fish
959 was then checked for PIT tags using a portable PIT tag reader. If no tag was detected, one was
960 injected beneath the fourth dorsal scute. Measurements of total length (mm) and weight (kg)
961 were then recorded for each fish. Prior to release a 0.5 – 1.0-cm section of the leading pectoral
962 fin spine was removed from a random sub-sample of 32 and 25 fish in 2005 and 2006
963 respectively for subsequent age determination.

964

965 *Data Analysis*

966 Ages of juvenile Atlantic sturgeon were determined based on modal distributions of
967 length-frequency histograms as described by Peterson et al. (2000) and subsequently, by McCord
968 et al. (2007). Accuracy of modal distribution age assignments was verified from fin spines
969 sections collected from a random sub-sample of captured juveniles. Using the basic methods
970 described by Cuerrier (1951), pectoral fin spine sections were first air dried for at least one

971 month, cross-sectioned using a Beulher Isomet[®] low-speed saw, and viewed under a dissecting
972 scope to reveal growth annuli.

973

974 *Modeling Overview*

975 The modeling approaches used to meet the objectives of the study involved the use of
976 robust design based model types. Traditional robust design models implement a combination of
977 open and closed model types (Kendall et al. 1995). Open population models, such as the
978 Cormack-Jolly-Seber model (or CJS; Cormack 1964, Jolly 1965, Seber 1965), are used between
979 primary occasions that are widely spaced, such as annual sampling, to provide estimates of
980 apparent survival. Apparent survival is defined as the probability of an individual surviving and
981 remaining in the study are during the interval from time i to time $i + 1$. Within primary occasions,
982 a series of sampling events, known as secondary occasions, are taken at shorter intervals, days or
983 a week, when the population is assumed closed, allowing the use of traditional closed population
984 abundance estimators (Otis et al. 1978). The assumptions of the traditional robust design are as
985 follows:

- 986 1. The conditional probability of surviving from primary period i to $i + 1$ is the same for
987 all fish
- 988 2. The conditional probability of being caught at each primary period is the same for all
989 marked fish
- 990 3. The fates of fish with respect to survival and capture are independent
- 991 4. Marks are retained and correctly recorded
- 992 5. Sampling periods are instantaneous, or very short, and recapture fish are released
993 immediately

994 6. All emigration is permanent

995 7. Within primary periods, the population is closed to birth, death, immigration, and
996 emigration

997 Two different modeling approaches were used to address the objectives of the study.

998 Robust design models have been modified to incorporate multi-state models among primary
999 periods, enabling the use of traditional closed capture models to estimate state specific

1000 abundance within primary periods, while allowing for state transitions between primary periods

1001 (Kendall and Bjorkland 2001, White et al. 2006). The closed robust design multi-state model

1002 type helped address the first objective by allowing us to estimate capture and recapture

1003 probabilities, determine factors influencing these probabilities, and therefore estimate state-

1004 specific abundance. The Pradel robust design model was used to estimate apparent survival, per

1005 capita recruitment, and factors influencing recruitment. Per capita recruitment was defined as

1006 the number of new juveniles in the population at time i per juvenile in the population at time $i -$

1007 1 . This is a relatively simple extension of the traditional robust design, where a Pradel model is

1008 used between primary periods rather than a CJS. Age-specific abundance estimates were not

1009 used to estimate these parameters because of potential for biased estimates. Both error in the age

1010 determination process and violations of assumptions could lead to biased age-specific abundance

1011 estimates, making them less useful than the direct estimates from the Pradel model. The

1012 assumptions of the Pradel robust design model are the same as the traditional robust design.

1013 We used a closed robust design multi-state model to estimate annual age-specific

1014 abundance and to identify factors influencing capture and recapture probabilities. Individual

1015 capture histories were constructed by using each sampling week during the summer as an

1016 individual sampling period. Eight secondary periods (4 weeks in June, and 4 weeks in July)

1017 within four primary periods (summers of 2004 – 2007) yielded a total of 32 sampling periods.
1018 Captured juveniles were first categorized into three different age strata: age-1, age-2, or age-3+.
1019 We then used the Huggins formulation of the multi-state robust design model (Huggins 1989;
1020 1991) to estimate annual abundance of each age class. The closed robust design multi-state
1021 model assumes the population is closed (i.e. no birth, death, immigration, emigration, or state
1022 transitions) within primary periods (summers), but open between primary periods. By using age
1023 as a state within the model, we were able to estimate annual abundance of each age class, while
1024 quantifying the effects of weekly sampling effort, water temperature, and river discharge on
1025 capture and recapture probabilities.

1026 A candidate set of models with different combinations of parameters for capture and
1027 recapture probabilities was constructed to identify potential differences among age-classes,
1028 behavioral responses, and to quantify influences of environmental predictor variables. Apparent
1029 survival and state transition probabilities were modeled as constant across time and ages in all
1030 models. Capture and recapture probabilities were modeled either as constant or as functions of
1031 predictor variables specific to secondary period sampling. Sampling effort was measured as
1032 number of nets set per week. Weekly means in water temperature and discharge were included as
1033 key environmental variables. Water temperature data were obtained from the Georgia Coastal
1034 Ecosystem – Long Term Ecological Research (GCE-LTER) monitoring station (~rkm 14, in
1035 South Altamaha River), while discharge data were obtained from the United States Geologic
1036 Survey (USGS) gauging station at rkm 100 (#02226000). All predictor variables were
1037 standardized, with a mean of zero and a standard deviation of one, across years before
1038 incorporation into models. The effects of predictor variables on capture and recapture
1039 probabilities were modeled as either constant or varying among summers. Behavioral response

1040 to capture (increased or decreased recapture rates after initial capture) was evaluated by
1041 including all models in the candidate set with capture and recapture probabilities set equal. To
1042 test for potential heterogeneity in capture and recapture probabilities among age classes, all
1043 models in the candidate set were rerun with separate parameters for each age class.

1044 The relative likelihood of each model was evaluated with an information theoretic
1045 approach (Burnham and Anderson 2002), by calculating Akaike's information criterion (Akaike
1046 1973) with a small sample size adjustment (AICc; Hurvich and Tsai 1989). As survival and state
1047 transition probabilities were consistent among models, assessing model likelihoods allowed us to
1048 identify sources of variation in capture and recapture probabilities. The most plausible model
1049 was then used for age-specific abundance estimates, with the corresponding parameterization of
1050 capture and recapture probabilities used in subsequent models to assess juvenile recruitment.

1051 Pradel temporal symmetry models with robust design were used to estimate parameters
1052 specific to the entire juvenile population (Kendall et al. 1995, Pradel et al. 1996). Open mark-
1053 recapture models are conditioned on first capture and use observed capture histories to estimate
1054 apparent survival and recapture probability. Reverse time models are conditioned on last
1055 observation of individuals and the reverse capture history is used to estimate the probability of an
1056 individual being in the population at a prior time (known as seniority probability) and
1057 recruitment of new individuals. Pradel temporal symmetry models use both forward and reverse
1058 time approaches simultaneously to estimate recruitment, population growth, and seniority
1059 probability (Pradel 1996). Like the closed robust design multi-state model, the Pradel robust
1060 design model also assumes the population is closed within primary periods (summers), but open
1061 between primary periods. Incorporation of Pradel models between primary periods (summers of

1062 2004 – 2007) of robust design models was used to estimate apparent survival, per capita
1063 recruitment, and juvenile population abundance.

1064 Per capita recruitment was defined as the number of new juveniles in the population at
1065 time i per juvenile in the population at time $i - 1$. Apparent survival was defined as the
1066 probability of an individual surviving and remaining in the river during the interval from time i
1067 to time $i + 1$. Apparent survival was modeled as constant or time varying. Capture and
1068 recapture probabilities were modeled using the same parameters as the best approximating closed
1069 robust design multi-state model.

1070 A candidate set of models with different combinations of recruitment parameters was
1071 constructed to evaluate the effect of various predictor variables on annual variation in juvenile
1072 recruitment. The candidate set also included models with recruitment time varying without
1073 predictor variables. Predictor variables used to explain annual variation in recruitment included
1074 spawner abundance and seasonal averages of water temperature and river discharge at time of
1075 age-0. Mean water temperature and discharge during March – May (spring), June – August
1076 (summer), and September – November (fall) were used as predictor variables because seasonal
1077 changes in flow and temperature have been previously recognized as important variables
1078 influencing Atlantic sturgeon recruitment (Secor and Gunderson 1998). Estimates of spawner
1079 abundance were derived from previous assessments of adult abundance by Peterson et al. (2008).
1080 All predictor variables were standardized among years, with a mean of zero and standard
1081 deviation of one.

1082 As in closed robust design multi-state models, the relative plausibility of each model was
1083 determined with an information theoretic approach (Burnham and Anderson 2002). Models with
1084 recruitment predictor variables were only considered important if they were more plausible than

time varying recruitment models lacking a predictor variable. As model weights were dispersed among several models, model-averaged parameter estimates were used to account for model selection uncertainty (Burnham and Anderson 2002). Model-averaged estimates and unconditional standard error were calculated for both the apparent survival and recruitment parameters and juvenile population abundance estimates.

1090

1091 **Results**

1092 In the four consecutive years of study, a total of 1,034 juvenile Atlantic sturgeon were
1093 tagged in a total of 391 net sets. A total of 86 individuals were recaptured at least once (Table
1094 1). During summer sampling, water temperature and discharge varied only slightly among years,
1095 except in 2005 when river discharge was higher and water temperature was lower. In all other
1096 years, summer water temperatures remained near 30° C and discharge varied from 70.5 to 154.6
1097 m³/s. Average number of nets set in a sampling week varied from 11.6 to 13.3 among sampling
1098 years. Catch-per-unit-effort varied from 2.04 to 3.75 juveniles per net from 2004 – 2007. Sizes
1099 of captured juveniles varied from 350 – 1050 mm total length, although 90% of juveniles
1100 measured less than 714 mm (Figure 2). While relative abundance of juvenile age-classes varied
1101 annually, the size distribution of juveniles within year classes was similar in each year of the
1102 study.

1103 Length frequency analyses of the catch identified a distinct modal distribution of
1104 juveniles. Length frequency analyses combined with age-determination from the random sub-
1105 sample of fin spines confirmed that age-1 juveniles measured 350 – 550 mm, age-2 juveniles
1106 measured 550 – 800 mm, while age-3+ juveniles measured 800 – 1050 mm (Figure 3). These
1107 results were consistent among all years of the study, except 2007 where the boundary between

1108 age-2 and age-3+ individuals was estimated to be 750 mm. After assigning ages to all juveniles
1109 captured in each year, we calculated that the total catch from 2004 to 2007 was comprised of 568
1110 age-1, 403 age-2, and 63 age-3+ juveniles (Table 2). Although annual abundance of the total
1111 juvenile population ranged from a low of 1,072 in 2004 to a high of 2,033 in 2006, ages 1-2
1112 comprised 87-96% of the juvenile population in all years of the study.

1113 Closed robust design multi-state models revealed the best-fitting model had capture and
1114 recapture probabilities equal and as a function of weekly effort varying annually (Table 3).
1115 Model comparisons showed that this model was 10.5 times more plausible than the second best
1116 model, which also had capture and recapture probabilities equal but as a function of temperature
1117 varying annually. These analyses indicated that there was no significant behavioral response to
1118 capture, and there was no evidence that capture and recapture probabilities differed among age
1119 groups.

1120 The best-fitting Pradel model indicated survival was time varying and that annual
1121 recruitment was significantly influenced by fall discharge, which had a positive relationship with
1122 recruitment (Table 4; Figure 4). In fact, this model was 1.69 times more plausible than the
1123 second best model, which had survival and recruitment time varying with no predictor variables.
1124 The third ranked model included recruitment as a function of spring Schnabel adult abundance
1125 estimates, but as this model was less likely than time varying recruitment lacking a predictor
1126 variable, it was not considered to be important. Model averaged parameters from Pradel models
1127 indicated that apparent survival and per capita recruitment estimates varied annually, with
1128 highest recruitment of 1.379 occurring in 2005 and highest apparent survival of 0.338 in the
1129 interval prior to 2006 (Table 5).

1130

1131 **Discussion**

1132 Length-frequency histograms were combined with ages determined from fin spines
1133 collected from randomly selected juveniles to estimate the ages of captured juveniles. There
1134 were some discrepancies between age determination methods. Ages determined from fin spines
1135 suggested that age-1 individuals could reach lengths of 600 mm; however, the length-frequency
1136 histograms from those years showed several distinct, non-overlapping modes. Because the
1137 modal distributions of age-1 juveniles predicted a maximum length of 550 mm for that age
1138 group, we used 550 mm as the upper limit for defining age-1 cohorts. This same approach was
1139 used by Peterson et al. (2000) who found that age-1 Atlantic sturgeon in the Hudson River were
1140 always <550 mm through the month of August (the end of our sampling season). Regardless,
1141 setting maximum size of age- cohorts in this study at 600 mm would only have changed the age
1142 assignment of a few individuals. As both approaches are subject to error, by combining length
1143 frequency analyses with fin spine collection we hoped to minimize any potential bias in our age
1144 estimates. Furthermore, average length at age-1 of Altamaha juveniles was virtually identical to
1145 that of age-1 juveniles from coastal rivers in South Carolina (McCord et al. 2007). Although
1146 these results suggest that age-estimates from length-frequency histograms and fin spines can be
1147 used to accurately identify age-1 cohorts in other southern rivers, spatial and temporal variations
1148 in growth could potentially complicate age assignment for older juveniles. Hence, future studies
1149 using known age juveniles, possibly from hatchery origin, are needed to validate age estimates of
1150 juveniles \geq age 2.

1151 Closed robust design multi-state models provided estimates of age specific juvenile
1152 abundance and identified potential sources of variation in capture probability. Model results
1153 showed that individuals of all age classes were equally likely to be captured or recaptured. The

analyses also confirmed the accuracy of the estimates by demonstrating that heterogeneity in capture probability was minimal, and hence, did not bias the abundance estimates. Consequently, we suggest that similar modeling approaches be used for other Atlantic sturgeon populations, so that results can be compared with those presented here. Provided that adequate numbers of juveniles can be captured over several consecutive years, such comparisons will greatly improve current knowledge of recruitment trends in many river systems.

The use of Pradel robust design models allowed for direct estimates of apparent survival and per capita recruitment, which together revealed a high turnover rate of the juvenile population. Apparent survival estimates were low, ranging from 0.03 to 0.34. Given that Atlantic sturgeon are a long lived species (Scott and Crossman 1973), low apparent survival values were most likely most caused by high rates of out-migration rather than true mortality. Per capita recruitment estimates in this study ranged from 0.82 to 1.38, indicating that annual recruitment to age-1 was nearly equal to, or greater than, the abundance of the entire juvenile population in the preceding year. Likewise, apparent survival was lowest when recruitment was highest, suggesting that a higher percentage of age-2 and older juveniles leave the river in years when newly recruited age-1 fish are more abundant. The surprisingly high turnover rate of river-resident juveniles observed in this study is consistent with findings of previous studies suggesting that the temporal scale of Atlantic sturgeon life history of is condensed in southern populations (Van Den Avyle 1984, Smith 1985,) compared to those of northern rivers where adults mature later and live longer (Scott and Crossman 1973, Van Eenennaam 1996). These findings also suggest that out-migration of river-resident juveniles older than age-1 may be influenced by density dependence. The source of density dependence could be competition with younger cohorts. Because early juveniles are intolerant of salinity, they are likely unable to seek

1177 alternative foraging habitats in coastal waters if riverine food resources become limited. Older
1178 juveniles, however, have no such constraints, but may prefer the relatively predator free
1179 environments of brackish water estuaries as long as food resources are not limited. To our
1180 knowledge, no research on competition among cohorts for river food sources has been
1181 researched in Atlantic sturgeon. Although further studies are needed, confirmation of density
1182 dependence in river-resident juvenile Atlantic sturgeon would have major implications for
1183 understanding ontogenetic variations in growth, survival, migration rates, and recruitment to
1184 marine life stages.

1185 Obtaining separate estimates of annual survival and out-migration rates was not possible
1186 in this study. In using the open population models to estimate apparent survival of juvenile
1187 cohorts in the Altamaha river, the requisite assumption was that emigration of juveniles was
1188 permanent (Williams et al. 2002). Consequently, apparent survival represented the probability of
1189 any individual surviving after time i and remaining in the river until time $i + 1$. As apparent
1190 survival was confounded by permanent emigration, mark-recapture methods were not capable of
1191 providing separate estimates of annual survival and out-migration, yet these rates are critical in
1192 understanding recruitment processes for the species. Future studies are needed to obtain
1193 quantified recruitment data using alternative methods such biotelemetry and known-fates
1194 modeling approaches (Cox and Oakes 1984).

1195 Although we examined the potential effects of several environmental variables, fall
1196 discharge was the only predictor variable that significantly explained annual variation in annual
1197 year class strength. The most plausible model was that with fall discharge as a predictor of
1198 recruitment, but the model with time-variation but no predictor variables also carried substantial
1199 relative weight. The fact that a model with time-variation but no predictor variables was the only

1200 other model to carry relative weight could indicate that other time varying factors not addressed
1201 in this study are important to the recruitment process. Adult abundance from the proceeding
1202 spring was the next best predictor variable, but these models were less likely than those with
1203 time varying recruitment lacking a predictor variable. Recent studies of Gulf sturgeon on the
1204 Suwannee River suggest that mean river flow during September and December may be
1205 positively related to recruitment of age-0 juveniles (Randall and Sulak 2007). The authors
1206 speculate that increased flow in fall and early winter may help increase dissolved oxygen and
1207 reduce salinity, thereby increasing potential foraging habitats available to age-0 juveniles. Given
1208 the number of hydro-generating facilities currently located on Atlantic coast rivers, future studies
1209 addressing the effects of flow on year class formation in Atlantic sturgeon should be considered
1210 as a high priority for long-term restoration of the species.

1211 The results of this study provide the first quantified recruitment data of a juvenile
1212 Atlantic sturgeon population in a southern river. Although further studies are needed to better
1213 understand recruitment mechanisms and variables affecting out-migration of river-resident
1214 juveniles, our results show that stage-based projection or population viability models can be used
1215 to assess population recovery of Atlantic sturgeon in the Altamaha and other Atlantic coast
1216 rivers. Similar approaches have been used in previous studies of other sturgeon species to
1217 project population trends (Pine et al. 2001), to identify survival bottlenecks at specific life history
1218 stages (Paragamian et al. 2005), and to quantify survival rates necessary to achieve recovery
1219 goals (Morrow et al. 1998). With regard to Atlantic sturgeon, however, current demographic
1220 data are needed to complete similar analyses. The results of this study provide quantified
1221 estimates of age-1 recruitment, apparent survival, and age-specific abundance, all of which could
1222 be used in simplified population viability analyses.

Despite the difficulties sampling juvenile sturgeons in large river systems, quantified recruitment data are essential to monitoring population recovery and to better understand the environmental variables that affect juvenile survival. Because juvenile Atlantic sturgeon remain in their natal rivers for at least 2 years after birth, quantified estimates of age-1 juveniles may offer the best opportunity to obtain these data. Similar approaches also may be possible for other sturgeon species, but the field methods employed must be developed based on a thorough understanding of specific life history traits and seasonal habitat needs. Thorough assessment of population status and recovery will require proper sampling designs and statistical approaches. Although future studies of sub-adult and adult life stages are needed, quantified assessment of river-resident juveniles can provide fisheries managers with the current data needed for evaluating population trends. Previous studies of Atlantic sturgeon on the Altamaha River have shown that population inference based on adult spawning runs can be confounded by the presence of non-spawning adults and immature fish (Peterson et al. 2008). The results of this and other studies show that sampling of river-resident juveniles, particularly the age-1 cohort, can provide reliable estimates of recruitment, a key aspect of evaluating population recovery (Bain et al. 1999, Peterson et al. 2000). The importance of monitoring juvenile populations is further supported by the finding that adult abundance does not accurately reflect variation in juvenile recruitment.

Acknowledgements

We would like to thank the Georgia Department of Natural Resources and the National Marine Fisheries Service for funding provided. We would also like to thank the many technicians and fellow graduate students who helped collect these data. Many thanks to Jim

1246 Peterson, David Higginbotham, and Justin Bezold for help with data analysis, field logistics, and
1247 editorial comments.

1248

1249

1250

1251 **References**

1252 Akaike, H. 1973. Information theory and an extension of the maximum likelihood principle.

1253 Pages 267-281. *in* B.N. Petrov and F. Csaki, editors. Second international symposium on
1254 information theory. . Akademiai Kiado, Budapest, Hungary.

1255 Atlantic Sturgeon Status Review Team. 2007. Status Review of Atlantic sturgeon (*Acipenser*
1256 *oxyrinchus oxyrinchus*). Report to National Marine Fisheries Service, northeast regional
1257 office. February 23, 2007.

1258 Bain, M.B. 1997. Atlantic and shortnose sturgeons of the Hudson River: Common and divergent
1259 life history attributes. *Environmental Biology of Fishes* 48: 347-358.

1260 Bain M. B, D. L. Peterson, K. A. Arend, and N. Haley. 1999. Atlantic sturgeon population
1261 monitoring for the Hudson River estuary: sampling design and gear recommendations.
1262 Final report to the Hudson River fisheries unit, New York Department of Environmental
1263 Conservation, New Paltz, NY and The Hudson River Foundation, New York, NY. New
1264 York Cooperative Fish and Wildlife Research Unit, Cornell University, Ithaca, NY.

1265 Burnham, K.P., and D.R. Anderson. 2002. Model selection and mulimodel inference: a practical
1266 information-theoretic approach. Springer-Verlag, New York, New York, USA.

1267 Caron, F., D. Hatin, and R. Fortin. 2002. Biological characteristics of adult Atlantic sturgeon
 1268 (*Acipenser oxyrinchus*) in the St. Lawrence River estuary and the effectiveness of
 1269 management rules. *Journal of Applied Ichthyology* 18: 580-585.

1270 Caroffino D. C., T. Sutton, R. Elliott, and M. Donofrio. 2010. Early life stage mortality rates of
 1271 lake sturgeon in the Peshtigo River, Wisconsin. *North American Journal of Fisheries*
 1272 *Management* 30:1, 295-304. Colligan, M., M. Collins, A. Hecht, M. Hendrix, A. Kahnle,
 1273 W. Laney, R. St. Pierre, R. Santos, and T. Squiers. 1998. Status review of Atlantic
 1274 sturgeon (*Acipenser oxyrinchus oxyrinchus*). Atlantic Sturgeon Status Review Team. 124.

1275 Collins, M.R., S.G. Rodgers, T.I.J. Smith, and M.L. Moser. 2000. Primary factors affecting
 1276 sturgeon populations in the southeastern United States: fishing mortality and degradation
 1277 of essential habitats. *Bulletin of Marine Sciences* 66: 917-928.

1278 Cox, D.R., and Oakes, D. 1984. *Analysis of Survival Data*. Chapman and Hall, London.

1279 Cuerrier, J.P. 1951. The use of pectoral fin rays for determining age of sturgeon and other
 1280 species of fish. *Canadian Fish Culturalist* 11: 10-18.

1281 Dovel, W.L., and T.J. Berggren. 1983. Atlantic sturgeon of the Hudson River estuary, New
 1282 York. *New York Fish and Game Journal* 30: 140-172.

1283 Hatin, D., R. Fortin, and F. Caron. 2002. Movements and aggregation areas of adult Atlantic
 1284 sturgeon (*Acipenser oxyrinchus*) in the St. Lawrence River estuary, Quebec, Canada.
 1285 *Journal of Applied Ichthyology* 18: 586-594.

1286 Huggins, R.M. 1989. On the statistical analysis of capture experiments. *Biometrika* 76:133-140.

1287 Huggins, R.M. 1991. Some practical aspects of conditional likelihood approach to capture
 1288 experiments. *Biometrics* 47:725-732.

1289 Hurvich, C.M., and C. Tsai. 1989. Regression and time series model selection in small samples.
1290 Biometrika 76: 297-307.

1291 Kendall, W.L., K.H. Pollock, and C. Brownie. 1995. A likelihood-based approach to capture-
1292 recapture estimation of demographic parameters under the robust design. Biometrics 51:
1293 293-308.

1294 Kynard, B., and M. Horgan. 2002. Ontogenetic behavior and migration of Atlantic sturgeon,
1295 *Acipenser oxyrinchus oxyrinchus*, and shortnose sturgeon, *A. brevirostrum*, with notes on
1296 social behavior. Environmental Biology of Fishes 63: 137-150.

1297 McCord, J.W., M.R. Collins, W.C. Post, and T.I.J. Smith. 2007 Attempts to develop an index of
1298 abundance for age-1 Atlantic sturgeon in South Carolina, USA. Pages 397 – 403 in J.
1299 Munro, D. Hatin, J.E. Hightower, K. McKown, K.J. Sulak, A.W. Kahnle, and F. Caron,
1300 editors. Anadromous sturgeons: habitat, threats, and management. American Fisheries
1301 Society, Symposium 56, Bethesda, MD.

1302 Morrow, J.V., J.P. Kirk, K.J. Killgore, H. Rogillio, and C. Knight. 1998. Status and recovery
1303 potential of Gulf sturgeon in the Pearl River system, Louisiana-Mississippi. North
1304 American Journal of Fisheries Management 18: 798-808.

1305 Moser, M.L., and S.W. Ross. 1995. Habitat use and movements of shortnose and Atlantic
1306 sturgeon in the lower Cape-Fear River, North Carolina. Transactions of the American
1307 Fisheries Society 124: 225-234.

1308 Niklitschek, E.J., and D.H. Secor. 2005. Modeling spatial and temporal variation of suitable
1309 nursery habitats for Atlantic sturgeon in the Chesapeake Bay. Estuarine Coastal and Shelf
1310 Science 64: 135-148.

1311 Paragamian, V.L., R. C. Beamesderfer, and S. C. Ireland. 2005. Status, population dynamics, and
 1312 future prospects of the endangered Kootenai River white sturgeon population with and
 1313 without hatchery intervention. *Transactions of the American Fisheries Society*. 134:518-
 1314 532.

1315 Peterson, D.L., M.B. Bain, and N. Haley. 2000. Evidence of declining recruitment of Atlantic
 1316 sturgeon in the Hudson River. *North American Journal of fisheries Management* 20: 231-
 1317 238.

1318 Peterson, D.L., P. Schueller, R. DeVries, J. Fleming, C. Grunwald, and I. Wirgin. 2008. Annual
 1319 run size and genetic characteristics of Atlantic sturgeon in the Altamaha River, Georgia.
 1320 *Transactions of the American Fisheries Society* 137: 393-401.

1321 Pine III, W.E., M. S. Allen, and V. J. Dreitz. 2001. Population viability of the Gulf of Mexico
 1322 sturgeon: Inferences from capture-recapture and age-structured models. *Transactions of*
 1323 *the American Fisheries Society* 130: 1164-1174.

1324 Pradel, R. 1996. Utilization of capture-mark-recapture for the study of recruitment and
 1325 population growth rates. *Biometrics* 52: 703-709.

1326 Randall, M.T., and K.J. Sulak. 2007. Relationship between recruitment of Gulf sturgeon and
 1327 water flow in the Suwannee River, Florida. Pages 69-83 *in* J. Munro, D. Hatin, J.E.
 1328 Hightower, K. McKown, K.J. Sulak, A.W. Kahnle, and F. Caron, editors. *Anadromous*
 1329 *sturgeons: habitat, threats, and management*. American Fisheries Society, Symposium 56,
 1330 Bethesda, MD.

1331 Rochard, E., G. Castelnaud, and M. Lepage. 1990. Sturgeons (Pisces: Acipenseridae); threats and
 1332 prospects. *Journal of Fish Biology* 37: 123-132.

1333 Scott, W.B., and E.J. Crossman. 1973. Freshwater fishes of Canada. Fisheries Research Board of
 1334 Canada Bulletin 184. 966 pp.

1335 Secor, D.H., and T.E. Gunderson. 1998. Effects of hypoxia and temperature on survival, growth,
 1336 and respiration of juvenile Atlantic sturgeon, *Acipenser oxyrinchus*. Fishery Bulletin 96:
 1337 603-613.

1338 Secor, D. H., P. J. Anders, W. Van Winkle, and D. Dixon. 2002. Can we study sturgeons to
 1339 extinction? What we do and don't know about the conservation of North American
 1340 sturgeons. Pages 3-10 in W. Van Winkle, P. J. Anders, D. H. Secor, and D. A. Dixon,
 1341 editors. Biology, management, and protection of North American sturgeons. American
 1342 Fisheries Society, Symposium 28, Bethesda, Maryland.

1343 Smith, T.I.J. 1985. The Fishery, biology, and management of Atlantic sturgeon, *Acipenser*
 1344 *oxyrinchus*, in North America. Environmental Biology of Fishes 14.1:61-72.

1345 William, B.K., J.D. Nichols, and M.J. Conroy. 2002. Analysis and management of animal
 1346 populations. Academic Press, New York.

1347 Van Den Avyle, M.J. 1984. Species profiles: life history and environmental requirements of
 1348 coastal fishes and invertebrates (South Atlantic) – Atlantic sturgeon. U.S. Fish Wildl. Sew.
 1349 FWS/OBS-82/11.25. U.S. Army Corps of Engineers, TR EL-82-4.

1350 Van Eenennaam, J.P., S.I. Doroshov, G.P. Moberg, J.G. Watson, D.S. Moore, and J. Linares.
 1351 1996. Reproductive conditions of the Atlantic sturgeon (*Acipenser oxyrinchus*) in the
 1352 Hudson River. Estuaries 19: 769-777.

Table 1. Number of fish tagged, number of fish recaptured, catch-per-unit-effort (CPUE), mean and range of effort (nets set per week), water temperature ($^{\circ}$ C), and discharge (m^3/s) values used to model capture probability of Atlantic sturgeon captured in the Altamaha River from June – August 2004 to 2007.

Year	Number Tagged	Number Recaptured	CPUE	Effort		Temperature		Discharge	
				Mean	Range	Mean	Range	Mean	Range
2004	174	15	2.04	11.6	3 - 21	29.8	29.1 - 30.8	154.6	80.2 - 258.3
2005	249	30	2.75	12.8	3 - 27	27.7	25.9 - 29.0	481.5	261.9 - 869.3
2006	315	18	3.72	11.3	5 - 15	30.0	28.6 - 31.5	70.5	54.3 - 90.4
2007	296	23	3.03	13.3	8 - 18	29.4	26.7 - 31.1	84.7	62.1 - 131.0

Table 2. Number of juvenile Atlantic sturgeon tagged in the Altamaha River per age class, age-specific abundance estimates from multi-state models, juvenile population abundance estimates from Pradel models, confidence intervals, and proportion of the population for 2004 to 2007.

Year	Age Class	Number Tagged	Abundance Estimate (95% CI)	Proportion of Population
2004	1	79	483 (368 – 643)	0.45
	2	89	544 (424 – 707)	0.51
	3+	6	37 (9 – 294)	0.03
	Total	174	1072 (815 – 1330)	
2005	1	226	1345 (1077 – 1697)	0.91
	2	18	107 (28 – 784)	0.07
	3+	5	30 (6 – 935)	0.02
	Total	249	1493 (1154 – 1833)	
2006	1	52	333 (246 – 460)	0.17
	2	250	1600 (1420 – 1808)	0.79
	3+	13	83 (38 – 209)	0.04
	Total	315	2033 (1582 – 2485)	
2007	1	211	1318 (1053 – 1668)	0.71
	2	46	287 (132 – 727)	0.16
	3+	39	244 (101 – 711)	0.13
	Total	296	1865 (1449 – 2282)	
Study Total	1	568		
	2	403		
	3+	63		

Table 3. Top five closed robust design multi-state models using predictor variables to describe variation in capture and recapture probability of Atlantic sturgeon in the Altamaha River for 2004 to 2007.

Capture Probability as a function of	Recapture Probability as a function of	AICc	AICc Weights	Model Likelihood	K
Weekly effort varying annually	Equal to capture probability	5251.59	0.845	1.000	7
Temperature varying annually	Equal to capture probability	5256.30	0.080	0.095	7
Weekly effort constant annually	Equal to capture probability	5258.15	0.032	0.038	4
Weekly effort varying annually	Weekly effort varying annually	5259.40	0.017	0.020	12
Weekly effort constant annually, varying by age class	Equal to capture probability	5259.75	0.014	0.017	6

Table 4. Top five Pradel robust design models using predictor variables (Fall discharge and adult abundance from two different model types, Schnabel and POPAN ;Schueller 2008) to describe variation in apparent survival and annual per capita recruitment of Atlantic sturgeon in the Altamaha River for 2004 to 2007.

Apparent Survival	Per Capita Recruitment	AICc	AICc Weights	Model Likelihood	K
Time varying	Fall discharge	8003.94	0.587	1.000	10
Time varying	Time varying	8004.99	0.347	0.592	11
Time varying	Schnabel adult abundance	8009.57	0.035	0.060	10
Constant	Time varying	8011.89	0.011	0.019	9
Time varying	POPAN adult abundance	8013.06	0.006	0.010	10
Constant	Fall discharge	8013.70	0.004	0.008	8

Table 5. Parameter estimates, and lower (LCI) and upper (UCI) 95% confidence intervals for annual apparent survival and per capita recruitment of Atlantic sturgeon in the Altamaha River for 2005 to 2007.

Parameter	Estimate	LCI	UCI
Apparent Survival '04 - '05	0.030	0.003	0.226
Apparent Survival '05 - '06	0.338	0.182	0.539
Apparent Survival '06 - '07	0.125	0.060	0.243
Per Capita Recruitment '05	1.379	1.071	1.687
Per Capita Recruitment '06	0.980	0.000	1.000
Per Capita Recruitment '07	0.823	0.609	0.933

Figure Captions

Figure 1. Netting locations (hollow triangles) and 10-km sampling strata (separated by black bars) for juvenile Atlantic sturgeon sampling within the Altamaha River, Georgia from 2004 to 2007.

Figure 2. Length (mm) frequency histogram and age assignments of all captured juvenile Atlantic sturgeon in the Altamaha River from summer sampling in 2004 to 2007.

Figure 3. Total length (mm) as a function of age, estimated from fin spines, of juvenile Atlantic sturgeon capture in the Altamaha River, Georgia.

Figure 4. Expected relationship (solid black line) and 95% confidence interval bands (dashed black line) between fall discharge and recruitment of juvenile Atlantic sturgeon based on pradel model averaged parameter estimates.




Figure 1.




Figure 2




Figure 3.




Figure 4.



Based on current regulations, areas open to commercial shad fishing in Georgia are highlighted in purple.

391-2-4-.02 Commercial Shad Fishing.

(1) **Purpose.** The purpose of these Rules is to implement the authority of the Board of Natural Resources to promulgate rules and regulations based on sound principles of wildlife research and management, establishing the seasons, days, places and methods for fishing commercially for shad.

(2) Areas Open to Commercial Shad Fishing.

(a) Nets shall be set or fished only in flowing water within the banks of the stream channels. Nets may not under any circumstances be set or fished in waters that are not flowing such as in sloughs or dead oxbow lakes.

(b) Waters of the Savannah River system open to commercial shad fishing are the Savannah River downstream of the U.S. Highway 301 bridge, Collis Creek, Albercorn Creek, Front River, Middle River, Steamboat River, McCoy's Cut, Housetown Cut, Back River upstream from Corps of Engineers New Savannah Cut, New Savannah Cut, North Channel Savannah River downstream to a line running due south of the easternmost tip of Oyster Bed Island, South Channel Savannah River downstream to a line running from the southeast tip of Cockspur Island to the mouth of Lazaretto Creek, and Elba Island Cut between North and South Channels of the Savannah River.

(c) Waters of the Ogeechee River system open to commercial shad fishing are the Ogeechee River downstream from Georgia Highway 204 bridge, Hell's Gate cut, and Ossabaw Sound upstream from the sound/beach boundary (see 391-2-4-.03) to a line running from the northwest tip of Raccoon Key across buoy R "86" to the southernmost tip of marsh adjacent to Green Island.

(d) Waters of the Altamaha River system open to commercial shad fishing are the Ochoopee River upstream to the U.S. Highway 1 bridge; the Altamaha River downstream of the from U.S. Highway 1 bridge including Cobb Creek Oxbow, Beards Creek from its mouth upstream to the Long-Tatnall County line (Big Lake), Sturgeon Hole from the Altamaha River to the lower mouth of Harper Slough, Old Woman's Pocket, South Branch, General's Cut, South Altamaha River, Champney River, Butler River, One Mile Cut, Wood Cut, Darien River upstream to the confluence Darien Creek and Cathead Creek, Buttermilk Sound upstream to the mouth of Hampton River, Hampton River, Altamaha sound to the sound/beach boundary (see 391-2-4-.03), Rockdedundy River, Little Mud River, South River, Back River, North River upstream to Hird Island Creek and Doboy Sound from the sound/beach boundary upstream to a line from range F1 R4 sec A across buoy R "178" to Sapelo Island. Old River and Mid Slough of the Penholoway River and Ellis Creek are closed to commercial shad fishing.

(e) Reserved.

(f) Reserved.

(3) **Seasons.** The commercial shad fishing season shall be open as provided in subparagraphs (a), (b) and (c) of this paragraph from 1 January to 31 March; however, the Commissioner of Natural Resources, in accordance with current, sound principles of

wildlife research and management, may at his discretion open or close the season 30 days after 31 March on any or all areas open to commercial shad fishing.

(a) The Altamaha River system downstream from the Seaboard Coastline Railroad bridge (at Altamaha Park) will be open to commercial shad fishing Monday through Friday each week. Upstream of this point will be open Tuesday through Saturday each week.

(b) The Savannah River system downstream from the I-95 bridge will be open to commercial shad fishing Tuesday through Friday each week. Upstream of the I-95 bridge it will be open Wednesday through Saturday each week.

(c) The Ogeechee River system will be open to commercial shad fishing Friday of each week.

(4) Gear and Methods for Taking Shad.

(a) Commercial Shad Fishing Gear.

1. Set nets and drift nets of at least four and one-half inch stretched mesh or trot lines (in accordance with O.C.G.A. 27-4-91) may be used to commercially fish for shad, provided, however, that only drift nets may be used in the Savannah River system downstream of a line between the mouth of Knoxboro Creek and McCoys Cut at Deadman's Point; the Ogeechee River; Altamaha Sound; and Doboy Sound.

2. Nothing in this section shall preclude the commercial use of pole and line gear as identified in O.C.G.A. 27-4-35.

(b) Methods for Taking Shad.

1. Set nets must be placed at least six hundred (600) feet apart and shall be limited to one hundred (100) feet in length. All set nets must have one end secured to the stream's bank and be buoyed at the outer (streamward) end so as to be clearly visible to boaters.

2. Set and drift nets must be situated so as to follow one-half the stream width open and free for the passage of fish.

3. Drift nets shall not be fished closer than three hundred (300) feet apart and shall be limited to a maximum of one thousand (1,000) feet in length in saltwaters.

Authority O.C.G.A. Title 27. **History.** Original Rule entitled "Commercial Shad Fishing" adopted. F. Dec. 28, 1979; eff. Jan. 17, 1980. **Amended:** F. Dec. 28, 1983; eff. Jan. 17, 1984. **Amended:** F. Dec. 2, 1987; eff. Dec. 22, 1987. **Amended:** F. June 19, 1989; eff. July 9, 1989. **Amended:** F. Dec. 9, 1994; eff. Dec. 29, 1994. **Amended:** F. Nov. 4, 2010; eff. Nov. 24, 2010.

Georgia Commercial Shad Fishery Regulation Options

The Georgia Department of Natural Resources (GA DNR) implemented new commercial shad regulations for the 2011 shad season. This action was taken in response to recent study findings that illustrated that potentially significant numbers of shortnose sturgeon could be incidentally captured in shad gill nets and the adoption of Amendment 3 to the Atlantic States Marine Fisheries Commission's (ASMFC) Interstate Fisheries Management Plan for Shad and River Herring. GA DNR utilized the best available data, results from Bahn and Peterson (2010) and GA DNR's commercial landings data, when evaluating changes to the commercial shad regulations. Bahn and Peterson's (2010) research analyzed the commercial shad set-net fishery in the Altamaha River from 2007-2009. Results from this study revealed that during 2007-2008 the bycatch rates of shortnose sturgeon in this fishery were relatively low, however, during 2009 bycatch rates of shortnose sturgeon greatly increased in the upper section of the Altamaha River. Factors, such as the periodic spawning behavior of sturgeon, location of potential spawning sites in the upper section of river, and environmental conditions (i.e. water level), may have all contributed to the increase in catch rates observed in 2009. In an attempt to reduce shortnose sturgeon bycatch in Georgia's commercial shad fishery and comply with Amendment 3 mandates, the following options were considered:

Option 1:

No change to existing commercial shad regulations. However, a status quo approach would not have provided any additional conservation measures for shortnose sturgeon nor satisfy mandates outlined in ASMFC's Amendment 3. Therefore, this option was not selected.

Option 2:

Establish new upper boundaries for commercial shad fishing on the Altamaha and Savannah rivers, while the Ogeechee, Satilla, and St. Marys rivers would have been completely closed to commercial shad fishing. It is believed that such actions would have provided adequate protection for shortnose sturgeon and satisfied Amendment 3 mandates. However, this option was not chosen due to the negative economic impacts that a total closure would have had on Ogeechee River commercial shad fishermen.

Option 3 (Preferred/Chosen Option):

Establish new upper boundaries for commercial shad fishing on the Altamaha, Ogeechee, and Savannah rivers and completely closed the Satilla and St. Marys rivers to commercial shad fishing. It is believed that these actions will provide adequate conservation measures for shortnose sturgeon and satisfied ASMFC Amendment 3 mandates.

The new upper boundary for the Altamaha River was set at the U.S. Hwy 1 bridge crossing and effectively closed commercial shad fishing on approximately 75% of the free flowing portions of the Altamaha River and its major tributaries (Ocmulgee and Oconee rivers). According to results reported by Bahn and Peterson (2010), this would decrease estimated sturgeon bycatch by up to 78% while only decreasing Altamaha River shad set-net landings by approximately 9%.

Other upper boundaries for the Altamaha River were considered (confluence of the Ochoopee River, U.S. Highway 84 bridge, and the Seaboard Coastline Railroad bridge). Utilizing 2009 creel estimates from Bahn and Peterson (2010), moving the upper boundary to one of these

lower points revealed minimal reductions in estimated shortnose sturgeon bycatch beyond those expected by setting the boundary at the U.S. Hwy 1 bridge, while having greater impacts to the commercial shad fishery. Due to the relatively small conservation advantages and larger impacts to the commercial shad fishery, GA DNR chose to set the upper commercial shad fishery boundary at U.S. Hwy 1.

No recent data on shortnose sturgeon bycatch was available for the Savannah and Ogeechee rivers. However, based on the findings from the Altamaha River it was presumed that closing the upper portions of these rivers would also likely provide greatly increased protection to shortnose sturgeon, while having relatively little impact on the commercial shad fisheries in these rivers. The upper commercial shad fishery boundary on the Savannah River was set at the U.S. Hwy 301 bridge crossing and resulted in closure of approximately 47% of the free flowing portion of the Savannah River. On the Ogeechee River, an upper commercial shad fishery boundary was established at the GA Hwy 204 bridge, which closed approximately 80% of the 245 miles of free flowing river. The number of days that the Ogeechee River remained open to commercial fishing was also reduced by 50% to one day per week and gear was limited to drift net only.

GA DNR does not have any reports off commercial shad landings on either the Satilla or St. Marys rivers since 1989. Therefore, it was concluded that entirely closing these two rivers would protect sturgeon in these two rivers and have no impact on commercial shad fishermen.