

HAWAIIAN MONK SEAL (*Neomonachus schauinslandi*)

STOCK DEFINITION AND GEOGRAPHIC RANGE

Hawaiian monk seals are distributed throughout the Northwestern Hawaiian Islands (NWHI), with subpopulations at French Frigate Shoals, Laysan Island, Lisianski Island, Pearl and Hermes Reef, Midway Atoll, Kure Atoll, and Necker and Nihoa Islands. They also occur throughout the main Hawaiian Islands (MHI). Genetic variation among monk seals is extremely low and may reflect a long-term history at low population levels and more recent human influences (Kretzmann *et al.* 1997, 2001, Schultz *et al.* 2009). Though monk seal subpopulations often exhibit asynchronous variation in demographic parameters (such as abundance trends and survival rates), they are connected by animal movement throughout the species' range (Johanos *et al.* 2013). Genetic analysis (Schultz *et al.* 2011) indicates the species is a single panmictic population. The Hawaiian monk seal is therefore considered a single stock. Scheel *et al.* (2014) established a new genus, *Neomonachus*, comprising the Caribbean and Hawaiian monk seals, based upon molecular and skull morphology evidence.

POPULATION SIZE

The best estimate of the total population size is 1,415 (95% confidence interval 1,348-1,525; CV = 0.03), (Table 1, Johanos 2018a,b,c). In 2016, new approaches were developed to estimate Hawaiian monk seal abundance, both range-wide and at individual subpopulations (Baker *et al.* 2016, Harting *et al.* 2017). In brief, methods for abundance estimation vary by site and year depending on the type and quantity of data available. Total enumeration is the favored method, but requires sufficient field presence to convincingly identify all the seals present, which is typically not achieved at most sites (Baker *et al.* 2006). When total enumeration is not possible, capture-recapture estimates (using Program CAPTURE) are conducted (Baker 2004; Otis *et al.* 1978, Rexstad & Burnham 1991, White *et al.* 1982). When no reliable estimator is obtainable in Program CAPTURE (i.e., the model selection criterion is < 0.75, following Otis *et al.* 1978), total non-pup abundance is estimated using pre-existing information on the relationship between proportion of the population identified and field effort hours expended (referred to as discovery curve analysis). At rarely visited sites (Necker, Nihoa, Niihau and Lehua Islands) where data are insufficient to use any of the above methods, beach counts are corrected for the proportion of seals at sea. In the MHI other than Niihau and Lehua Islands, abundance is estimated as the minimum tally of all individuals identified by an established sighting network during the calendar year. At all sites, pups are tallied. Finally, site-specific abundance estimates and their uncertainty are combined using Monte Carlo methods to obtain a range-wide abundance estimate distribution. All the above methods are described or referenced in Baker *et al.* (2016) and Harting *et al.* (2017). Note that because some of the abundance estimation methods utilize empirical distributions which are updated as new data accrue, previous years' estimates can change slightly when recalculated using these updated distributions.

In 2016, total enumeration was achieved only at Lisianski Island, and capture-recapture estimates were obtained for Midway Atoll and Pearl and Hermes Reef. At French Frigate Shoals, Laysan Island, and Kure Atoll abundance estimates were obtained using discovery curve analysis. Counts at Necker and Nihoa Islands are conducted from zero to a few times per year. Pups are born over the course of many months and have very different haulout patterns compared to older animals. Therefore, pup production at Necker and Nihoa Islands is estimated as the mean of the total pups observed in the past 5 years, excluding counts occurring early in the pupping season when most have yet to be born. In 2016, a single count was conducted at Necker Island and four counts were conducted at Nihoa Island.

In the MHI, NMFS collects information on seal sightings reported throughout the year by a variety of sources, including a volunteer network, the public, and directed NMFS observation effort. In recent years, a small number of surveys of Ni'ihau and nearby Lehua Islands have been conducted through a collaboration between NMFS, Ni'ihau residents and the US Navy. Total MHI monk seal abundance is estimated by adding the number of individually identifiable seals documented in 2016 on all MHI other than Ni'ihau and Lehua to an estimate for these latter two islands based on counts expanded by a haulout correction factor. A recent telemetry study (Wilson *et al.*, 2017) found that MHI monk seals (N=23) spent a greater proportion of time ashore than Harting *et al.* (2017) estimated for NWHI seals. Therefore, the total non-pup estimate for Ni'ihau and Lehua Islands was the total beach count at those sites (less individual seals already counted at other MHI) divided by the mean proportion of time hauled out in the MHI (Wilson *et al.*, 2017). The total pups observed at Ni'ihau and Lehua Islands were added to obtain the total (Table 1).

Table 1. Total and minimum estimated abundance (N_{min}) of Hawaiian monk seals by location in 2016. The estimation method is indicated for each site. Methods used include DC: discovery curve analysis, EN: total enumeration; CR: capture-recapture; CC: counts corrected for the proportion of seals at sea; Min: minimum tally. Median values are presented. Note that the median range-wide abundance is not equal to the total of the individual sites' medians, because the median of sums may differ from the sum of medians for non-symmetrical distributions. N_{min} for individual sites are either the minimum number of individuals identified or the 20th percentile of the abundance distribution (the latter applies to Necker, Nihoa, Ni'ihau/Lehua, and range-wide).

Location	Total			N_{min}			Method
	Non-pups	Pups	Total	Non-pups	Pups	Total	
French Frigate Shoals	164	35	199	162	35	197	DC
Laysan	208	31	239	208	31	239	DC
Lisianski	142	23	165	142	23	165	EN
Pearl and Hermes Reef	135	29	164	133	29	162	CR
Midway	61	12	73	59	12	71	CR
Kure	78	20	98	78	20	98	DC
Necker	63	7	70	53	7	60	CC
Nihoa	104	7	111	88	7	95	CC
MHI_(without Ni'ihau/Lehua)	124	16	140	124	16	140	Min
Ni'ihau/Lehua	122	21	143	103	21	124	CC
Range-wide	1214	201	1415	1183	201	1384	

Minimum Population Estimate

The total numbers of seals identified at the NWHI subpopulations other than Necker and Nihoa, and in the MHI other than Ni'ihau and Lehua, are the best estimates of minimum population size at those sites. Minimum population sizes for Necker, Nihoa, Ni'ihau, and Lehua Islands are estimated as the lower 20th percentiles of the non-pup abundance distributions generated using haulout corrections as described above, plus the pup estimates. The minimum abundance estimates for each site and for all sites combined (1,384) are presented in Table 1.


Current Population Trend

Range-wide abundance estimates are available from 2013 to 2016 (Figure 1). While these estimates remain somewhat negatively-biased for reasons explained in Baker et al. (2016), they provided a much more comprehensive assessment of status and trends than has been previously available. A Monte Carlo approximation of the annual multiplicative rate of realized population growth during 2013-2016 was generated by fitting 10,000 log-linear regressions to randomly selected values from each year's abundance distributions. The median rate (and 95% confidence limits) is 1.04 (1.01, 1.08). Thus, the best estimate is that the population grew at an average rate of about 4% per year from 2013 to 2016. Only 1% of the distribution was below 1, indicating that there is a 99% chance that the monk seal population increased during 2013-2016.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Mean non-pup beach counts are used as a long-term index of abundance for years when data are insufficient to estimate total abundance as described above. Prior to 1999, beach count increases of up to 7% annually were observed at Pearl and Hermes Reef, and this is the highest estimate of the maximum net productivity rate (R_{max}) observed for this species (Johanos 2017a). Consistent with this value, a life table analysis representing a time when the MHI monk seal population was apparently expanding, yielded an estimated intrinsic population growth rate of 1.07 (Baker et al. 2011).

Figure 1. Range-wide abundance of Hawaiian monk seals, 2013-2016. Medians and 95% confidence limits are shown.



POTENTIAL BIOLOGICAL REMOVAL

Past reports have concluded that Hawaiian monk seal stock dynamics did not conform to the underlying model for calculating PBR such that PBR for the Hawaiian monk seal has been undetermined. That conclusion was based on the fact that the stock was declining despite being well below OSP (Optimum Sustainable Population level). The trend since 2013 indicates the stock has recently been increasing so that PBR may be determined. Using current minimum population size (1,384), R_{max} (0.07) and a recovery factor (F_r) for ESA endangered stocks (0.1), yields a Potential Biological Removal (PBR) of 4.8.

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Human-related mortality has caused two major declines of the Hawaiian monk seal (Ragen 1999). In the 1800s, this species was decimated by sealers, crews of wrecked vessels, and guano and feather hunters (Dill and Bryan 1912; Wetmore 1925; Bailey 1952; Clapp and Woodward 1972). Following a period of at least partial recovery in the first half of the 20th century (Rice 1960), most subpopulations again declined. This second decline has not been fully explained, but long-term trends at several sites appear to have been driven both by variable oceanic productivity (represented by the Pacific Decadal Oscillation) and by human disturbance (Baker *et al.* 2012, Ragen 1999, Kenyon 1972, Gerrodette and Gilmartin 1990). Currently, human activities in the NWHI are limited and human disturbance is relatively rare, but human-seal interactions, have become an important issue in the MHI. Intentional killing of seals in the MHI is an ongoing and serious concern (Table 2).

Table 2. Intentional and potentially intentional killings of MHI monk seals, and anthropogenic mortalities not associated with fishing gear during 2012-2016 (Johanos 2018d). There were no confirmed cases in 2016; an adult female died due to trauma which may or may not have been anthropogenic.

Year	Age/sex	Island	Cause of Death	Comments
2012	Juvenile male	Kauai	Gunshot wound	
2012	Subadult male	Kauai	Skull fracture	Intent unconfirmed
2014	Adult male	Oahu	Suspected trauma	Intent unconfirmed
2014	Pup female	Kauai	Skull fracture, blunt force trauma	Likely intentional
2015	Pup male	Kauai	Dog attack/bite wounds	4 other seals injured during this event
2015	Juvenile male	Kauai	Probable boat strike	
2015	Adult male	Laysan	Research handling	Accidental, specific cause undetermined

Fishery Mortality Rate

Total fishery mortality and serious injury is not considered to be insignificant and approaching zero mortality and serious injury rate. Monk seals are being hooked and entangled in the MHI at a rate that has not been reliably assessed but is certainly greater than zero. The information above represents only reported direct interactions, and without directed observation effort, the true interaction rate cannot be estimated. Monk seals also die from entanglement in fishing gear and other debris throughout their range (likely originating from various sources outside of Hawaii), and NMFS along with partner agencies is pursuing a program to mitigate entanglement (see below).

Entanglement in Marine Debris

Hawaiian monk seals become entangled in fishing and other marine debris at rates higher than reported for other pinnipeds (Henderson 2001). Several hundred cases of marine debris entanglement with monk seals are documented (nearly all in the NWHI), including 9 mortalities (Henderson 2001; Henderson 2018b). The fishing gear fouling the reefs and beaches of the NWHI and entangling monk seals only rarely includes types used in Hawaii fisheries. For example, trawl net and monofilament gillnet accounted for approximately 35% and 34%, respectively, of the debris removed from reefs in the NWHI by weight, and trawl net alone accounted for 88% of the debris by frequency (Donohue *et al.* 2001), despite the fact that trawl fisheries have been prohibited in Hawaii since the 1980s.

The NMFS and partner agencies continue to mitigate impacts of marine debris on monk seals as well as turtles, coral reefs and other wildlife. Marine debris is removed from beaches and seals are disentangled during annual population assessment activities at the main reproductive sites. Since 1996, annual debris survey and removal efforts in the NWHI coral reef habitat have been ongoing (Donohue *et al.* 2000, Donohue *et al.* 2001, Dameron *et al.* 2007).

Other Mortality

Sources of mortality that impede recovery include food limitation (see Habitat Issues), single and multiple-male intra-species aggression (mobbing), shark predation, and disease/parasitism. Male seal aggression has caused episodes of mortality and injury. Past interventions to remove aggressive males greatly mitigated, but have not eliminated, this source of mortality (Johanos *et al.* 2010). Galapagos shark predation on monk seal pups has been a chronic and significant source of mortality at French Frigate Shoals since the late 1990s, despite mitigation efforts by NMFS (Gobush 2010). Infectious disease effects on monk seal demographic trends are low relative to other stressors. However, land-to-sea transfer of *Toxoplasma gondii*, a protozoal parasite shed in the feces of cats, is of growing concern. A case definition for toxoplasmosis and other protozoal-related mortalities was developed and retrospectively applied to 306 cases of monk seal mortality from 1982-2015 (Barbieri *et al.* 2016). Eight monk seal mortalities (and 1 suspect mortality) have been directly attributed to toxoplasmosis from 2001 to 2016. The number of mortalities from this pathogen are likely underrepresented, given that more seals disappear each year than are found dead and examined. Furthermore, *T. gondii* can be transmitted vertically from dam to fetus, and failed pregnancies are difficult to detect in wild, free-ranging animals. Unlike threats such as hook ingestion or malnutrition, which can often be mitigated through rehabilitation, options for treating seals with toxoplasmosis are severely restricted. The accumulating number of monk seal deaths from toxoplasmosis in recent years is a growing concern given the increasing geographic overlap between humans, cats, and Hawaiian monk seals in the MHI. Furthermore, the consequences of a disease outbreak introduced from livestock, feral animals, pets or other carrier wildlife may be catastrophic to the immunologically naïve monk seal population. Key disease threats include West Nile virus, morbillivirus and influenza.

Habitat Issues

Poor juvenile survival rates and variability in the relationship between weaning size and survival suggest that prey availability has limited recovery of NWHI monk seals (Baker and Thompson 2007, Baker *et al.* 2007, Baker 2008). Multiple strategies for improving juvenile survival, including translocation and captive care are being implemented (Baker and Littnan 2008, Baker *et al.* 2013, Norris 2013). The impact and effectiveness of past actions to improve monk seal survival is reflected in that approximately one-third of the monk seal population alive in 2012 was made up of seals that either had been intervened with to mitigate life-threatening situations, or were descendants of such seals (Harting *et al.* 2014). In 2014, NMFS produced a final Programmatic Environmental Impact Statement (PEIS) on current and future anticipated research and enhancement activities and issued a permit covering the activities described in PEIS preferred alternative. A major habitat issue involves loss of terrestrial habitat at French Frigate Shoals, where some pupping and resting islets have shrunk or virtually disappeared (Antonelis *et al.* 2006). Projected increases in global average sea level may further significantly reduce terrestrial habitat for monk seals in the NWHI (Baker *et al.* 2006, Reynolds *et al.* 2012).

Goodman-Lowe (1998) provided information on prey selection using hard parts in scats and spewings. Information on at-sea movement and diving is available for seals at all six main subpopulations in the NWHI using

satellite telemetry (Stewart *et al.* 2006). Cahoon (2011) and Cahoon *et al.* (2013) described diet and foraging behavior of MHI monk seals, and found no difference in prey selection between the NWHI and MHI.

The seawall at Tern Island, French Frigate Shoals, continues to degrade and poses an increasing entrapment hazard for monk seals and other fauna. The situation has worsened since 2012, when the USFWS ceased operations on Tern Island, thus leaving the island unmanaged for most of the year. Previously, daily surveys were conducted throughout the year to remove entrapped animals. Now this only occurs when NMFS monk seal field staff are on site. Furthermore, sea wall breaches are allowing sections of the island to erode and undermine buildings and other infrastructure. Several large water tanks have collapsed, exposing pipes and wiring that may entangle or entrap seals. Strategies to mitigate these threats are under consideration and there are discussions of USFWS supporting the extension of monk seal field camps to allow for entrapment mitigation beyond the regular spring/summer field season.

Monk seal juvenile survival rates are favorable in the MHI (Baker *et al.* 2011). Further, the excellent condition of pups weaned on these islands suggests that there are ample prey resources available, perhaps in part due to fishing pressure that has reduced monk seal competition with large fish predators (sharks and jacks) (Baker and Johanos 2004). Yet, there are many challenges that may limit the potential for growth in this region. The human population in the MHI is approximately 1.4 million compared to fewer than 100 in the NWHI, such that anthropogenic threats in the MHI are considerable. Intentional killing of seals is a very serious concern. Also, the same fishing pressure that may have reduced the monk seal's competitors is a source of injury and mortality. Vessel traffic in the populated islands entails risk of collision with seals and impacts from oil spills. A mortality in 2015 was deemed most likely due to boat strike. Finally, as noted above, toxoplasmosis is now recognized as a serious anthropogenic threat to seals in the MHI.

STATUS OF STOCK

In 1976, the Hawaiian monk seal was designated depleted under the Marine Mammal Protection Act of 1972 and as endangered under the Endangered Species Act of 1973. Therefore, the Hawaiian monk seal is a strategic stock. The species is well below its optimum sustainable population and has not recovered from past declines. Annual human-caused mortality for the most recent 5-year period (2012-2016) was ≥ 3.0 animals, including fishery-related mortality in nearshore gillnets and hook-and-line gear ($\geq 1.6/\text{yr}$, Table 3), intentional killings and other human-caused mortalities ($\geq 1.4/\text{yr}$, Table 2).

REFERENCES

- Antonelis, G.A., J.D. Baker, T.C. Johanos, R.C. Braun, and A.L. Harting. 2006. Hawaiian monk seal (*Monachus schauinslandi*): Status and Conservation Issues. *Atoll Res. Bull.* 543:75-101.
- Bailey, A.M. 1952. The Hawaiian monk seal. *Museum Pictorial*, Denver Museum of Natural History 7:1-32.
- Baker, J.D. 2004. Evaluation of closed capture-recapture methods to estimate abundance of Hawaiian monk seals, *Monachus schauinslandi*. *Ecol. Appl.* 14:987-998.
- Baker J.D. 2008. Variation in the relationship between offspring size and survival provides insight into causes of mortality in Hawaiian monk seals. *Endangered Species Research* 5:55-64.
- Baker J.D., A.L. Harting, and C.L. Littnan. 2013. A two-stage translocation strategy for improving juvenile survival of Hawaiian monk seals. *Endang. Species Res.* 21:33-44.
- Baker, J.D., A.L. Harting, and T.C. Johanos. 2006. Use of discovery curves to assess abundance of Hawaiian monk seals. *Mar. Mamm. Sci.* 22:847-861.
- Baker J.D., A.L. Harting, T.C. Johanos, and C.L. Littnan. 2016. Estimating Hawaiian monk seal range-wide abundance and associated uncertainty. *Endang. Species Res.* 31:317-324.
- Baker J.D., A. L. Harting, T.A. Wurth, and T.C. Johanos. 2011. Dramatic shifts in Hawaiian monk seal distribution predicted from divergent regional trends. *Mar. Mamm. Sci.* 27: 78-93.
- Baker, J.D. and T.C. Johanos. 2004. Abundance of Hawaiian monk seals in the main Hawaiian Islands. *Biol. Conserv.* 116:103-110.
- Baker J.D., and Littnan C.L. 2008. Report of the Hawaiian Monk Seal Captive Care Workshop, Honolulu, Hawaii, June 11-13, 2007. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96822-2396. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-08-02, 42 p.
- Baker J.D., E.A. Howell, and J.J. Polovina. 2012. Relative influence of climate variability and direct anthropogenic impact on a sub-tropical Pacific top predator, the Hawaiian monk seal. *Mar. Ecol. Prog. Ser.* 469:175-189.
- Baker J.D., C.L. Littnan, and D.W. Johnston. 2006. Potential effects of sea-level rise on the terrestrial habitats of endangered and endemic megafauna in the Northwestern Hawaiian Islands. *Endang. Species Res.* 4:1-10.
- Baker, J.D., J.J. Polovina, and E.A. Howell. 2007. Effect of variable oceanic productivity on the survival of an upper trophic predator, the Hawaiian monk seal, *Monachus schauinslandi*. *Mar. Ecol. Prog. Ser.* 346:277-283.

- Baker J.D. and P.M. Thompson. 2007. Temporal and spatial variation in age-specific survival rates of a long-lived mammal, the Hawaiian monk seal. *Proc. Roy. Soc. B* 274:407-415.
- Barbieri, M.M., L. Kashinsky, D.S. Rotstein, K.M. Colegrove, K.H. Haman, S.L. Magargal, A.R. Sweeny, A.C. Kaufman, M.E. Grigg, and C.L. Littnan. 2016. Protozoal-related mortalities in endangered Hawaiian monk seals *Neomonachus schauinslandi*. *Dis. Aquat. Organisms* 121:85-95.
- Cahoon, M.K. 2011. The foraging ecology of monk seals in the main Hawaiian Islands. MSc thesis, University of Hawaii, 172 p.
- Cahoon M.K., C.L. Littnan, K. Longenecker, and J.R. Carpenter. 2013. Dietary comparison of two Hawaiian monk seal populations: the role of diet as a driver of divergent population trends. *Endang. Species Res.* 20:137-146.
- Clapp, R.B., and P.W. Woodward. 1972. The natural history of Kure Atoll, Northwestern Hawaiian Islands, *Atoll Res. Bull.* 164:303-304.
- Dameron O.J., M. Park, M. Albins, and R. Brainard. 2007. Marine debris accumulation in the Northwestern Hawaiian Islands: An examination of rates and processes. *Mar. Pollut. Bull.* 54(4): 423-433.
- Dill, H.R., and W.A. Bryan. 1912. Report on an expedition to Laysan Island in 1911. U.S. Dept. of Agric. *Surv. Bull.* 42:1-30.
- Donohue, M.J., R. Brainard, M. Parke, and D. Foley. 2000. Mitigation of environmental impacts of derelict fishing gear through debris removal and environmental monitoring. *In* Hawaiian Islands Humpback Whale National Marine Sanctuary, Proceedings of the International Marine Debris Conference on Derelict Fishing Gear and the Ocean Environment, 6-11 August 2000, Honolulu, Hawaii. p. 383-402.
- Donohue, M.J., R.C. Boland, C.M. Sramek, and G.A. Antonelis. 2001. Derelict fishing gear in the Northwestern Hawaiian Islands: diving surveys and debris removal in 1999 confirm threat to coral reef ecosystems. *Mar. Pollut. Bull.* 42:1301_1312.
- Gerrodette, T.M., and W.G. Gilmartin. 1990. Demographic consequences of changed pupping and hauling sites of the Hawaiian monk seal. *Conserv. Biol.* 4:423-430.
- Gobush, K.S. 2010. Shark predation on Hawaiian monk seals: Workshop II & post-workshop developments, November 5-6, 2008. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-PIFSC-21, 43 p. + Appendices.
- Gobush, K.S., Wurth, T.A., Henderson, J. R., Becker, B. L., and C.L. Littnan. 2016. Prevalence of interactions between Hawaiian monk seals (*Neomonachus schauinslandi*) and nearshore fisheries in the main Hawaiian Islands. *Pac. Conserv. Biol.*
- Goodman-Lowe, G.D. 1998. Diet of the Hawaiian monk seal (*Monachus schauinslandi*) from the northwestern Hawaiian islands during 1991 to 1994. *Mar. Biol.* 132:535-546.
- Harting AL, Baker JD, Johanos TC. 2017. Estimating population size for Hawaiian monk seals using haulout data. *Journal of Wildlife Management.* 81:1202-9.
- Harting A.L., T.C. Johanos, and C.L. Littnan. 2014. Benefits derived from opportunistic survival-enhancing interventions for the Hawaiian monk seal: the silver BB paradigm. *Endang. Spec. Res.* 25: 89-96.
- Henderson, J.R. 1990. Recent entanglements of Hawaiian monk seals in marine debris. *In* R. S. Shomura and M. L. Godfrey (eds.), Proceedings of the Second International Conference on Marine Debris, April 2-7, 1989, Honolulu, Hawaii, p. 540-553. U.S. Dep. Commer., NOAA, Tech. Memo. NMFS-SWFSC-154.
- Henderson, J.R. 2001. A Pre and Post MARPOL Annex V Summary of Hawaiian Monk Seal Entanglements and Marine Debris Accumulation in the Northwestern Hawaiian Islands, 1982_1998. *Mar. Pollut. Bull.* 42:584-589.
- Henderson, J.R. 2018a. Hawaiian Monk Seal Research Program Hawaiian monk seal fisheries interaction data collected in the Hawaiian Archipelago, 1976-2016. US National Oceanographic Data Center.
- Henderson, J.R. 2018b. Hawaiian Monk Seal Research Program Hawaiian monk seal entanglement data collected in the Hawaiian Archipelago, 1974-2016. US National Oceanographic Data Center.
- Iverson, S., J. Piché, and W. Blanchard. 2011. Hawaiian monk seals and their prey: assessing characteristics of prey species fatty acid signatures and consequences for estimating monk seal diets using Quantitative Fatty Acid Signature Analysis. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-PIFSC-23, 114 p. + Appendices.
- Johanos, T.C. 2018a. Hawaiian Monk Seal Research Program Hawaiian monk seal survey data collected in the Hawaiian Archipelago, 1981-2016. US National Oceanographic Data Center.
- Johanos, T.C. 2018b. Hawaiian Monk Seal Research Program Hawaiian monk seal master identification records (annual) collected in the Hawaiian Archipelago, 1981-2016. US National Oceanographic Data Center.

- Johanos, T.C. 2018c. Hawaiian Monk Seal Research Program Hawaiian monk seal master identification records (seal) collected in the Hawaiian Archipelago, 1981-2016. US National Oceanographic Data Center.
- Johanos, T.C. 2018d. Hawaiian Monk Seal Research Program Hawaiian monk seal survival factors collected in the Hawaiian Archipelago, 1977-2016. US National Oceanographic Data Center.
- Johanos, T.C. and J.D. Baker (editors). 2001. The Hawaiian monk seal in the Northwestern Hawaiian Islands, 1999. U.S. Dep. Commer., NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-310, 130 p.
- Johanos T.C., B.L. Becker, J.D. Baker, T.C. Ragen, W.G. Gilmartin, and T. Gerrodette. 2010. Impacts of sex ratio reduction on male aggression in the critically endangered Hawaiian monk seal *Monachus schauinslandi*. *Endang. Species Res.* 11: 123–132.
- Johanos, T.C., A.L. Harting, T.A. Wurth, and J.D. Baker. 2013. Range-wide movement patterns of Hawaiian monk seals. *Mar. Mamm. Sci.* 30(3):1165-1174.
- Kenyon, K.W. 1972. Man versus the monk seal. *J. Mammal.* 53(4):687-696.
- Kretzmann, M.B., W.G. Gilmartin, A. Meyer, G.P. Zegers, S.R. Fain, B.F. Taylor, and D.P. Costa. 1997. Low genetic variability in the Hawaiian monk seal. *Conserv. Biol.* 11(2):482-490.
- Kretzmann, M.B., N.J. Gemmill, and A. Meyer. 2001. Microsatellite analysis of population structure in the endangered Hawaiian monk seal. *Conserv. Biol.* 15(2):457-466.
- Mercer, T.A. 2017. Hawaiian Monk Seal Research Program Hawaiian monk seal fisheries interaction data collected in the main Hawaiian Islands, 1976-2016. US National Oceanographic Data Center.
- Norris, T.A. 2013. Foraging behavior, habitat, health, and survival of resident and translocated Hawaiian monk seals at Nihoa Island, Hawaii. Msc thesis, San Jose State University, 121 p.
- Otis, D. L., K.P. Burnham, G.C. White, and D.R. Anderson. 1978. Statistical inference from capture data on closed animal populations. *Wildl. Monogr.* 62:1-135.
- Punt, A.E., and P.R. Wade. 2010. Population status of the eastern North Pacific stock of gray whales in 2009. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-AFSC-207, 43 p.
- Parrish, F.A., M.P. Craig, T.J. Ragen, G.J. Marshall, and B.M. Buhleier. 2000. Identifying diurnal foraging habitat of endangered Hawaiian monk seals using a seal-mounted video camera. *Mar. Mamm. Sci.* 16:392-412.
- Peltier, H., W. Dabin, P. Daniel, O. Van Canneyt, G. Dorémus, M. Huon, and V. Ridoux. 2012. The significance of stranding data as indicators of cetacean populations at sea: Modelling the drift of cetacean carcasses. *Ecol. Indicators* 18:278-290.
- Perrin, W.F., J.L. Thieleking, W.A. Walker, F.I. Archer, and K.M. Robertson. 2011. Common bottlenose dolphins (*Tursiops truncatus*) in California waters: Cranial differentiation of coastal and offshore ecotypes. *Mar. Mamm. Sci.* 27:769-792.
- Ragen, T.J. 1993. Status of the Hawaiian monk seal in 1992. Admin. Rep. H-93-05. Southwest Fisheries Science Center, National Marine Fisheries Service, 2570 Dole St., Honolulu, HI 96822-2396. 79 pp.
- Ragen, T.J. 1999. Human activities affecting the population trends of the Hawaiian monk seal. Pages 183-194 in J.A. Musick, ed. *Life in the slow lane: Ecology and conservation of long-lived marine animals*. American Fisheries Society Symposium 23, American Fisheries Society, Bethesda, MD.
- Reynolds, M.H., P. Berkowitz, K.N. Courtot, and C.M. Krause, eds. 2012. Predicting sea-level rise vulnerability of terrestrial habitat and wildlife of the Northwestern Hawaiian Islands: U.S. Geological Survey Open-File Report 2012-1182, 139 p.
- Rexstad, E.A., and K.P. Burnham. 1991. User's manual for interactive Program CAPTURE. Colorado Cooperative Fish and Wildlife Research Unit, Colorado State University, Fort Collins, CO. 29 pp.
- Rice, D.W. 1960. Population dynamics of the Hawaiian monk seal. *J. Mammal.* 41:376-385.
- Scheel, D.M., G.J. Slater, S.O. Kolokotronis, C.W. Potter, D.S. Rotstein, K. Tsangaras, A.D. Greenwood, and K.M. Helgen. 2014. Biogeography and taxonomy of extinct and endangered monk seals illuminated by ancient DNA and skull morphology. *ZooKeys* 409:1-33.
- Schultz J.K., Baker J.D., Toonen R.J., Bowen B.W. 2009. Extremely low genetic diversity in the endangered Hawaiian monk seal (*Monachus schauinslandi*). *J. Heredity* 100:25-33.
- Schultz J.K., J.D. Baker, R.J. Toonen, A.L. Harting, and B.W. Bowen. 2011. Range-wide genetic connectivity of the Hawaiian monk seal and implications for translocation. *Conserv. Biol.* 25:124-132.
- Stewart B.S., G.A. Antonelis, J.D. Baker, and P.Y. Yochem. 2006. Foraging biogeography of the Hawaiian monk seal in the Northwestern Hawaiian Islands. *Atoll Res Bull* 543:131-145. Wade, P.R. and R.P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Wade, P.R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Mar. Mamm. Sci.* 14:1-37.

- Wetmore, A. 1925. Bird life among lava rock and coral sand. *The Natl. Geograp. Mag.* 48:77-108.
- White, G.C., D.R. Anderson, K.P. Burnham, and L. Otis. 1982. Capture-recapture and removal methods for sampling closed populations. Los Alamos National Laboratory, Los Alamos, New Mexico.
- Williams, R., S. Gero, L. Bejder, J. Calambokidis, S.D. Kraus, D. Lusseau, A.J. Read, and J. Robbins. 2011. Underestimating the damage: interpreting cetacean carcass recoveries in the context of the Deepwater Horizon/BP incident. *Conserv. Letters* 4:228-233.
- Wilson K, Littnan C, Read AJ. 2017. Movements and home ranges of monk seals in the main Hawaiian Islands. *Marine Mammal Science.* 33:1080-96.