



Atlantic halibut – Northwestern Atlantic Coast




Blue king crab – Pribilof Islands


There has been no directed fishing since 1999 and a number of other measures have been implemented to protect this resource, but the stock has made no progress towards rebuilding. This failure to recover is likely due to environmental conditions that are unfavorable to the blue king crab's reproduction and survival rates. The current rebuilding time does not have a projected time to rebuild.



Ocean pout – Northwestern Atlantic Coast




Sandbar Shark - Atlantic and Gulf of Mexico




Thorny skate – Gulf of Maine

Overfishing occurs when the 3-year moving average of a survey index (kg/tow) declines 30% or more, or if the survey biomass declines for 3 consecutive years. A ratio < 1 represents a stock that is not subject to overfishing.




Windowpane – Gulf of Maine / Georges Bank



Winter flounder – Georges Bank

The 2018 estimates of biomass ($B/B_{MSY}=0.24$) and fishing mortality ($F/F_{MSY}=0.43$) were adjusted using Mohn's rho, to account for the retrospective pattern of overestimating biomass and underestimating fishing mortality. These adjustments were judged to be the best measure of stock size and fishing mortality and are the ratios reported in support of stock status (they are more conservative estimates).



Winter flounder – Southern New England / Mid-Atlantic

