

LONGMAN'S BEAKED WHALE (*Indopacetus pacificus*): Hawaii Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Longman's beaked whale is considered one of the least known cetacean species (Jefferson *et al.* 1993; Rice 1998; Dalebout *et al.* 2003). Until recently, it was known only from two skulls found in Australia and Somalia (Longman 1926; Azzaroli 1968). Recent genetic studies (Dalebout *et al.* 2003) have revealed that sightings of 'tropical bottlenose whales' (*Hyperoodon* sp.; Pitman *et al.* 1999) in the Indo-Pacific region were in fact Longman's beaked whales, providing the first description of the external appearance of this species. Although originally described as *Mesoplodon pacificus* (Longman 1926), it has been proposed that this species is sufficiently unique to be placed within its own genus, *Indopacetus* (Moore 1968; Dalebout *et al.* 2003).

The distribution of Longman's beaked whale, as determined from stranded specimens and sighting records of 'tropical bottlenose whales', includes tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa. A single stranding of Longman's beaked whale has been reported in Hawaii, in 2010 near Hana, Maui (West *et al.* 2012), and there was a single sighting off Kona over 13 years of nearshore surveys off in the leeward waters of the main Hawaiian Islands (Baird *et al.* 2013). Summer/fall shipboard surveys of the waters within the U.S. Exclusive Economic Zone (EEZ) of the Hawaiian Islands, resulted in one sighting in 2002 and three in 2010 (Barlow 2006, Bradford *et al.* 2017; Figure 1).

For the Marine Mammal Protection Act (MMPA) stock assessment reports, there is one Pacific stock of Longman's beaked whales, found within waters of the Hawaiian Islands EEZ. This stock includes animals found both within the Hawaiian Islands EEZ and in adjacent high seas waters; however, because data on abundance, distribution, and human-caused impacts are largely lacking for high seas waters, the status of this stock is evaluated based on data from U.S. EEZ waters of the Hawaiian Islands (NMFS 2005).

POPULATION SIZE

Encounter data from a 2010 shipboard line-transect survey of the entire Hawaiian Islands EEZ was recently evaluated using Beaufort sea-state-specific trackline detection probabilities for Longman's beaked whales, resulting in an abundance estimate of 7,619 (CV = 0.66) Longman's beaked whales (Bradford *et al.* 2017) in the Hawaii stock. A 2002 shipboard line-transect survey of the same area resulted in an abundance estimate of 1,007 (CV=1.25) Longman's beaked whales (Barlow 2006). Species abundances estimated from the 2002 HICEAS survey used pooled small dolphin, large dolphin, and large whale $g(0)$ (the probability of sighting and recording an animal directly on the track line) estimates stratified by group size (Barlow 1995). Since then, Barlow (2015) developed a more robust method for estimating species-specific $g(0)$ values that are adjusted for the Beaufort sea states that are encountered during a survey. This new method was used for analyzing the data from the 2010 survey, but has not yet been used to analyze the 2002 data.




Figure 1. Sighting locations of Longman's beaked whale during the 2002 (open diamond) and 2010 (black diamonds) shipboard cetacean surveys of U.S. waters surrounding the Hawaiian Islands (Barlow 2006, Bradford *et al.* 2017; see Appendix 2 for details on timing and location of survey effort). Outer line indicates approximate boundary of survey area and U.S. EEZ. Gray shading indicates area of Papahānaumokuākea Marine National Monument. Dotted line represents the 1000 m isobath.

Minimum Population Estimate

The minimum population size is calculated as the lower 20th percentile of the log-normal distribution (Barlow *et al.* 1995) around the 2010 abundance estimate, or 4,592 Longman's beaked whales within the Hawaiian Islands EEZ.

Current Population Trend

Abundance analyses of the 2002 and 2010 datasets used different $g(0)$ values. The 2002 survey data have not been reanalyzed using this method. This change precludes evaluation of population trends at this time. Assessment of population trend will likely require additional survey data and reanalysis of all datasets using comparable methods.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

No data are available on current or maximum net productivity rate for Longman's beaked whales.

POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) level for this stock is calculated as the minimum population size within the U.S. EEZ of the Hawaiian Islands (4,592) times one half the default maximum net growth rate for cetaceans ($\frac{1}{2}$ of 4%) times a recovery factor of 0.50 (for a stock of unknown status with no known fishery mortality or serious injury within the Hawaiian Islands EEZ; Wade and Angliss 1997), resulting in a PBR of 46 Longman's beaked whales per year.

HUMAN CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

Information on fishery-related mortality and serious injury of cetaceans in Hawaiian waters is limited, but the gear types used in Hawaiian fisheries are responsible for marine mammal mortality and serious injury in other fisheries throughout U.S. waters. No interactions between nearshore fisheries and Longman's beaked whales have been reported in Hawaiian waters. No estimates of human-caused mortality or serious injury are currently available for nearshore hook and line fisheries because these fisheries are not observed or monitored for protected species bycatch. There are currently two distinct longline fisheries based in Hawaii: a deep-set longline (DSL) fishery that targets primarily tunas, and a shallow-set longline fishery (SSL) that targets swordfish. Both fisheries operate within U.S. waters and on the high seas. Between 2011 and 2015, no Longman's beaked whales were observed hooked or entangled in the SSL fishery (100% observer coverage) or the DSL fishery (20-22% observer coverage) (Bradford 2017, Bradford and Forney 2017, McCracken 2017). However, four unidentified cetaceans, which may have been a Longman's beaked whale, were taken in the DSL fishery, and one unidentified cetacean, one unidentified Mesoplodon, and two unidentified beaked whale, which may have been Longman's beaked whales were taken in the SSL fishery.

Other Mortality

Anthropogenic sound sources, such as military sonar and seismic testing have been implicated in the mass strandings of beaked whales, including atypical events involving multiple beaked whale species (Simmonds and Lopez-Jurado 1991, Frantiz 1998, Anon. 2001, Jepson *et al.* 2003, Cox *et al.* 2006). While D'Amico *et al.* (2009) note that most mass strandings of beaked whales are unassociated with documented sonar activities, lethal or sub-lethal effects of such activities would rarely be documented, due to the remote nature of such activities and the low probability that an injured or dead beaked whale would strand. Filadelpho *et al.* (2009) reported statistically significant correlations between military sonar use and mass strandings of beaked whales in the Mediterranean and Caribbean Seas, but not in Japanese and Southern California waters, and hypothesized that regions with steep bathymetry adjacent to coastlines are more conducive to stranding events in the presence of sonar use. In Hawaiian waters, Faerber & Baird (2010) suggest that the probability of stranding is lower than in some other regions due to nearshore currents carrying animals away from beaches, and that stranded animals are less likely to be detected due to low human population density near many of Hawaii's beaches. Actual and simulated sonar are known to interrupt the foraging dives and echolocation activities of tagged beaked whales (Tyack *et al.* 2011, DeRuiter *et al.* 2013). Cuvier's beaked whales tagged and tracked during simulated mid-frequency sonar exposure showed avoidance reactions, including prolonged diving, cessation of echolocation click production associated with foraging, and directional travel away from the simulated sonar source (DeRuiter *et al.* 2013). Blainville's beaked whale presence

was monitored on hydrophone arrays before, during, and after sonar activities on a Caribbean military range, with evidence of avoidance behavior: whales were detected throughout the range prior to sonar exposure, not detected in the center of the range coincident with highest sonar use, and gradually returned to the range center after the cessation of sonar activity (Tyack *et al.* 2011). Fernández *et al.* (2013) report that there have been no mass strandings of beaked whales in the Canary Islands following a 2004 ban on sonar activities in that region. The absence of beaked whale bycatch in California drift gillnets following the introduction of acoustic pingers into the fishery implies additional sensitivity of beaked whales to anthropogenic sound (Carretta *et al.* 2008, Carretta and Barlow 2011). No estimates of potential mortality or serious injury are available for U.S. waters.

STATUS OF STOCK

The Hawaii stock of Longman's beaked whales is not considered strategic under the 1994 amendments to the MMPA. The status of Longman's beaked whales in Hawaiian waters relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. Longman's beaked whales are not listed as "threatened" or "endangered" under the Endangered Species Act (1973), nor designated as "depleted" under the MMPA. Given the absence of recent recorded fishery-related mortality or serious injuries, the total fishery mortality and serious injury can be considered to be insignificant and approaching zero. The impacts of anthropogenic sound on beaked whales remain a concern (Barlow and Gisiner 2006, Cox *et al.* 2006, Hildebrand *et al.* 2005, Weilgart 2007). The first confirmed case of *morbillivirus* in a Hawaiian cetacean was found in a subadult Longman's beaked whale stranded on Maui in 2010 (West *et al.* 2012). The presence of *morbillivirus* in all 3 known species of beaked whales in Hawaiian waters (Jacob *et al.* 2016), raises concerns about the history and prevalence of this disease in Hawaii and the potential population impacts, including cumulative impacts of disease with other stressors.

REFERENCES

- Anon. 2001. Joint Interim Report, Bahamas Marine Mammal Stranding Event of 15_16 March 2000. Available from NOAA, NMFS, Office of Protected Resources, Silver Spring, MD.
- Azzaroli, M. L. 1968. Second specimen of *Mesoplodon pacificus*, the rarest living beaked whale. *Monitore Zoologico Italiano (N.S.)* 2:67-79.
- Baird, R.W., D.L. Webster, J.M. Aschettino, G.S. Schorr, D.J. McSweeney. 2013. Odontocete cetaceans around the main Hawaiian Islands: Habitat use and relative abundance from small-boat sighting surveys. *Aquatic Mammals* 39:253-269.
- Barlow 1995. The abundance of cetaceans in California waters. Part I: ship surveys in summer and fall of 1991. *Fish.Bull.* 93:1-14.
- Barlow, J. 2006. Cetacean abundance in Hawaiian waters estimated from a summer/fall survey in 2002. *Marine Mammal Science* 22: 446-464.
- Barlow 2015. Inferring trackline detection probabilities, $g(0)$, for cetaceans from apparent densities in different survey conditions. *Mar. Mamm. Sci.* 31:923-943.
- Barlow, J. and R. Gisiner. 2006. Mitigating, monitoring, and assessing the effects of anthropogenic sound on beaked whales. *J. Cet. Res. Manage.* 7(3):239-249.
- Barlow, J., S.L. Swartz, T.C. Eagle, and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 p.
- Bradford, A.L. 2017. Injury Determinations for Marine Mammals Observed Interacting with Hawaii and American Samoa Longline Fisheries During 2015-2016. NOAA Tech Memo NMFS-PIFSC-xxx.
- Bradford, A.L. and K.A. Forney. 2017. Injury determinations for cetaceans observed interacting with Hawaii and American Samoa longline fisheries during 2010-2014. NOAA-TM-NMFS-PIFSC-62, doi:10.7289/V5/TM-PIFSC-62
- Bradford, A.L., K.A. Forney, J. E.M. Oleson, J. Barlow. 2017. Abundance estimates of cetaceans from a line-transect survey within the U.S Hawaiian Islands Exclusive Economic Zone. *Fishery Bulletin* 115: 129-142.
- Carretta, J., J. Barlow, and L. Enriquez. 2008. Acoustic pinger eliminate beaked whale bycatch in a gillnet fishery. *Marine Mammal Science* 24(4): 956-961.
- Carretta, J.V. and J. Barlow. 2011. Long-term effectiveness, failure rates, and "dinner bell" properties of acoustic pingers in a gillnet fishery. *Marine technology Society Journal* 45(5): 7-19.
- Cox, T.M., T.J. Ragen, A.J. Read, E. Vos, R.W. Baird, K. Balcomb, J. Barlow, J. Caldwell, T. Cranford, L. Crum, A. D'Amico, G. D'Spain, A. Fernandez, J. Finneran, R. Gentry, W. Gerth, F. Gulland, J.A. Hildebrand, D. Houser, T. Hullar, P.D. Jepson, D. Ketten, C.D. Macleod, P. Miller, S. Moore, D. Mountain, D. Palka, P.

- Ponganis, S. Rommel, T. Rowles, B. Taylor, P. Tyack, D. Wartzok, R. Gisiner, J. Mead, and L. Brenner. 2006. Understanding the impacts of anthropogenic sound on beaked whales. *J.Cetacean Res. Manag.* 7: 177-187.
- Dalebout, M. L., G. J. B. Ross, C. S. Baker, R. C. Anderson, P. B. Best, V. G. Cockcroft, H. L. Hinsz, V. Peddemors and R. L. Pitman. 2003. Appearance, distribution and genetic distinctiveness of Longman's beaked whale, *Indopacetus pacificus*. *Marine Mammal Science* 19:421-461.
- D'Amico A., Gisiner R.C., Ketten D.R., Hammock J.A., Johnson C., *et al.*. 2009. Beaked whale strandings and naval exercises. *Aquat. Mamm.* 34: 452-472.
- DeRuiter, S.L., Southall B.L., Calambokidis J., Zimmer W.M.X., Sadykova D., Falcone E.A., Friedlaender A.S., Joseph J.E., Moretti D., Schorr G.S., Thomas L., Tyack P.L. 2013. First direct measurements of behavioural responses by Cuvier's beaked whales to mid-frequency active sonar. *Biol Lett* 9: 20130223. <http://dx.doi.org/10.1098/rsbl.2013.0223>
- Faerber, M.M. and R.W. Baird. 2010. Does a lack of observed beaked whale strandings in military exercise areas mean no impacts have occurred? A comparison of stranding and detection probabilities in the Canary and main Hawaiian Islands. *Mar. Mamm. Sci.* 26(3); 602-613.
- Fernández, A., Arbelo, M. and Martín, V. 2013. No mass strandings since sonar ban. *Nature* 497:317.
- Filadelfo R., Mintz J., Michlovich E., D'Amico A., Tyack P.L. 2009. Correlating military sonar use with beaked whale mass strandings: what do the historical data show? *Aquat Mamm* 34: 435-444.
- Frantzis, A. 1998. Does acoustic testing strand whales? *Nature* 392(5):29.
- Hildebrand J.A. 2005. Impacts of anthropogenic sound. In: Reynolds III JE, Perrin WF, Reeves RR, Montgomery S, Ragen TJ, editors. *Marine mammal research: conservation beyond crisis*. Baltimore: Johns Hopkins University. pp. 101 – 123.
- Jacob, J.M., K.L. West, G. Levine, S. Sanchez, B.A. Jensen. 2016. Initial characterization of novel beaked whale morbillivirus in Hawaiian cetaceans. *Disease of Aquatic Organisms* 117:215-227. doi:10.3354/dao02941.
- Jefferson, T. A., S. Leatherwood, and M. A. Webber. 1993. *FAO species identification guide: marine mammals of the world*. United States Environment Programme; Food and Agriculture Organization of the United Nations (FAO), Rome. 320p.
- Jepson, P.D., M. Arbelo, R. Deaville, I. A.P. Patterson, P. Castro, J.R. Baker, E. Degollada, H.M. Ross, P. Herraiez, A.M. Pocknell, F. Rodriguez, F.E. Howie, A. Espinoza, R.J. Reid, J.R. Jaber, V. martin, A.A. Cunningham, and A. Fernandez. 2003. "Gas-bubble lesions in stranded cetaceans." *Nature* 425, no. 6958 (2003): 575-576.
- Longman, H.A. 1926. New records of Cetacea, with a list of Queensland species. *Memoirs of the Queensland Museum* 8:266-278.
- McCracken, M. 2017. Preliminary assessment of incidental interactions with marine mammals in the Hawaii longline deep and shallow set fisheries from 2011 to 2015. PIFSC Internal Report IR-17-003.
- Moore J. C. 1968. Relationships among the living genera of beaked whales. *Fieldiana Zoology* 53:209-298.
- NMFS. 2005. [Revisions to Guidelines for Assessing Marine Mammal Stocks. 24 pp.](#)
- NMFS. 2012. [NOAA Fisheries Policy Directive 02-038-01 Process for Injury Determinations \(01/27/12\).](#)
- Pitman, R. L., D. M. Palacios, P. L. Brennan, K.C. III. Balcomb, and T. Miyashita. 1999. Sightings and possible identity of a bottlenose whale in the tropical Indo-Pacific: *Indopacetus pacificus*? *Marine Mammal Science* 15:531-549.
- Rice, D. W. 1998. *Marine Mammals of the world: systematics and distribution*. Special Publication 4. The Society for Marine Mammalogy, Lawrence, KS, USA.
- Richardson, W. J., C. R. Greene, Jr., C. I. Malme, and D. H. Thompson. 1995. *Marine Mammals and Noise*. Academic Press, San Diego. 576 p.
- Simmonds, M. P., and L.F. Lopez-Jurado. 1991. Whales and the military. *Nature*, 351(6326): 448.
- Tyack, P. L., W.M.X. Zimmer, D. Moretti, B.L. Southall, D.E. Claridge, J.W. Durban, C. W. Clark, A. D'Amico, N. DiMarzio, S. Jarvis, E. McCarthy, R. Morrissey, J. Ward, and I.L. Boyd. 2011. Beaked whales respond to simulated and actual navy sonar. *PLoS One* 6(3): e17009.
- Wade, P. R. and R. P. Angliss. 1997. *Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington*. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- West, K.L., S. Sanchez, D. Rothstein, K.M. Robertson, S. Dennison, G. Levine, N. Davis, D. Schoffield, C.W. Potter, and B. Jensen. 2012. A Longman's beaked whale (*Indopacetus pacificus*) strands in Maui, Hawaii, with the first case of morbillivirus in the central Pacific. *Mar. Mamm. Sci.* DOI: 10.1111/j.1748-7692.2012.00616.x.

Weilgart, L.S. 2007. The impacts of anthropogenic ocean noise on cetaceans and implications for management. *Canadian Journal of Zoology* 85:1091-1116.