


## **MELON-HEADED WHALE (*Peponocephala electra*): Hawaiian Islands Stock Complex: Hawaiian Islands & Kohala Resident Stocks**

### **STOCK DEFINITION AND GEOGRAPHIC RANGE**


Melon-headed whales are found in tropical and warm-temperate waters throughout the world. The distribution of reported sightings suggests that the oceanic habitat of this species is primarily equatorial waters (Perryman *et al.* 1994). Small numbers have been taken in the tuna purse-seine fishery in the eastern tropical Pacific, and they are occasionally killed in direct fisheries in Japan and elsewhere in the western Pacific. Summer/fall shipboard surveys of the waters within the U.S. Exclusive Economic Zone (EEZ) of the Hawaiian Islands during 2002 and 2010 resulted in only one sighting each year (Figure 1; Barlow 2006, Bradford *et al.* 2017). Little is known about this species elsewhere in its range, and most knowledge about its biology comes from mass strandings (Perryman *et al.* 1994).

Photo-identification and telemetry studies suggest there are two demographically-independent populations of melon-headed whales in Hawaiian waters, the Hawaiian Islands stock and the Kohala resident stock. Resighting data and social network analyses of photographed individuals indicate very low rates of interchange between these populations (0.0009/yr) (Aschettino *et al.* 2012). This finding is supported by preliminary genetic analyses that suggest restricted gene flow between the Kohala residents and other melon-headed whales sampled in Hawaiian waters (Oleson *et al.* 2013). Some individuals in each population have been seen repeatedly for more than a decade, implying high site-fidelity for both populations. Individuals in the larger Hawaiian Islands stock have been resighted throughout the main Hawaiian Islands. Satellite telemetry data revealed distant offshore movements, nearly to the edge of the U.S. EEZ around the Hawaiian Islands (Figure 2), with apparent foraging near cold and warm-core eddies (Woodworth *et al.* 2012). Individuals in the smaller Kohala resident stock have a range restricted to shallower waters of the Kohala shelf and west side of Hawaii Island (Aschettino *et al.* 2012, Schorr *et al.* unpublished data). Satellite telemetry data indicate they occur in waters less than 2500m depth around the northwest and west shores of Hawaii Island, west of 156° 45' W and north of 19° 15' N (Oleson *et al.* 2013). The northern boundary between the two stocks provisionally runs through the Alenuihaha Channel between Hawaii Island and Maui, bisecting the distance between the 1000 m depth contours (Oleson *et al.* 2013).

For the Marine Mammal Protection Act (MMPA) stock assessment reports, there are two Pacific management stocks within the Hawaiian Islands EEZ (Oleson *et al.* 2013): 1) the Kohala resident stock, which includes melon-headed whales off the Kohala Peninsula and west coast of Hawaii Island and in less than 2500m of water, and 2) the Hawaiian Islands stock, which includes melon-headed whales inhabiting waters throughout the U.S. EEZ of the Hawaiian Islands, including the area of the Kohala resident stock, and adjacent high seas waters. At this time, assignment of individual melon-headed whales within the overlap area to either stock requires photographic-identification of the animal. Because data on abundance, distribution, and human-caused impacts are largely lacking for high seas waters, the status of the Hawaiian Islands stock is evaluated based on data from U.S. EEZ waters of the Hawaiian Islands (NMFS 2005).



**Figure 1.** Melon-headed whale sighting location during the 2002 (open diamond) and 2010 (black diamond) shipboard surveys of U.S. EEZ waters surrounding the Hawaiian Islands (Barlow 2006, Bradford *et al.* 2017; see Appendix 2 for details on timing and location of survey effort). Outer line represents approximate boundary of survey area and U.S. EEZ. Gray shading indicates area of Papahānaumokuākea Marine National Monument. Dotted line represents the 1000 m isobath.



**Figure 2.** Sighting locations of melon-headed whales identified as being part of the Kohala resident stock (crosses) and telemetry records of Kohala resident (dark gray triangles) and Hawaiian Islands (light gray squares) melon-headed whale stocks (Schorr *et al.*, unpublished data). The dotted line around waters adjacent to the northwest and west shores of Hawaii Island represents the provisional stock boundary for the Kohala resident stock (Oleson *et al.* 2013). The Kohala resident stock and the Hawaiian Islands stocks overlap throughout the range of the Kohala resident stock. Outer line represents U.S. EEZ. Gray shading indicates area of Papahānaumokuākea Marine National Monument.

## HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

### Fishery Information

Information on fishery-related mortality and serious injury of cetaceans in U.S. EEZ of the Hawaiian Islands waters is limited, but the gear types used in Hawaii fisheries are responsible for marine mammal mortality and serious injury in other fisheries throughout U.S. waters. Entanglement in gillnets and hooking or entanglement in various hook and line fisheries have been reported for small cetaceans in Hawaii (Nitta & Henderson, 1993). No interactions between nearshore fisheries and melon-headed whales have been reported in Hawaiian waters. No estimates of human-caused mortality or serious injury are currently available for nearshore hook and line or gillnet fisheries because these fisheries are not observed or monitored for protected species bycatch. Long-term photo-identification studies have noted individuals from both the Kohala Resident and Hawaiian Islands stocks with bullet holes in their dorsal fin or with linear scars on their fins or bodies (Aschettino 2010) which may be consistent fisheries interactions.

There are currently two distinct longline fisheries based in Hawaii: a deep-set longline (DSL) fishery that targets primarily tunas, and a shallow-set longline (SSL) fishery that targets swordfish. Both fisheries operate within U.S. waters and on the high seas. Between 2011 and 2015, no melon-headed whales were observed hooked or entangled in the SSL fishery (100% observer coverage) or the DSL fishery (20-21% observer coverage) (Bradford 2017, Bradford and Forney 2017, McCracken 2017). However, four unidentified cetaceans were taken in the DSL fishery, and one unidentified cetacean was taken in the SSL fishery, some of which may have been melon-headed whales.

### Other Mortality

In recent years, there has been increasing concern that loud underwater sounds, such as active sonar and seismic operations, may be harmful to beaked whales (Cox *et al.* 2006) and other cetaceans, including melon-headed whales (Southall *et al.* 2006) and pygmy killer whales (*Feresa attenuata*) (Wang and Yang 2006). The use of active sonar from military vessels has been implicated in mass strandings of beaked whales and recent mass-stranding reports suggest some delphinids may be impacted as well. A 2004 mass-stranding of 150-200 melon-headed whales in Hanalei Bay, Kauai occurred during a multi-national sonar training event around Hawaii (Southall *et al.* 2006). Although data limitations regarding the position of the whales prior to their arrival in the Bay, the

magnitude of sonar exposure, behavioral responses of melon-headed whales to acoustic stimuli, and other possible relevant factors preclude a conclusive finding regarding the role of Navy sonar in triggering this event, sonar transmissions were considered a plausible cause of the mass stranding based on the spatiotemporal link between the sonar exercises and the stranding, the direction of movement of the transmitting vessels near Hanalei Bay, and propagation modeling suggesting the sonar transmissions would have been audible at the mouth of Hanalei Bay (Southall *et al.* 2006; Brownell *et al.* 2009). In 2008 approximately 100 melon-headed whales stranded within a lagoon off Madagascar during high-frequency multi-beam sonar use by oil and gas companies surveying offshore. Although the multi-beam sonar cannot be conclusively deemed the cause of the stranding event, the very close temporal and spatial association and directed movement of the sonar use with the stranding event, the unusual nature of the stranding event, and that all other potential causal factors were considered unlikely to have contributed, an Independent Scientific Review panel found that multi-beam sonar transmissions were a “plausible, if not likely” contributing factor (Southall *et al.* 2013) in this mass stranding event. This examination together with that of Brownell *et al.* (2009) suggests melon-headed whale may be particularly sensitive to impacts from anthropogenic sounds. No estimates of potential mortality or serious injury are available for U.S. waters.

## **KOHALA RESIDENT STOCK**

### **POPULATION SIZE**

Using the photo-ID catalog of individuals encountered between 2002 and 2009, Achettino (2010) used a POPAN open-population model to produce a mark-recapture abundance estimate of 447 (CV=0.12) individuals. A portion of the data used in that analysis is more than 8 years old; however, full sighting histories were required to produce a valid model for mark-recapture analyses, such that an estimate restricted to only the later years of the period is not available. Although this estimate includes individuals that have died since 2002 it is currently the best available abundance estimate for the resident stock.

### **Minimum Population Estimate**

The minimum population size is calculated as the lower 20th percentile of the log-normal distribution (Barlow *et al.* 1995) around the 2002-2009 mark-recapture abundance estimate (Achettino 2010), or 404 melon-headed whales in the Kohala resident stock.

### **Current Population Trend**

Photographic mark-recapture data will be evaluated in the future to assess whether sufficient data exists to assess trends.

## **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

No data are available on current or maximum net productivity rate.

## **POTENTIAL BIOLOGICAL REMOVAL**

The potential biological removal (PBR) level for this stock is calculated as the minimum population estimate (404) times one half the default maximum net growth rate for cetaceans ( $\frac{1}{2}$  of 4%) times a recovery factor of 0.50 (for a species of unknown status with no known fishery mortality; Wade and Angliss 1997), resulting in a PBR of 4.0 Kohala resident melon-headed whales per year.

## **STATUS OF STOCK**

The Kohala resident stock of melon-headed whales is not considered strategic under the MMPA. The status of this stock relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. Melon-headed whales are not listed as “threatened” or “endangered” under the Endangered Species Act (1973), nor designated as “depleted” under the MMPA. There have been no reports of recent mortality or serious injuries; however, there is no systematic monitoring of takes in near-shore fisheries that may take this species. Given noted bullet holes and potential line injuries on individuals from this stock, insufficient information is available to determine whether the total fishery mortality and serious injury for Kohala Resident melon-headed whales is insignificant and approaching zero mortality and serious injury rate. The very restricted range and small population size of Hawaii Island resident melon-headed whales suggests this population may be at risk due to its proximity to U.S. Navy training, including sonar transmissions, in the Alenuihaha Channel between Hawaii Island and Maui (Anonymous 2006). Although a 2004 mass-stranding in Hanalei Bay, Kauai could not be conclusively linked to

Naval training events in the region (Southall *et al.* 2006), the spatiotemporal link between sonar exercises and the stranding does raise concern on the potential impact on the Kohala Resident population due to of sonar training nearby.

## **HAWAIIAN ISLANDS STOCK**

### **POPULATION SIZE**

Encounter data from a 2010 shipboard line-transect survey of the entire Hawaiian Islands EEZ was recently evaluated using Beaufort sea-state-specific trackline detection probabilities for bottlenose dolphins, resulting in an abundance estimate of 8,666 (CV = 1.00) melon headed whales (Bradford *et al.* 2017) in the Hawaiian Islands stock. Using the photo-ID catalog of individuals encountered between 2002 and 2009 near the main Hawaiian Islands, Achettino (2010) used a POPAN open-population model to produce a mark-recapture abundance estimate of 5,794 (CV=0.20) individuals. A 2002 shipboard line-transect survey of the Hawaiian EEZ resulted in an abundance estimate of 2,950 (CV=1.17) melon-headed whales (Barlow 2006). Species abundances estimated from the 2002 HICEAS survey used pooled small dolphin, large dolphin, and large whale  $g(0)$  (the probability of sighting and recording an animal directly on the track line) estimates stratified by group size (Barlow 1995). Since then, Barlow (2015) developed a more robust method for estimating species-specific  $g(0)$  values that are adjusted for the Beaufort sea states that are encountered during a survey. This new method was used for analyzing the data from the 2010 survey, but has not yet been used to analyze the 2002 data. An abundance estimate of melon-headed whales is available for the eastern tropical Pacific (Wade and Gerrodette 1993), but it is not known whether any of these animals are part of the same population that occurs around the Hawaiian Islands.

### **Minimum Population Estimate**

The minimum population size is calculated as the lower 20th percentile of the log-normal distribution (Barlow *et al.* 1995) of the 2010 line-transect abundance estimate (Bradford *et al.* 2017) or 4,299 melon-headed whales. This log-normal 20<sup>th</sup> percentile minimum population size is similar to the log-normal 20<sup>th</sup> percentile mark-recapture estimate (4,904) from Aschettino (2010).

### **Current Population Trend**

Abundance analyses of the 2002 and 2010 datasets used different  $g(0)$  values. The 2002 survey data have not been reanalyzed using this method. This change precludes evaluation of population trends at this time. Assessment of population trend will likely require additional survey data and reanalysis of all datasets using comparable methods.

## **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

No data are available on current or maximum net productivity rate.

## **POTENTIAL BIOLOGICAL REMOVAL**

The potential biological removal (PBR) level for this stock is calculated as the minimum population estimate for the U.S. EEZ of the Hawaiian Islands (4,299) times one half the default maximum net growth rate for cetaceans ( $\frac{1}{2}$  of 4%) times a recovery factor of 0.50 (for a species of unknown status with no known fishery mortality; Wade and Angliss 1997), resulting in a PBR of 43 melon-headed whales per year.

## **STATUS OF STOCK**

The Hawaiian Islands stock of melon-headed whales is not considered strategic under the 1994 amendments to the MMPA. The status of this stock relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. Melon-headed whales are not listed as “threatened” or “endangered” under the Endangered Species Act (1973), nor designated as “depleted” under the MMPA. There have been no reports of recent mortality or serious injuries; however, there is no systematic monitoring of takes in near-shore fisheries that may take this species. Given noted bullet holes and potential line injuries on individuals from this stock, insufficient information is available to determine whether the total fishery mortality and serious injury for Hawaiian Islands melon-headed whales is insignificant and approaching zero mortality and serious injury rate. A 2004 mass-stranding of melon-headed whales in Hanalei Bay, Kauai occurred during a multi-national sonar training event around Hawaii (Southall *et al.* 2006). Although the event could not be conclusively linked to Naval training events in the region (Southall *et al.* 2006), the spatiotemporal link between sonar exercises and the stranding does raise concern on the

potential impact on the Hawaiian Islands population due to its frequent use of nearshore areas within the main Hawaiian Islands.

## REFERENCES

- Anonymous. 2006. Supplement to the 2002 Rim of the Pacific (RIMPAC) Programmatic Environmental Assessment.
- Aschettino, J.M. 2010. Population size and structure of melon-headed whales (*Peponocephala electra*) around the main Hawaiian Islands: evidence of multiple populations based on photographic data. Master's Thesis. Hawaii Pacific University.
- Aschettino, J.M., R.W. Baird, D.J. McSweeney, D.L. Webster, G.S. Schorr, J.L. Huggins, K.K. Martien, S.D. Mahaffy, and K.L. West. 2012. Population structure of melon-headed whales (*Peponocephala electra*) in the Hawaiian Archipelago: evidence of multiple populations based on photo identification. *Marine Mammal Science* 28(4): 666-689. doi: 10.1111/j.1748-7692.2011.00517.x
- Barlow 1995. The abundance of cetaceans in California waters. Part I: ship surveys in summer and fall of 1991. *Fish.Bull.* 93:1-14.
- Barlow, J. 2006. Cetacean abundance in Hawaiian waters estimated from a summer/fall survey in 2002. *Marine Mammal Science* 22(2): 446-464.
- Barlow 2015. Inferring trackline detection probabilities,  $g(0)$ , for cetaceans from apparent densities in different survey conditions. *Mar. Mamm. Sci.* 31:923-943.
- Barlow, J., S.L. Swartz, T.C. Eagle, and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 p.
- Bradford, A.L. 2017. Injury Determinations for Marine Mammals Observed Interacting with Hawaii and American Samoa Longline Fisheries During 2015-2016. NOAA Tech Memo NMFS-PIFSC-xxx.
- Bradford, A.L. and K.A. Forney. 2017. Injury determinations for cetaceans observed interacting with Hawaii and American Samoa longline fisheries during 2010-2014. NOAA-TM-NMFS-PIFSC-62, doi:10.7289/V5/TM-PIFSC-62
- Bradford, A.L., K.A. Forney, J. E.M. Oleson, J. Barlow. 2017. Abundance estimates of cetaceans from a line-transect survey within the U.S Hawaiian Islands Exclusive Economic Zone. *Fishery Bulletin* 115: 129-142.
- Brownell, R.L., Jr., K. Ralls, S. Baumann-Pickering, and M.M. Poole. 2009. Behavior of melon-headed whales, *Peponocephala electra*, near oceanic islands. *Marine Mammal Science* 25(3):639-658.
- Cox, T.M., T.J. Ragen, A.J. Read, E. Vos, R.W. Baird, K. Balcomb, J. Barlow, J. Caldwell, T. Cranford, L. Crum, A. D'Amico, G. D'Spain, A. Fernandez, J. Finneran, R. Gentry, W. Gerth, F. Gulland, J.A. Hildebrand, D. Houser, T. Hullar, P.D. Jepson, D. Ketten, C.D. Macleod, P. Miller, S. Moore, D. Mountain, D. Palka, P. Ponganis, S. Rommel, T. Rowles, B. Taylor, P. Tyack, D. Wartzok, R. Gisiner, J. Mead, and L. Brenner. 2006. Understanding the impacts of anthropogenic sound on beaked whales. *J.Cetacean Res. Manag.* 7: 177-187.
- Kobayashi, D. R. and K. E. Kawamoto. 1995. Evaluation of shark, dolphin, and monk seal interactions with Northwestern Hawaiian Island bottomfishing activity: a comparison of two time periods and an estimate of economic impacts. *Fisheries Research* 23:11-22.
- Maldini, D., L. Mazzuca, and S. Atkinson. 2005. Odontocete stranding patterns in the Main Hawaiian Islands (1937-2002): How do they compare with live animal surveys? *Pacific Science* 59(1):55-67.
- McCracken, M. 2017. Preliminary assessment of incidental interactions with marine mammals in the Hawaii longline deep and shallow set fisheries from 2011 to 2015. PIFSC Internal Report IR-17-003.
- Nishiwaki, M. and K. S. Norris. 1966. A new genus, *Peponocephala*, for the odontocete cetacean species *Electra electra*. *Sci. Rep. Whales Res. Inst.* 20:95-100.
- Nitta, E. 1991. The marine mammal stranding network for Hawaii: an overview. *In*: J.E. Reynolds III, D.K. Odell (eds.), *Marine Mammal Strandings in the United States*, pp.56-62. NOAA Tech. Rep. NMFS 98, 157 pp.
- Nitta, E. and J. R. Henderson. 1993. A review of interactions between Hawaii's fisheries and protected species. *Mar. Fish. Rev.* 55(2):83-92.
- NMFS, Pacific Islands Region, Observer Program, 1602 Kapiolani Blvd, Suite 1110, Honolulu, HI 96814.
- NMFS. 2005. [Revisions to Guidelines for Assessing Marine Mammal Stocks. 24 pp.](#)
- NMFS. 2012. [NOAA Fisheries Policy Directive 02-038-01 Process for Injury Determinations \(01/27/12\).](#)
- Oleson, E.M., R.W. Baird, K.K. Martien, and B.L. Taylor. 2013. Island-associated stocks of odontocetes in the main Hawaiian Islands: A synthesis of available information to facilitate evaluation of stock structure. PIFSC Working Paper WP-13-003. Perrin, W.F., G. P. Donovan and J. Barlow. 1994. Gillnets and Cetaceans. Rep.

- Int. Whal. Commn., Special Issue 15, 629 pp.
- Perryman, W. L., D. W. K. Au, S. Leatherwood, and T. A. Jefferson. 1994. Melon-headed whale *Peponocephala electra* Gray, 1846. In: S. H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Vol.5: The First Book of Dolphins, pp. 363-386. Academic Press, 416 pp.
- Shallenberger, E.W. 1981. The status of Hawaiian cetaceans. Final report to U.S. Marine Mammal Commission. MMC-77/23, 79pp.
- Southall, B. R. Braun, F.M.D. Gulland, A.D. Heard, R.W. Baird, S.M. Wilkin, T.K. Rowles. 2006. Hawaiian melon-headed whale (*Peponocephala electra*) mass stranding event of July 3-4, 2004. NOAA Technical Memorandum NMFS-OPR-31. 73 pp.
- Southall, B.L., T. Rowles, F. Gulland, R.W. Baird, and P.D. Jepson. 2013. Final report of the Independent Scientific Review Panel investigating potential contributing factors to a 2008 mass stranding of melon-headed whales (*Peponocephala electra*) in Antsohihy, Madagascar. Report to the U.S. Marine Mammal Commission, 75 pp.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Wade, P. R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. Rep. Int. Whal. Commn. 43:477-493.
- Wang, J.Y., and S.-C. Yang. 2006. Unusual cetacean stranding events in Chinese waters in early 2004 and 2005. J. Cetacean Res. and Manag. 8:283-292.
- Woodworth, P.A., G.S. Schorr, R.W. Baird, D.J. McSweeney, M.B. Hanson, R.D. Andrews, and J.J. Polovina. 2012. Eddies as offshore foraging grounds for melon-headed whales (*Peponocephala electra*). Marine Mammal Science 28(3); 638-647.