


## SPINNER DOLPHIN (*Stenella longirostris*): Hawaiian Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

Spinner dolphins are found throughout the world in tropical and warm-temperate waters (Perrin and Gilpatrick 1994). They are common and abundant throughout the entire Hawaiian archipelago (Shallenberger 1981; Norris and Dohl 1980; Norris et al. 1994). Recent sighting locations around the main Hawaiian Islands (Mobley et al. 2000) are shown in Figure 1. There is some suggestion from an intensive study of spinner dolphins off the Kona Coast of Hawaii that the waters surrounding this island may have a large, relatively stable "resident" population (Norris et al. 1994).

Hawaiian spinner dolphins belong to a stock that is separate from those involved in the tuna purse-seine fishery in the eastern tropical Pacific (Perrin 1975; Dizon et al. 1994). The Hawaiian form is referable to the subspecies *S. longirostris longirostris*, which occurs pantropically (Perrin 1990). For the Marine Mammal Protection Act (MMPA) stock assessment reports, there is a single Pacific management stock including only animals found within the U.S. Exclusive Economic Zone of the Hawaiian Islands. Spinner dolphins involved in eastern tropical Pacific tuna purse-seine fisheries are managed separately under the MMPA.



**Figure 1.** Spinner dolphin sighting locations during 1993-98 aerial surveys within about 25 nmi of the main Hawaiian Islands (see Appendix 2 for details on timing and location of survey effort). Outer line indicates approximate boundary of survey area.

### POPULATION SIZE

Although spinner dolphins are clearly among the most abundant cetaceans in Hawaiian waters, previously available population estimates apply only to the west coast of Hawaii. Norris et al. (1994) photoidentified 192 individuals along the west coast of Hawaii and estimated 960 animals for this area in 1979-1980. Östman (1994) photoidentified 677 individual spinner dolphins in the same area from 1989 to 1992. Using the same estimation procedures as Norris et al. (1994), Östman (1994) estimated a population size of 2,334 for his study area along the Kona coast of Hawaii. As part of the Marine Mammal Research Program of the Acoustic Thermometry of Ocean Climate (ATOC) study, a total of twelve aerial surveys were conducted within about 25 nmi of the main Hawaiian Islands in 1993, 1995 and 1998. An abundance estimate of 3,184 (CV=0.37) spinner dolphins was recently calculated from the combined survey data (Mobley et al. 2000). This abundance underestimates the total number of spinner dolphins within the U.S. EEZ off Hawaii, because areas around the Northwest Hawaiian Islands (NWHI) and beyond 25 nautical miles from the main islands were not surveyed.

#### Minimum Population Estimate

The log-normal 20th percentile of the combined 1993-98 abundance estimate is 2,355 spinner dolphins. As with the best abundance estimate above, this includes only areas within about 25 nmi of the main Hawaiian Islands and is therefore an underestimate.

#### Current Population Trend

No data on current population trend are available.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

No information on current or maximum net productivity rate is currently available for the Hawaiian stock.

## POTENTIAL BIOLOGICAL REMOVAL


The potential biological removal (PBR) level for this stock is calculated as the minimum population size (2,355) times one half the default maximum net growth rate for cetaceans ( $\frac{1}{2}$  of 4%) times a recovery factor of 0.50 (for a species of unknown status with no estimated fishery mortality; Wade and Angliss 1997), resulting in a PBR of 24 spinner dolphins per year.

## HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

### Fishery Information

Although some mortality of spinner dolphins has been observed in inshore gillnets, no estimate of annual human-caused mortality and serious injury is available. The gear types used in Hawaiian fisheries are responsible for marine mammal mortality and serious injury in other fisheries throughout U.S. waters. Gillnets are used in Hawaiian waters and appear to capture marine mammals wherever they are used, and float lines from lobster traps and longlines can be expected to occasionally entangle whales (Perrin et al. 1994).

Interactions with cetaceans have been reported for all Hawaiian pelagic fisheries, and there are records of spinner dolphins taken in inshore monofilament gillnets and net fragments in Hawaiian waters (Nitta and Henderson 1993). One spinner dolphin was observed hooked in the Hawaiian longline fishery between 1994 and 1998 in waters outside the U.S. EEZ, with approximately 4.4% of all effort (measured as the number of hooks fished) observed. This interaction rate extrapolates to a total 5-year estimate of 23 (95% CI = 1-108) spinner dolphins, or an average of 4.6 interactions per year (Kleiber 1999). The single observed spinner dolphin was reported to have been hooked in the fluke. Following the guidelines of a 1997 Serious Injury Workshop (Angliss and DeMaster 1998), this animal would not be considered seriously injured (defined under the MMPA as likely to result in mortality). Reports for other odontocetes indicate they may also become hooked in the mouth or ingest the hook, and they may occasionally become entangled in the fishing line. Interaction rates between dolphins and the NWHI bottomfish fishery have been estimated based on studies conducted in 1990-1993, indicating that an average of 2.67 dolphin interactions, most likely involving bottlenose and rough-toothed dolphins, occurred for every 1000 fish brought on board (Kobayashi and Kawamoto 1995). Fishermen claim interactions with dolphins who steal bait and catch are increasing. It is not known whether these interactions result in serious injury or mortality of dolphins, nor whether spinner dolphins are involved.



**Figure 2.** Locations of observed cetacean interactions in the Hawaiian longline fishery, 1994-98 (modified from Kleiber 1999). Dashed line is the U.S. Exclusive Economic Zone (EEZ); SL = spinner dolphin.

### Other Removals

At least 85 spinner dolphins were live-captured in Hawaiian waters from 1962 to 1981 (Shallenberger 1981). The main capture area was around Oahu.

## STATUS OF STOCK

The status of spinner dolphins in Hawaiian waters relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. A habitat issue of increasing concern is the potential effect of swim-with-dolphin programs and other tourism activities on spinner dolphins around the main Hawaiian Islands. Spinner dolphins are not listed as “threatened” or “endangered” under the Endangered Species Act (1973), nor as “depleted” under the MMPA. The Hawaiian stock is not considered a strategic stock under the 1994 amendments to the MMPA, because there are no estimates of mortality within the U.S. EEZ. However, there is no systematic monitoring of gillnet fisheries that may

take this species, and the potential effect of interactions with the Hawaiian longline fishery in international waters is not known. Insufficient information is available to determine whether the total fishery mortality and serious injury for spinner dolphins is insignificant and approaching zero mortality and serious injury rate.

## REFERENCES

- Angliss, R. P. and D. P. DeMaster. 1998. Differentiating Serious and Non-Serious Injury of Marine Mammals Taken Incidental to Commercial Fishing Operations: Report of the Serious Injury Workshop 1-2 April 1997, Silver Spring, Maryland. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Dizon, A. E., W. F. Perrin, and P. A. Akin. 1994. Stocks of dolphins (*Stenella* spp. and *Delphinus delphis*) in the eastern tropical Pacific: a phylogeographic classification. NOAA Tech. Rep. NMFS 119, 20 pp.
- Kleiber, P. 1999. Estimates of marine mammal takes in the Hawaiian longline fishery. (Unpublished). Southwest Fisheries Science Center, NMFS, 2570 Dole St, Honolulu, HI, 96822-2396.
- Kobayashi, D. R. and K. E. Kawamoto. 1995. Evaluation of shark, dolphin, and monk seal interactions with Northwestern Hawaiian Island bottomfishing activity: a comparison of two time periods and an estimate of economic impacts. Fisheries Research 23: 11-22.
- Mobley, J. R., Jr, S. S. Spitz, K. A. Forney, R. A. Grotefendt, and P. H. Forestall. 2000. Distribution and abundance of odontocete species in Hawaiian waters: preliminary results of 1993-98 aerial surveys Admin. Rep. LJ-00-14C. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 26 pp.
- Nitta, E. and J. R. Henderson. 1993. A review of interactions between Hawaii's fisheries and protected species. Mar. Fish. Rev. 55(2):83-92.
- Norris, K. S., B. Würsig, R. S. Wells, and M. Würsig. 1994. The Hawaiian Spinner Dolphin. University of California Press, 408 pp.
- Norris, K. S. and T. P. Dohl. 1980. Behavior of the Hawaiian spinner dolphin, *Stenella longirostris*. Fish. Bull. 77:821-849.
- Östman, J. S. O. 1994. Social organization and social behavior of Hawaiian spinner dolphins (*Stenella longirostris*). Ph.D. dissertation, University of California, Santa Cruz, 114 pp.
- Perrin, W. F. 1975. Variation of spotted and spinner porpoise (genus *Stenella*) in the eastern tropical Pacific and Hawaii. Bull. Scripps Inst. Oceanogr. 21, 206 pp.
- Perrin, W. F. 1990. Subspecies of *Stenella longirostris* (Mammalia: Cetacea: Delphinidae). Proc. Biol. Soc. Wash. 103:453-463.
- Perrin, W. F. and J. W. Gilpatrick, Jr. 1994. Spinner dolphin *Stenella longirostris* (Gray, 1828). In: S. H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Vol.5: The First Book of Dolphins, pp. 99-128. Academic Press, 416 pp.
- Perrin, W.F., G. P. Donovan and J. Barlow. 1994. Gillnets and Cetaceans. Rep. Int. Whal. Commn., Special Issue 15, 629 pp.
- Shallenberger, E.W. 1981. The status of Hawaiian cetaceans. Final report to U.S. Marine Mammal Commission. MMC-77/23, 79pp.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Wade, P. R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. Rep. Int. Whal. Commn. 43:477-493.