

LONG-BEAKED COMMON DOLPHIN (*Delphinus capensis*): California Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Long-beaked common dolphins have only recently been recognized as a distinct species (Heyning and Perrin 1994; Rosel et al. 1994). Along the U.S. west coast, their distribution overlaps with that of the short-beaked common dolphin, and much historical information has not distinguished between these two species. Long-beaked common dolphins are commonly found within about 50 nmi of the coast, from Baja California (including the Gulf of California) northward to about central California (Figure 1). Stranding data and sighting records indicate that the relative abundance of this species off California changes both seasonally and inter-annually, with highest densities observed during warm-water events (Heyning and Perrin 1994). Although long-beaked common dolphins are not restricted to U.S. waters, cooperative management agreements with Mexico exist only for the tuna purse seine fishery and not for other fisheries which may take this species (e.g. gillnet fisheries). Under the Marine Mammal Protection Act (MMPA), long-beaked ("Baja neritic") common dolphins involved in eastern tropical Pacific tuna fisheries are managed separately as part of the 'northern common dolphin' stock (Perrin et al. 1985), and these animals are not included in the assessment reports. For the MMPA stock assessment reports, there is a single Pacific management stock including only animals found within the U.S. Exclusive Economic Zone of California.

POPULATION SIZE

Aerial line transect surveys conducted in winter and spring of 1991 and 1992 resulted only in a combined abundance estimate of 305,694 (CV=0.34) long-beaked and short-beaked common dolphins, because species-level identification was not possible from the air (Forney et al. 1995). Based on sighting locations, the majority of these animals were probably short-beaked common dolphins. More recent, species-specific abundance estimates are available based on two summer/fall shipboard surveys that were conducted within 300 nmi of the coasts of California, Oregon and Washington in 1996 (Barlow 1997) and 2001 (NMFS, unpublished data). The distribution and abundance of long-beaked common dolphins off California appears to be variable on interannual and seasonal time scales (Heyning and Perrin 1994). As oceanographic conditions change, long-beaked common dolphins may spend time in Mexican waters, and therefore a multi-year average abundance estimate is the most appropriate for management within the U.S. waters. The 1991-96 weighted average abundance estimate for California, Oregon and Washington waters based on the three ship surveys is 43,360 (CV=0.72) long-beaked common dolphins (Barlow 2003).

Minimum Population Estimate

The log-normal 20th percentile of the weighted average abundance estimate is 25,163 long-beaked common dolphins.




Figure 1. Long-beaked common dolphin sightings based on shipboard surveys off California, Oregon, and Washington, 1991-2001 (see Appendix 2 for data sources and information on timing and location of survey effort). No *Delphinus* sightings have been made off Washington. Dashed line represents the U.S. EEZ, thick line indicates the outer boundary of all surveys combined.

Current Population Trend

Due to the historical lack of distinction between the two species of common dolphins, it is difficult to establish trends in abundance for this species. In the past, common dolphins have been shown to increase in abundance off California during the warm-water months (Dohl et al. 1986). Surveys conducted during both cold-water and warm-water conditions in 1991 and 1992 (Barlow 1995, Forney et al. 1995) resulted in overall abundance estimates (for both types of common dolphins combined) which were considerably greater than historical estimates (Dohl et al. 1986). The combined abundance estimate for the 1991-96 summer/fall surveys (Barlow 1997) is the highest and most precise to date. An ongoing decline in the abundance of 'northern common dolphins' (including both long-beaked and short-beaked common dolphins) in the eastern tropical Pacific and along the Pacific coast of Mexico (IATTC 1997) suggests a possible northward shift in the distribution of common dolphins during this period of gradual warming of the waters off California (Roemmich 1992). However, it is unclear how much of this increase reflects an increase in the abundance of the long-beaked common dolphin. Heyning and Perrin (1994) have detected changes in the proportion of short-beaked to long-beaked common dolphins stranding along the California coast, with the short-beaked common dolphin stranding more frequently prior to the 1982-83 El Niño (which increased water temperatures off California), and the long-beaked common dolphin more commonly observed for several years afterwards. Thus, it appears that both relative and absolute abundance of these species off California may change with varying oceanographic conditions.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

There are no estimates of current or maximum net productivity rates for long-beaked common dolphins.

POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) level for this stock is calculated as the minimum population size (25,163) times one half the default maximum net growth rate for cetaceans ($\frac{1}{2}$ of 4%) times a recovery factor of 0.48 (for a species of unknown status with a mortality rate $CV \geq 0.30$ and ≤ 0.60 ; Wade and Angliss 1997), resulting in a PBR of 242 long-beaked common dolphins per year.

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

A summary of recent fishery mortality and injury for long-beaked common dolphins is shown in Table 1. More detailed information on these fisheries is provided in Appendix 1. Mortality estimates for the California drift gillnet fishery are included for the five most recent years of monitoring, 1997-2001 (Cameron and Forney 1999, 2000; Carretta 2001, 2002). Because of the difficulty in distinguishing short-beaked and long-beaked common dolphins in the field, tissue samples have been collected for most of the animals observed killed. These tissue samples have enabled positive identification using genetic techniques for all except two of the common dolphins killed (NMFS, unpublished data). Based on past patterns (Barlow et al. 1997), these two animals are likely to have been a short-beaked common dolphin, and they have not been included in the mortality calculations below for long-beaked common dolphins. After the 1997 implementation of a Take Reduction Plan, which included skipper education workshops and required the use of pingers and minimum 6-fathom extenders, common dolphin entanglement rates in the drift gillnet fishery dropped considerably (Barlow and Cameron 2003). However, because of interannual variability in entanglement rates additional years of data will be required to fully evaluate the effectiveness of pingers for reducing mortality of this species in the long term. Mean annual takes in Table 1 are based on 1997-2001 data. This results in an average estimate of 11 ($CV=0.50$) long-beaked common dolphins taken annually.

Additional common dolphin mortality has been reported for set gillnets in California (Julian and Beeson 1998); however, because of a 1994 ban on gillnets in nearshore areas of Southern California, the size of this fishery decreased by about a factor of two (see Appendix 1), and the observer program was discontinued. Approximately 4% and 1.8% of the entire fishery was observed in Monterey Bay in 1999 and 2000, respectively, and no common dolphin were observed taken. Marine Mammal Authorization Permit (MMAP) fisher self-reports for 1997-2001 indicate that at least two common dolphins (type not specified) were killed between 1997 and 2001. Although these reports are considered unreliable (see Appendix 4 of Hill and DeMaster 1998) they represent a minimum mortality for this fishery.

Nine common dolphins (type not specified) stranded with evidence of fishery interaction (NMFS, Southwest Region, unpublished data) between 1997-2001. It is not known which fisheries were responsible for these deaths.

Table 1. Summary of available information on the incidental mortality and injury of long-beaked common dolphins (California Stock) and prorated unidentified common dolphins in commercial fisheries that might take this species. All observed entanglements resulted in the death of the animal. The observer program for the set gillnet fishery was discontinued during 1994 and later resumed in Monterey Bay from 1999-2000. Coefficients of variation for mortality estimates are provided in parentheses, when available. Mean annual takes are based on 1997-2001 data unless noted otherwise.

Fishery Name	Data Type	Year(s)	Percent Observer Coverage	Observed	Estimated Annual Mortality	Mean Annual Takes (CV in parentheses)	
CA/OR thresher shark/swordfish drift gillnet fishery	observer data	1997	23.0%	4	25 (0.74)	8.4 (0.50)	
		1998	20.0%	0	0		
		1999	20.0%	1	8 (0.93)		
		2000	22.9%	1	9 (0.76)		
		2001	20.4%	0	0		
CA angel shark/ halibut and other species large mesh (>3.5in) set gillnet fishery	observer data	Common dolphins, species not determined				3 ¹	0.6 (n/a)
		1997-2001	0 - 4%	0			
	MMAP self-reporting	1997	-	0	0	≥ 0.4 (n/a)	
		1998	-	2	≥ 2		
		1999	-	0	0		
		2000	-	0	0		
2001		-	0	0			
Undetermined	strandings	1997-2001	9 common dolphins (species not determined) stranded with evidence of fishery interactions			≥ 1.8 (n/a)	
Minimum total annual takes						11 (0.50)	

The set gillnet fishery was observed from 1991-94 and then only in Monterey Bay during 1999-2000, where 20-25% of the local fishery was observed. Recent mortality estimates for common dolphin in this fishery are based on kill rates observed from 1991-94 and current levels of fishing effort.

Drift gillnet fisheries for swordfish and sharks exist along the entire Pacific coast of Baja California, Mexico and may take animals from this population. Quantitative data are available only for the Mexican swordfish drift gillnet fishery, which uses vessels, gear, and operational procedures similar to those in the U.S. drift gillnet fishery, although nets may be up to 4.5 km long (Holts and Sosa-Nishizaki 1998). The fleet increased from two vessels in 1986 to 31 vessels in 1993 (Holts and Sosa-Nishizaki 1998). The total number of sets in this fishery in 1992 can be estimated from data provided by these authors to be approximately 2700, with an observed rate of marine mammal bycatch of 0.13 animals per set (10 marine mammals in 77 observed sets; Sosa-Nishizaki et al. 1993). This overall mortality rate is similar to that observed in California driftnet fisheries during 1990-95 (0.14 marine mammals per set; Julian and Beeson, 1998), but species-specific information is not available for the Mexican fisheries. Previous efforts to convert the Mexican swordfish driftnet fishery to a longline fishery have resulted in a mixed-fishery, with 20 vessels alternately using longlines or driftnets, 23 using driftnets only, 22 using longlines only, and seven with unknown gear type (Berdegúe 2002).

Other Mortality

In the eastern tropical Pacific, 'northern common dolphins' have been incidentally killed in international tuna purse seine fisheries since the late 1950's. Cooperative international management programs have dramatically reduced overall dolphin mortality in these fisheries during the last decade (Joseph 1994). Between 1997 and 2001, annual mortality of northern common dolphins (potentially including both short-beaked and long-beaked common dolphins) ranged between 9 and 261 animals, with an average of 101 (IATTC, in prep). Although it is unclear whether these animals are part of the same population as short-beaked common dolphins found off California, they are managed separately under a section of the MMPA written specifically for the management of dolphins involved in eastern tropical

Pacific tuna fisheries.

STATUS OF STOCK

The status of long-beaked common dolphins in California waters relative to OSP is not known, and there are insufficient data to evaluate potential trends in abundance of this species of common dolphin. No habitat issues are known to be of concern for this species. They are not listed as "threatened" or "endangered" under the Endangered Species Act nor as "depleted" under the MMPA. Including driftnet mortality only for years after implementation of the Take Reduction Plan (1997-98), the average annual human-caused mortality in 1997-2001 (14 animals) is estimated to be less than the PBR (226), and therefore they are not classified as a "strategic" stock under the MMPA. The average total fishery mortality and injury for long-beaked common dolphins is less than 10% of the PBR and, therefore, can be considered to be insignificant and approaching zero mortality and serious injury rate.

REFERENCES

- Barlow, J. 1995. The abundance of cetaceans in California waters. Part I: Ship surveys in summer and fall of 1991. *Fish. Bull.* 93:1-14.
- Barlow, J. 1997. Preliminary estimates of cetacean abundance off California, Oregon and Washington based on a 1996 ship survey and comparisons of passing and closing modes. Administrative Report LJ-97-11, Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 25p.
- Barlow, J. and G. A. Cameron. 2003. Field experiments show that acoustic pingers reduce marine mammal bycatch in the California drift gillnet fishery. *Marine Mammal Science* 19(2):265-283.
- Barlow, J., K. A. Forney, P. S. Hill, R. L. Brownell, Jr., J. V. Carretta, D. P. DeMaster, F. Julian, M. S. Lowry, T. Ragen, and R. R. Reeves. 1997. U.S. Pacific Marine Mammal Stock Assessments: 1996. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-248. 223p.
- Barlow, J. and T. Gerrodette. 1996. Abundance of cetaceans in California waters based on 1991 and 1993 ship surveys. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-233.
- Barlow, J. 2003. Preliminary estimates of the abundance of cetaceans along the U.S. west coast: 1991-2001. Southwest Fisheries Science Center Administrative Report LJ-03-03. Available from SWFSC, 8604 La Jolla Shores Dr., La Jolla CA 92037. 31p.
- Berdegú, J. 2002. Depredación de las especies pelágicas reservadas a la pesca deportiva y especies en peligro de extinción con uso indiscriminado de artes de pesca no selectivas (palangres, FAD's, trampas para peces y redes de agallar fijas y a la deriva) por la flota palangrera Mexicana. Fundación para la conservación de los picudos. A.C. Mazatlán, Sinaloa, 21 de septiembre.
- Cameron, G., and K. A. Forney. 1999. Estimates of cetacean mortality in the California gillnet fisheries for 1997 and 1998. Paper SC/51/O4 presented to the International Whaling Commission, 1999 (unpublished). 14pp.
- Cameron, G.A. and K.A. Forney. 2000. Preliminary estimates of cetacean mortality in California/Oregon Gillnet Fisheries for 1999. Report SC/52/O24 presented to the Scientific Committee of the International Whaling Commission, June 2000 (unpublished). 12p. [Available from Southwest Fisheries Science Center, National Marine Fisheries Service, 8604 La Jolla Shores Drive, La Jolla, CA 92037, USA.]
- Carretta, J.V. 2001. Preliminary estimates of cetacean mortality in California gillnet fisheries for 2000. Report SC/53/SM9 presented to the Scientific Committee of the International Whaling Commission, June 2001 (unpublished). 21p. [Available from Southwest Fisheries Science Center, National Marine Fisheries Service, 8604 La Jolla Shores Drive, La Jolla, CA 92037, USA.]
- Carretta, J.V. 2002. Preliminary estimates of cetacean mortality in California gillnet fisheries for 2001. Report SC/54/SM12 presented to the Scientific Committee of the International Whaling Commission, April 2002 (unpublished). 22p. [Available from Southwest Fisheries Science Center, National Marine Fisheries Service, 8604 La Jolla Shores Drive, La Jolla, CA 92037, USA.]
- Dohl, T. P., M. L. Bonnell, and R. G. Ford. 1986. Distribution and abundance on common dolphin, *Delphinus delphis*, in the Southern California Bight: A quantitative assessment based upon aerial transect data. *Fish. Bull.* 84:333-343.
- Forney, K. A., J. Barlow and J. V. Carretta. 1995. The abundance of cetaceans in California waters. Part II: Aerial surveys in winter and spring of 1991 and 1992. *Fish. Bull.* 93:15-26.
- Heyning, J. E. and W. F. Perrin. 1994. Evidence for two species of common dolphins (Genus *Delphinus*) from the eastern North Pacific. *Contr. Nat. Hist. Mus. L.A. County*, No. 442.

- Hill, P. S. and D. P. DeMaster. 1998. Alaska Marine Mammal Stock Assessments, 1998. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-97. 166 pp.
- Holts, D. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038.
- Holts, D. and O. Sosa-Nishizaki. 1998. Swordfish, *Xiphias gladius*, fisheries of the eastern North Pacific Ocean. In: I. Barrett, O. Sosa-Nishizaki and N. Bartoo (eds.). Biology and fisheries of swordfish, *Xiphias gladius*. Papers from the International Symposium on Pacific Swordfish, Ensenada Mexico, 11-14 December 1994. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 142, 276 pp.
- IATTC. 1997. Annual Report of the Inter-American Tropical Tuna Commission, 1995, La Jolla, California.
- IATTC. (in prep). Report on the International Dolphin Conservation Program. Document MOP-7-04. 19 p.
- Joseph, J. 1994. The tuna-dolphin controversy in the eastern Pacific Ocean: biological, economic and political impacts. Ocean Dev. Int. Law 25:1-30.
- Julian, F. 1997. Cetacean mortality in California gill net fisheries: Preliminary estimates for 1996. Paper SC/49/SM02 presented to the International Whaling Commission, 1997 (unpublished). 13 pp.
- Julian, F. and M. Beeson. 1998. Estimates of mammal, turtle and bird mortality for two California gillnet fisheries: 1990-1995. Fish. Bull. 96:271-284.
- NMFS, Southwest Fisheries Science Center, P.O. Box 271, La Jolla, CA 92038-027.
- NMFS, Southwest Region, 501 West Ocean Blvd, Long Beach, CA 90802-4213.
- Perrin, W. F., M. D. Scott, G. J. Walker and V. L. Cass. Review of geographical stocks of tropical dolphins (*Stenella* spp. and *Delphinus delphis*) in the eastern Pacific. NOAA Technical Report NMFS 28. Available from NMFS, Southwest Fisheries Science Center, P.O. Box 271, La Jolla, California, 92038. 28p.
- Roemmich, D. 1992. Ocean warming and sea level rise along the southwest U.S. coast. Science 257:373-375.
- Rosel, P. E., A. E. Dizon and J. E. Heyning. 1994. Population genetic analysis of two forms of the common dolphin (genus *Delphinus*) utilizing mitochondrial DNA control region sequences. Marine Biology 119:159-167.
- Sosa-Nishizaki, O., R. De la Rosa-Pacheco, R. Castro-Longoria, M. Grijalva Chon, and J. De la Rosa Velez. 1993. Estudio biologico pesquero del pez (*Xiphias gladius*) y otras especies de picudos (marlins y pez vela). Rep. Int. CICESE, CTECT9306.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.